



US79419457E (140905) ADAMENT 20 OZ ETL - 08/23/19 Colors: cmyk













Copy Area







- 2.75"-

Copy Area

**←** 0.125"








- 2.75"-

Copy Area

**←** 0.125"



**USE DIRECTIONS FOR SPECIFIC CROPS** 

ADAMENT 50 WG Fungicide provides control or suppression of several important diseases of almonds, pistachio and tree nuts. When reference is made to disease suppression, suppression can mean either erratic control from good to fair, or consistent control at a level below that obtained with the best commercial disease control products.


Almonds			
Disease Control	Rate (oz./acre)	Application Timing	
Brown Rot Blossom Blight (Monilinia laxa, M. fructicola)	6 to 8	Begin applications when conditions are favorable for	
Jacket Rot, Green Fruit Rot (Botrytis cinerea)		disease but before infection.  Apply on a 7- to 14-day spray schedule.	
Alternaria (Alternaria alternata)		Use higher rates and shorter intervals when disease pressure is severe.	
Anthracnose (Colletotrichum acutatum)			
Shot Hole (Wilsonomyces carpophilus)		Refer to University and/or extension guidelines for best application timings.	


(continued)





3.25" 3.5"











Tree Nuts: (continued)		
Disease Control	Rate (oz./acre)	Application Timing
Botryosphaeria Panicle & Shoot Blight (Botryosphaeria dothidea)	6 to 8	Begin applications when conditions are favorable for disease but before infection.
Septoria Leaf Spot (Septoria pistaciarum)		Apply on a 14- to 21-day spray schedule.
		Use higher rates and shorter intervals when disease pressure is severe.

### Restrictions:

Restrictions:

Do not apply more than 32 oz. of ADAMENT 50 WG FUNGICIDE per acre per season. Do not make more than 4 total applications of ADAMENT 50 WG FUNGICIDE per season. Do not apply ADAMENT 50 WG FUNGICIDE within 60 days of harvest or after hull split. Do not cut cover crops in treated areas for feed, or allow livestock to graze treated areas. ADAMENT 50 WG FUNGICIDE is a mixture of a Group 11 and a Group 3 fungicide. To limit the potential for the development of resistance, do not apply more than 2 sequential applications of ADAMENT 50 WG FUNGICIDE or other Group 11 -containing fungicides.

### ROTATIONAL RESTRICTIONS

Treated areas may be replanted immediately following harvest with any crop listed on this label. For crops not listed on this label, do not plant back within 120 days of last application.


# **STORAGE AND DISPOSAL**


Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross-contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary









Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Dispose of pesticide as directed. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer CropScience Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer CropScience Emergency Response telephone number is 1-800-334-7577.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.


Container Handling: Non-refillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.


Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.


Offer for recycling, if available. If not recycled, then puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.


IMPORTANT: READ BEFORE USE
Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at



3.25" 3.5"







# **BASE**

- 3.5"



ADAMENT® 50 WG Fungicide GROUP 3 11 FUNGICIDE For control of certain diseases on almonds, pistachio and tree nuts ACTIVE INGREDIENTS:

 Tebuconazole
 25.0%

 Trifloxystrobin
 25.0%

 OTHER INGREDIENTS:
 50.0%

EPA Reg. No. 264-1052

# KEEP OUT OF REACH OF CHILDREN CAUTION

For MEDICAL And TRANSPORTATION Emergencies ONLY
Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Causes moderate eye irritation. Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. 140905E

Net Contents: 1 LB. 4 OZ. (20 OZ.)

Bayer CropScience LP 800 N. Lindbergh Blvd. St. Louis, M0 63167 ADAMENT is a registered trademark of Bayer. ©2019 Bayer CropScience

3.5"

US79419457E

### **BACK BASE LABEL**

## US79424531E (140905) ADAMENT 08/23/19

color: black

3.5"

### ADAMENT® 50 WG Fungicide

FIRST AID

IF IN EYES: • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

IF SWALLOWED: • Call a poison control center or doctor immediately for treatment advice. • Do not induce or omitting unless told to do so by a poison control center or doctor. • Have person sip a glass of water if able to swallow. • Do not give anything to an unconscious person. IF ON SKIN: • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.

IF INHALED: • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice. In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.

NOTE TO PHYSICIAN: No specific antidote. Treat Symptomatically.

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not use this product until you have read the entire label.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **Pesticide Storage:** Store in a cool, dry place and in such a manner as to prevent cross-contamination with GROUP 3 11 FUNGICIDE

other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Dispose of pesticide as directed. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer CropScience Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer CropScience Emergency Response telephone number is 1-800-334-7577.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Container Handling: Non-refillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank. Indid container upiside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Offer for recycling, if available. If not recycled, then puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

US79424531E <u>140905E</u> 08/19

3.5"