Product Bulletin

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Transform® WG

EPA Reg. No. 62719-625

Special 2(ee) Recommendation[†] (For Distribution and Use in ID, MN, ND, OR, and WA)

For Control of Plant Bugs (Lygus spp) in Potatoes (Crop Groups 1C and 1D)

ATTENTION

- [†]This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.
- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Transform® WG insecticide before applying. Carefully follow all
 precautionary statements and applicable use directions except as specified below.
- Use of Transform WG according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Transform WG.

Directions for Use

Refer to product label for Transform WG for General Use Precautions, Mixing Directions, and Application Directions.

Pests and Application Rates

Pests	Transform WG (oz/acre)
Plant Bugs	1.5 – 2.25 (0.047 – 0.071 lb ai/acre)

Application Timing: Treat in accordance with local economic thresholds. Consult your Corteva Agriscience representative, cooperative extension service, certified crop advisor or state agricultural experiment station for any additional local use recommendations for your area.

Application Rate: Use a higher rate in the rate range for heavy pest populations

Restrictions:

- Preharvest Interval: Do not apply within 7 day of harvest.
- Minimum Treatment Interval: Do not make applications less than 14 days apart.
- Do not make more than four applications per crop.
- Do not make more than two consecutive applications per crop.
- Do not apply more than a total of 8.5 oz of Transform WG (0.266 lb ai of sulfoxaflor) per acre per year.

™®Trademarks of Corteva Agriscience and its affiliated companies

R396-278

Issued: 08/19/22

Replaces Label: R396-048