

Mefentrifluconazole	Group	3	Fungicide
Fluxapyroxad	Group	7	Fungicide
Pyraclostrobin	Group	11	Fungicide

Supplemental Label

Revytek™

Fungicide

For disease control in cotton, grasses, grass grown for seed, legume vegetables, non-grass forages, oilseeds, rapeseed (canola), and sugarcane

This supplemental label expires October 30, 2023, and must not be used or distributed after this date.

Active Ingredients*:

mefentrifluconazole: 2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1H-1,2,4-triazole-1-yl)propan-2-ol	11.61%
pyraclostrobin: (carbamic acid, [2-[[[1-(4-chlorophenyl)-1H-pyrazol-3-yl]oxy]methyl]phenyl]methoxy-, methyl ester).....	15.49%
fluxapyroxad: 1H-Pyrazole-4-carboxamide, 3-(difluoromethyl)-1-methyl-N-(3',4',5'-trifluoro[1,1'-biphenyl]-2-yl)-.....	7.74%
Other Ingredients:	65.16%
Total:	100.00%

* **Revytek™ fungicide** contains 1.11 lbs mefentrifluconazole, 1.48 lbs pyraclostrobin, and 0.74 lb fluxapyroxad per gallon.

EPA Reg. No. 7969-406
CAUTION/PRECAUCION

Directions For Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Revytek™ fungicide** container label, EPA Reg. No. 7969-406, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Revytek** before applying.
- Use of **Revytek** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Revytek**.

Application Instructions

Refer to the **Revytek** container label for complete application instructions and restrictions.

Crop Rotation Restrictions

Rotational Crops: Please see the following table for crop rotational restrictions.

Crop or Crop Group	Rotation Interval
Brassicas including broccoli, cauliflower and head cabbages (crop group 5-16) Bulb vegetables including bulb onion, spring onion and garlic (crop group 3-07) Cereals, including wheat, oats, barley, triticale, rye, rice and corn (crop group 15 and crop group 16) Cucurbits including cucumber, squash and melons (crop group 9) Fruiting vegetables including tomato, eggplant, peppers (crop group 8-10) Fresh herbs Grass and non-grass animal feeds including alfalfa and clover (crop group 17 and crop group 18) Leafy vegetables including lettuces, spinach and leafy cabbages (crop group 4-16) Legume vegetables including soybean (crop group 6 and crop group 7) Low-growing berries including strawberries (crop group 13-07G) Oilseeds including cotton, sunflower and canola (crop group 20) Peanut Root and tuber vegetables including carrot, potato, beets and sugar beet (crop group 1 and crop group 2) Stalk, stem and leaf petiole vegetables including celery and asparagus (crop group 22) Sugarcane Any other crop labeled for direct application of a product containing mefentrifluconazole	May be planted immediately following the last application
Other food and feed crops, not listed above	May not be planted in rotation

Detailed Use Directions

Labeled Crops

Foliar

- When conditions favor disease or disease pressure is high:
 - Use the shorter interval.
 - For a rate range, use the higher specified rate.
- **DO NOT** make more than two (2) sequential applications of **Revvytek™ fungicide** before alternating to a labeled **non-Group 3**, **non-Group 7**, or **non-Group 11** fungicide, unless otherwise specified in the following crop-specific use sections.

Use Rate Conversion

fl ozs product/A	lb mefentrifluconazole/A	lb pyraclostrobin/A	lb fluxapyroxad/A
8	0.07	0.09	0.05
12	0.10	0.14	0.07
13	0.11	0.15	0.08
15	0.13	0.17	0.09
26	0.23	0.30	0.15
30	0.26	0.35	0.17
45	0.39	0.52	0.26

State-specific Restrictions - Not for sale, distribution, or use in **Nassau and Suffolk counties in New York State**.

Cotton

Disease Controlled		Rate per Acre (fl ozs product) per application	Rate per Acre (fl ozs product) maximum per year	PHI (days)
Foliar	Alternaria leaf spot, boll rot <i>Alternaria</i> spp.	8 to 15	30	30
	Anthrachnose, boll rot <i>Glomerella</i> spp.			
	Ascochyta blight, boll rot <i>Ascochyta</i> spp.			
	Cercospora blight and leaf spot <i>Cercospora</i> spp.			
	Diplodia boll rot <i>Diplodia</i> spp.			
	Hard lock, boll rot <i>Fusarium</i> spp.			
	Phoma blight, boll rot <i>Phoma</i> spp.			
	Rust <i>Phakopsora</i> spp., <i>Puccinia</i> spp.			
	Stemphylium leaf spot <i>Stemphylium</i> spp.			
	Target spot and Corynespora leaf spot <i>Corynespora cassiicola</i>			
• Spray Interval - Apply before onset of disease and on a minimum interval of 7 days.				

Use Restrictions

- **DO NOT** apply more than 15 fl ozs (0.13 lb mefentrifluconazole, 0.17 lb pyraclostrobin, 0.087 lb fluxapyroxad) per acre per application.
- **DO NOT** make more than 2 applications at 15 fl ozs per acre per year.
- **DO NOT** apply more than 30 fl ozs (0.26 lb mefentrifluconazole, 0.34 lb pyraclostrobin, 0.17 lb fluxapyroxad) per acre per year.
- **DO NOT** apply more than a cumulative total of 0.26 lb ai/acre/year of mefentrifluconazole-containing products.
- **DO NOT** apply more than a cumulative total of 0.58 lb ai/acre/year of pyraclostrobin-containing products.
- **DO NOT** apply more than a cumulative total of 0.18 lb ai/acre/year of fluxapyroxad-containing products.

Grasses

Disease Controlled		Rate per Acre (fl ozs product) per application	Rate per Acre (fl ozs product) maximum per year	PHI (days)
Foliar	Anthracnose <i>Colletotrichum</i> spp.	8 to 13	39	0
	Leaf Spot/Blight <i>Bipolaris</i> spp., <i>Cochliobolus</i> spp., <i>Drechslera</i> spp., <i>Stagonospora</i> spp.			
	Net Blotch <i>Drechslera</i> spp., <i>Helminthosporium</i> spp.			
	Powdery Mildew <i>Erysiphe</i> spp.			
	Purple Leaf Spot <i>Stagonospora</i> spp.			
	Rust <i>Puccinia</i> spp., <i>Uromyces dactylidis</i>			
	Septoria Leaf Spot <i>Septoria</i> spp.			
• Spray Interval - Apply at 14-day to 21-day intervals.				

Use Restrictions

- **DO NOT** apply more than 13 fl ozs (0.13 lb mefentrifluconazole, 0.15 lb pyraclostrobin, 0.075 lb fluxapyroxad) per acre per application.
- **DO NOT** make more than 3 applications at 15 fl ozs per acre per year.
- **DO NOT** apply more than 39 fl ozs (0.33 lb mefentrifluconazole, 0.45 lb pyraclostrobin, 0.22 lb fluxapyroxad) per acre per year.
- **DO NOT** apply more than a cumulative total of 0.39 lb ai/acre/year of mefentrifluconazole-containing products.
- **DO NOT** apply more than a cumulative total of 0.45 lb ai/acre/year of pyraclostrobin-containing products.
- **DO NOT** apply more than a cumulative total of 0.22 lb ai/acre/year of fluxapyroxad-containing products.

Detailed Grasses (crop group 17) Crop List - any grass, Gramineae family (either green or cured) that will be fed to or grazed by livestock, all pasture and range grasses and grasses grown for hay or silage, except sugarcane and those included in the cereal grains group.

Grass Grown for Seed

Disease Controlled		Rate per Acre (fl ozs product) per application	Rate per Acre (fl ozs product) maximum per year	PHI (days)
Foliar	Powdery mildew <i>Erysiphe</i> spp.	8 to 13	26	0
	Rust <i>Puccinia</i> spp.			
	• Spray Interval - Apply at 14-day to 21-day intervals.			
Use Restrictions				
• DO NOT apply more than 13 fl ozs (0.13 lb mefentrifluconazole, 0.15 lb pyraclostrobin, 0.075 lb fluxapyroxad) per acre per application. • DO NOT make more than 2 applications at 15 fl ozs per acre per year. • DO NOT apply more than 26 fl ozs (0.22 lb mefentrifluconazole, 0.30 lb pyraclostrobin, 0.15 lb fluxapyroxad) per acre per year. • DO NOT apply more than a cumulative total of 0.26 lb ai/acre/year of mefentrifluconazole-containing products. • DO NOT apply more than a cumulative total of 0.39 lb ai/acre/year of pyraclostrobin-containing products. • DO NOT apply more than a cumulative total of 0.15 lb ai/acre/year of fluxapyroxad-containing products.				
Detailed Grass Grown for Seed (crop group 17) Crop List - any grass, Gramineae family (either green or cured) that will be fed to or grazed by livestock, all pasture and range grasses and grasses grown for hay or silage, except sugarcane and those included in the cereal grains group.				

Legume Vegetables (crop group 6)
(except soybean and edamame)

Disease Controlled		Rate per Acre (fl ozs product) per application	Rate per Acre (fl ozs product) maximum per year	PHI (days)
Foliar	Alternaria leaf and pod spot <i>Alternaria</i> spp.	8 to 13	26	21
	Anthracnose <i>Colletotrichum</i> spp.			
	Ascochyta blight <i>Ascochyta</i> spp., <i>Phoma exigua</i>			
	Asian soybean rust <i>Phakopsora pachyrhizi</i>			
	Botrytis gray mold <i>Botrytis cinerea</i>			
	Cercospora leaf spot <i>Cercospora</i> spp.			
	Mycosphaerella blight <i>Mycosphaerella</i> spp.			
	Powdery mildew <i>Erysiphe polygoni</i>			
	Rust <i>Uromyces appendiculatus</i>			
	* Downy mildew <i>Peronospora</i> spp., <i>Phytophthora</i> spp.			
• Spray Interval - Apply at 7-day to 14-day intervals. • Additives and Adjuvants - EXCEPT FOR - Chickpea, lentil, and field pea - DO NOT use with emulsifiable concentrate (EC) or solvent-based formulation products, crop oil concentrate (COC), methylated seed oil (MSO), organosilicone (OS), or MSO/OS blend adjuvants.				
* Suppression only				
Use Restrictions • DO NOT apply more than 13 fl ozs (0.11 lb mefentrifluconazole, 0.15 lb pyraclostrobin, 0.08 lb fluxapyroxad) per acre per application. • DO NOT make more than 2 applications per acre per year. • DO NOT apply more than 26 fl ozs (0.23 lb mefentrifluconazole, 0.30 lb pyraclostrobin, 0.15 lb fluxapyroxad) per acre per year. • DO NOT apply more than a cumulative total of 0.39 lb ai/acre/year of mefentrifluconazole-containing products. • DO NOT apply more than a cumulative total of 0.29 lb ai/acre/year of pyraclostrobin-containing products. • DO NOT apply more than a cumulative total of 0.18 lb ai/acre/year (0.36 lb for dried beans) of fluxapyroxad- containing products. • Bean forage, bean hay, pea vines, and pea hay may be fed no sooner than 21 days after last application.				

(continued)

Legume Vegetables (crop group 6)

(except soybean and edamame) *(continued)*

Detailed Legume Vegetables (crop group 6) Crop List (except soybean and edamame; see **Soybean and Edamame** detailed use directions)

- broad bean (fava bean), chickpea (garbanzo bean), guar, jackbean, lablab bean, lentil, pigeon pea, sword bean.
- ***Lupinus spp.***: grain lupin, sweet lupin, white lupin, white sweet lupin.
- ***Phaseolus spp.***: field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean.
- ***Vigna spp.***: adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean.
- ***Pisum spp.***: dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea.

Non-grass Forages

Disease Controlled		Rate per Acre (fl ozs product) per application	Rate per Acre (fl ozs product) maximum per year	PHI (days)
Foliar	Anthracnose <i>Colletotrichum trifolii</i>	8 to 13	39	14
	Common leaf spot <i>Pseudopeziza medicaginis</i>			
	Leaf spot <i>Leptosphaerulina briosiani</i>			
	Powdery mildew <i>Erysiphe</i> spp.			
	Rhizoctonia blight/black patch <i>Rhizoctonia</i> spp.			
	Rust <i>Phakopsora</i> spp., <i>Puccinia</i> spp., <i>Uromyces</i> spp.			
	Spring black stem and leaf spot <i>Ascochyta medicaginicola</i>			
	Stagonospora leaf spot <i>Stagonospora meliloti</i>			
	Stemphylium leaf spot <i>Stemphylium</i> spp.			
	Summer black stem and leaf spot <i>Cercospora medicaginis</i>			
	Yellow leaf blotch <i>Leptotrichia medicaginis</i>			
	• Spray Interval - Apply at 14-day to 21-day intervals.			

Use Restrictions

- **DO NOT** apply more than 13 fl ozs (0.13 lb mefentrifluconazole, 0.15 lb pyraclostrobin, 0.075 lb fluxapyroxad) per acre per application.
- **DO NOT** make more than 3 applications at 15 fl ozs per acre per year.
- **DO NOT** apply more than 39 fl ozs (0.33 lb mefentrifluconazole, 0.45 lb pyraclostrobin, 0.22 lb fluxapyroxad) per acre per year.
- **DO NOT** apply more than a cumulative total of 0.39 lb ai/acre/year of mefentrifluconazole-containing products.
- **DO NOT** apply more than a cumulative total of 0.45 lb ai/acre/year of pyraclostrobin-containing products.
- **DO NOT** apply more than a cumulative total of 0.22 lb ai/acre/year of fluxapyroxad-containing products.
- **DO NOT** apply within 14 days of grazing or harvest for forage or hay.
- **DO NOT** use on rangeland.

Detailed Non-grass Forages (crop group 18) Crop List - alfalfa; bean, velvet; clover (*Trifolium* spp., *Melilotus* spp.); kudzu; lespedeza; lupin; sainfoin; trefoil; vetch; vetch, crown; vetch, milk.

Oilseeds

Disease Controlled		Rate per Acre (fl ozs product) per application	Rate per Acre (fl ozs product) maximum per year	PHI (days)
Foliar	Alternaria leaf spot <i>Alternaria</i> spp.	8 to 15	30	21
	Blackleg <i>Leptosphaeria maculans</i>			
	Blackspot <i>Alternaria</i> spp.			
	Cercospora leaf spot <i>Cercospora</i> spp.			
	Pasmo <i>Septoria linicola</i>			
	Powdery mildew <i>Erysiphe</i> spp.			
	Rust <i>Puccinia</i> spp., <i>Uromyces</i> spp.			
	Septoria leaf spot <i>Septoria</i> spp.			
	White rust <i>Albugo tragopogonis</i>			
	• Spray Interval - Apply at 7-day to 14-day intervals. - Blackleg: Apply at the 2-leaf to 4-leaf stage. - Blackspot: Apply at early pod development.			

Use Restrictions

- **DO NOT** apply more than 15 fl ozs (0.13 lb mefentrifluconazole, 0.17 lb pyraclostrobin, 0.087 lb fluxapyroxad) per acre per application.
- **DO NOT** make more than 2 applications at 15 fl ozs per acre per year.
- **DO NOT** apply more than 30 fl ozs (0.25 lb mefentrifluconazole, 0.34 lb pyraclostrobin, 0.174 lb fluxapyroxad) per acre per year.
- **DO NOT** apply more than a cumulative total of 0.26 lb ai/acre/year of mefentrifluconazole-containing products.
- **DO NOT** apply more than a cumulative total of 0.35 lb ai/acre/year of pyraclostrobin-containing products.
- **DO NOT** apply more than a cumulative total of 0.17 lb ai/acre/year of fluxapyroxad-containing products.

Detailed Oilseeds (crop group 20) Crop List - borage; calendula; castor oil plant; Chinese tallowtree; cottonseed; crambe; cuphea; echium; euphorbia; evening primrose; flax seed; gold of pleasure; hare's ear mustard; jojoba; lesquerella; lunaria; meadowfoam; milkweed; mustard seed; niger seed; oil radish; poppy seed; rapeseed; rose hip; safflower; sesame; stokes aster; sunflower; sweet rocket; tallowwood; tea oil plant; veronica; cultivars, varieties, and/or hybrids of these.

Rapeseed (canola) (subgroup 20A)

Disease Controlled		Rate per Acre (fl ozs product) per application	Rate per Acre (fl ozs product) maximum per year	PHI (days)
Foliar	Blackleg <i>Leptosphaeria maculans</i>	8 to 15	30	21
	Blackspot <i>Alternaria</i> spp.			
	* White mold/Sclerotinia stem rot <i>Sclerotinia sclerotiorum</i>	15		
• Spray Interval - Apply at 14-day intervals. - Blackleg: Apply at the 2-leaf to 4-leaf stage. - Blackspot: Apply at early pod development. - Sclerotinia: Apply before disease develops or at 20% to 50% flowering. A second application may be made 14-days later if weather conditions are favorable for disease.				

* Suppression only

Use Restrictions

- **DO NOT** apply more than 15 fl ozs (0.13 lb mefentrifluconazole, 0.17 lb pyraclostrobin, 0.09 lb fluxapyroxad) per acre per application.
- **DO NOT** make more than 2 applications per acre per year.
- **DO NOT** apply more than 30 fl ozs (0.26 lb mefentrifluconazole, 0.35 lb pyraclostrobin, 0.17 lb fluxapyroxad) per acre per year.
- **DO NOT** apply more than a cumulative total of 0.26 lb ai/acre/year of mefentrifluconazole-containing products.
- **DO NOT** apply more than a cumulative total of 0.39 lb ai/acre/year of pyraclostrobin-containing products.
- **DO NOT** apply more than a cumulative total of 0.18 lb ai/acre/year of fluxapyroxad-containing products.

Detailed Rapeseed (subgroup 20A) Crop List - borage; crambe; cuphea; echium; flax seed; gold of pleasure; hare's ear mustard; lesquerella; lunaria; meadowfoam; milkweed; mustard seed; oil radish; poppy seed; rapeseed, canola; sesame; sweet rocket; cultivars, varieties, and/or hybrids of these.

Sugarcane

Disease Controlled		Rate per Acre (fl ozs product) per application	Rate per Acre (fl ozs product) maximum per year	PHI (days)
Infurrow	Rhizoctonia seed and seedling rot <i>Rhizoctonia solani</i> Sugarcane pineapple disease <i>Ceratocystis paradoxa</i> * Fusarium seed rot, Seedling blight <i>Fusarium</i> spp.	11 to 15	30	14
	Brown rust <i>Puccinia melanocephala</i> Orange rust <i>Puccinia kuehnii</i>	8 to 15		
Foliar	• Spray Interval - Apply at 14-day to 28-day intervals.			
* Suppression only				
Use Restrictions • DO NOT apply more than 15 fl ozs (0.13 lb mefentrifluconazole, 0.17 lb pyraclostrobin, 0.087 lb fluxapyroxad) per acre per application. • DO NOT make more than 2 applications at 15 fl ozs per acre per year. • DO NOT apply more than 30 fl ozs (0.25 lb mefentrifluconazole, 0.35 lb pyraclostrobin, 0.17 lb fluxapyroxad) per acre per year. • DO NOT apply more than a cumulative total of 0.25 lb ai/acre/year of mefentrifluconazole-containing products. • DO NOT apply more than a cumulative total of 0.78 lb ai/acre/year of pyraclostrobin-containing products. • DO NOT apply more than a cumulative total of 0.27 lb ai/acre/year of fluxapyroxad-containing products. • DO NOT apply this product to sugarcane through any type of irrigation system. • DO NOT feed treated sugarcane commodities to livestock. • Aerial Application - DO NOT use less than 5 gallons of spray solution per acre. Revytek™ fungicide can be applied by ground or air.				

SUGARCANE ONLY Infurrow and Soil-directed Banded Rates

Rate per 1000 Row Feet (fl ozs product)	Product Rate (fl ozs/A) Row Width (feet)	
	5 feet	6 feet
1.26	11.0	
1.38	12.0	
1.49	13.0	
1.52		11.0
1.61	14.0	
1.65		12.0
1.72	15.0	
1.79		13.0
1.93		14.0
2.07		15.0

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

Revytek is a trademark of BASF.

© 2020 BASF Corporation
All rights reserved.

007969-00406.20201119.**NVA 2020-04-562-0230**
Based on: NVA 2020-04-562-0148

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709



We create chemistry