



PENOXsulAM	GROUP	2	HERBICIDE
OXYfluORFEN	GROUP	14	HERBICIDE

# FIFRA Section 24(c) Special Local Need (SLN) Label

Corteva Agriscience LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

**Cleantraxx™**  
EPA Reg. No. 62719-702

**For Control of Herbaceous Broadleaf Weeds, Grasses and Woody  
Plant Seedlings in Actively Managed Conifer Forests**

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF IDAHO

Expiration Date: This label is valid until December 31, 2026, or until otherwise amended,  
withdrawn, canceled, or suspended

24(c) Special Local Need Registration EPA SLN No. ID-220011

**ATTENTION**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label and the federal label for this product must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, and precautions on this supplemental label and the main EPA-registered label

Active Ingredient:  
 penoxsulam: 2-(2,2-difluoroethoxy)-N-(5,8-dimethoxy[1,2,4] triazolo[1,5c]pyrimidin-2-yl)-6-(trifluoromethyl)benzenesulfonamide.....0.85%  
 oxyfluorfen: 2-chloro-1-(3-ethoxyl-4-nitrophenoxy)-4-(trifluoromethyl) benzene .....40.31%  
 Other Ingredients .....58.84%  
 Total .....100.00%

Contains 0.083 lb penoxsulam and 3.93 lb oxyfluorfen active ingredient per gallon

## Keep Out of Reach of Children

# CAUTION

### Environmental Hazards

This product is toxic to aquatic invertebrates and wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. See Directions for Use for additional restrictions. Do not contaminate water when disposing of equipment wash water or rinsate.

### GROUNDWATER ADVISORY STATEMENT

This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

### NON-TARGET ORGANISM ADVISORY STATEMENT

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks
- Safety glasses

## Directions for Use

### Product Information

Cleantraxx™ herbicide is a selective herbicide for preemergence and postemergence residual weed control of certain broadleaf and grass weeds in forestry applications. Apply Cleantraxx from early fall to late winter or in early spring, prior to germination of targeted weeds, as per directions below. The best weed control is obtained by application to weeds either preemergence or early postemergence when weeds are small and actively growing. Any cultural practices that disturb or redistribute surface soil following treatment with Cleantraxx will reduce weed control effectiveness.

## Forestry Use Sites

Cleantraxx herbicide may be applied to forestry sites for site preparation prior to planting or for conifer release as broadcast over the top of trees. Not for use in Christmas tree plantations or in ornamental plantings. Applications may be made as aerial or ground broadcast applications, spot or directed spray applications for preemergent and early postemergent control of herbaceous broadleaf weeds, grasses and preemergent control of germinating seeds of sensitive woody plants.

Tank-mix combinations with labeled rates of other herbicides can be made provided: (1) the tank mix product is labeled for the timing and method of application for the use site to be treated and (2) mixing is not prohibited by the label of the registered tank mixed products. Use as directed in the Directions of Use section of all herbicides in the mix.

## Application Directions

- Preemergence control: Cleantraxx controls susceptible weeds germinating from seed. Cleantraxx is stable on the soil surface for up to 21 days, but must be incorporated by moisture to provide effective preemergence control of susceptible weeds. A single rainfall of 0.5 inches or more, within 21 days after application, is necessary to activate Cleantraxx.
- Postemergence control: The addition of an adjuvant at 1 quart per 100 gallons of spray volume (0.25% v/v) of a crop oil concentrate, methylated seed oil or 80% active (minimum) nonionic surfactant is required for effective postemergence control of susceptible emerged weeds. For applications over-the-top of conifers, use only adjuvants/surfactants for which the desired conifers are known to be tolerant.
- When rate ranges are given, use a lower rate in the rate range on coarse textured soils low in organic matter, lighter weed infestations, and for reduced lengths of residual control. Use a higher rate in the rate range on medium to fine textured soils, soils containing higher organic matter, heavy weed infestations, or for extended residual preemergence weed control.
- Repeat treatments may be made, but the total amount of Cleantraxx applied must not exceed 4.5 pints per broadcast acre per year.
- For broad spectrum applications use Cleantraxx tank mixed with an appropriate herbicide such as Opensight, Accord XRT II, Rodeo, Garlon 4 Ultra, Garlon XRT, Transline, Vastlan or other herbicides registered for use on these sites and for these application methods. When tank mixing, read and carefully follow all applicable use directions, precautions and limitations on the respective product labels.

## Application Precautions

- For the best weed control, apply Cleantraxx as a preemergence application prior to weed emergence during the winter dormant period.
- Postemergent applications made when weeds are less than 4 inches in height are most effective.
- For best control from broadcast and spot or directed spray applications of Cleantraxx, recommended spray volumes are 10 to 25 gallons per acre by air or 10 to 50 gallons per acre by ground. For ground applications lower spray volumes (5 gallons per acre) may be used, however lower spray volumes may reduce weed control or conifer tolerance so users should gain local experience to see if results are acceptable.
- Any cultural practices, cultivation, or disturbance of the soil surface after application will decrease the weed control provided by Cleantraxx.
- To prevent misapplication, spot treatments and directed spray treatments should be applied using calibrated application methods.
- To prevent misapplication, spot or directed treatments should be applied with a boom, boomless spray systems, hand-held or backpack sprayers.

## Site Preparation Applications

Use Cleantraxx at 3 to 4.5 pints per acre in site preparation applications prior to planting to control herbaceous weeds and sensitive woody brush seedlings. Split applications of 2.25 pints per acre may be made in the fall and following spring for control of sensitive weeds such as woodland groundsel. Tank mixes with other herbicides labeled for use in forestry or conifer plantations can be made to broaden the spectrum of control of various weedy plants. Conifers may be planted into areas treated with Cleantraxx, but disturbance of the treated soil can break the herbicide barrier and decrease weed control. Also, planters should avoid placing treated soil into the rooting zone of the tree seedlings. When tank mixtures of herbicides are used, labels for all products in the mixture must be followed and the longest specified waiting period before planting observed.

## Woody Plant Seedling Control

Apply Cleantraxx at 3 to 4.5 pints per acre for preemergent control of sensitive woody brush seedlings, such as squaw carpet, deerbrush and manzanita.

## Conifer Release Applications

Cleantraxx at 3 to 4.5 pints per acre may be applied broadcast by aerial or ground applications or as a spot or directed spray application over-the-top of conifers such as Douglas-fir, Ponderosa pine, sugar pine, incense cedar, white fir and grand fir for preemergent and early postemergence control of weeds. Split applications of 2.25 pints per acre may be made in the fall and following spring for control of sensitive weeds such as woodland groundsel. In general, applications made before bud break or after terminals harden off have the least potential for injury to crop trees. Applications made during active growth stages may injure desirable trees. Users should determine through field trials whether applications of Cleantraxx give acceptable tree tolerance under local conditions for tree species not listed above.

## Use Restrictions

- Do not apply more than 4.5 pints of Cleantraxx per acre per year.
- Do not apply Cleantraxx when weather conditions favor drift. Avoid drift to all non-target crops and areas.
- Do not treat inside banks or bottoms of irrigation ditches, either dry or containing water, or other channels that carry water that may be used for irrigation or domestic purposes. Do not contaminate water used for irrigation or domestic purposes.
- Do not graze or harvest plants from areas treated with Cleantraxx for feed or forage for domestic livestock.
- Do not apply to frozen soil or snow-covered soil.

### **Conditions and Risks of Use for Special Local Need**

**USE OF CLEANTRAXX HERBICIDE (THE "PRODUCT") ON ACTIVELY MANAGED CONIFER FORESTS (THE "CROP") FOR THIS SPECIAL LOCAL NEED MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS AS FURTHER DISCUSSED BELOW. READ AND UNDERSTAND THESE CONDITIONS AND RISKS OF USE FOR SPECIAL LOCAL NEED BEFORE USING THE PRODUCT ON THE CROP.**

**This Product is available for use in the manner described in this Special Local Need (SLN) label on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury, Crop yield reduction or Crop loss. The decision to use this Product in the manner described in this SLN Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the potential risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls and (iv) any other relevant factors.**

**By purchasing the Product for use, or using the Product, in the manner described in this SLN Labeling, you acknowledge and accept that to the extent permitted by law:**

- (1) you assume all risk of Crop injury, Crop yield reduction and Crop loss;**
- (2) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this SLN labeling;**
- (3) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaim all warranties, expressed or implied, including any implied warranty of merchantability;**
- (4) Corteva Agriscience disclaims all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to Crop injury, Crop yield reduction and/or Crop loss;**
- (5) these Conditions and Risks of Use for Special Local Need supersede any contrary representations or recommendations by Corteva Agriscience, or its respective agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.**

**If these Conditions and Risks of Use for Special Local Need are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container.**

**These Conditions and Risks of Use for Special Local Need are required by Corteva Agriscience and not specified by the US EPA or the State of Idaho.**

**24(c) Registrant:**

**Corteva Agriscience LLC  
9330 Zionsville Road  
Indianapolis, IN 46268**

<sup>TM</sup>® Trademarks of Corteva Agriscience and its affiliated companies

R419-009  
Accepted: 06/17/22  
Initial Printing