

**RESTRICTED USE PESTICIDE
DUE TO ACUTE TOXICITY**

TO BE USED BY CERTIFIED APPLICATORS ONLY – NOT TO BE USED BY UNCERTIFIED PERSONS WORKING UNDER THE SUPERVISION OF A CERTIFIED APPLICATOR, EXCEPT THAT UNCERTIFIED PERSONS MAY TRANSPORT CONTAINERS AS PROVIDED UNDER DIRECTIONS FOR USE.

syngenta.

HERA

Section 24(c) Special Local Need Label

PARAQUAT DICHLORIDE

GROUP

22

HERBICIDE

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF ARIZONA

**Gramoxone® SL 3.0
EPA Reg. No. 100-1652
EPA SLN No. AZ-230101**

For Use on Guayule to Control Weeds and for Desiccation to Facilitate Harvest in the Following Counties Only – Cochise, Gila, Graham, Greenlee, La Paz, Maricopa, Mojave, Pima, Pinal, Yuma

This label for Gramoxone SL 3.0 expires and must not be distributed or used in accordance with this SLN registration after December 31, 2027

KEEP OUT OF REACH OF CHILDREN / MANTENER FUERA DEL ALCANCE DE LOS NIÑOS.



DANGER / POISON

PELIGRO / VENENO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)
See additional precautionary statements and directions for use inside booklet

DIRECTIONS FOR USE

- It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.

- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

Specific Crop Use Directions

Guayule (Pre and Postemergence Directed Spray and Harvest Aid)

Crop	Use Pattern	Rate Per Treated Acre	Minimum Spray Volume gal/acre	Preharvest Interval (Days)
Guayule	Pre- and Postemergence - Directed Spray	1.3 – 2.7 pt	Ground: 20 gal	See precautions below
Desiccation of Guayule to facilitate harvest.	Harvest-Aid	1.3 – 2.7 pt	Ground: 20 gal Air: 10 gal	

ALWAYS ADD ONE OF THE FOLLOWING:

Add a nonionic surfactant containing 80% or more surface-active agent at 0.25% v/v (2 pint per 100 gallons) of the finished spray volume for ground applications. For aerial applications, add a nonionic surfactant at 0.25% v/v (2 pints per 100 gallons) of the finished spray volume.

Specific Use Restrictions

DO NOT allow livestock to graze treated fields. DO NOT use treated crops for livestock feed. Crop residues are not to be fed to livestock or baled for such use. If land used for guayule production changes ownership or use pattern, measures must be taken to ensure no crop residues are used for livestock feed. This information must be passed along to anyone using this crop.

Precautions, Restrictions, and Comments

- DO NOT harvest until at least 4 days after application.
- Maximum number of applications per 12 month period is 5.
- DO NOT apply this product in a manner that allows spray to drift from the application target site and/or cause harm to humans, animals, or other non-target sites.
- DO NOT apply by ground equipment within 25 ft or by air within 75 ft of lakes; reservoirs; rivers; permanent streams, marshes or natural ponds; estuaries; and commercial fish ponds.
- Make aerial or ground applications when the wind velocity favors on target product deposition, (approximately 3-10 mph). Do not apply when wind velocity exceeds 10 mph for ground applications or 7 mph for aerial applications.
- For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wingtip or rotor vortices. Boom length must not exceed 75% of wingspan or rotor diameter.
- For directed spray applications, do not allow spray to contact green stems or foliage as injury may result.

- Use flat fan nozzles for the most effective application.
 - Follow recommended spray volumes listed. These are minimum volumes only, and spray volumes should be increased as necessary to obtain complete coverage of the target weed or plant without runoff from the foliage.
 - DO NOT apply this product through any type of irrigation system.
 - DO NOT enter treated field within 12 hours of application.
 - The user must refer to the federal approved Gramoxone SL 3.0 label and read and follow direction for use, restrictions, and precautions.
-

©2022 Syngenta

Gramoxone® and the Syngenta logo are trademarks of a Syngenta Group Company

24(c) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300

Label Code: AZ1652047AA1222