

---

## SUPPLEMENTAL LABELING

---

**Syngenta Crop Protection, LLC**  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300

**SCP 828A-S7 0222**

CYPRODINIL	GROUP	<b>9</b>	FUNGICIDE
------------	-------	----------	-----------

**Vangard® WG**

### Fungicide

**This supplemental label expires on 02/04/2025 and must not be used or distributed after this date.**

Active Ingredient:	
Cyprodinil: 4-cyclopropyl-6-methyl-N-phenyl-pyrimidinamine*	75.0%
Other Ingredients:	25.0%
Total:	100.0%

\*CAS No. 121552-61-2

Vangard WG is a water-dispersible granule containing 75% cyprodinil.

**KEEP OUT OF REACH OF CHILDREN.**

## CAUTION

**EPA Reg. No. 100-828**

**All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Vangard WG as permitted according to this Supplemental Labeling, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.**

**syngenta**

## DIRECTIONS FOR USE

Crop	Disease	Product Rate oz/Acre	Use Directions
<b>Tropical and Subtropical, Small fruit, inedible peel subgroup 24A**</b>  Lychee Longan Spanish lime  And cultivars and/or hybrids of these	Botrytis fruit rot ( <i>Botrytis</i> spp.)  Alternaria fruit rot ( <i>Alternaria</i> spp.)  Anthracnose ( <i>Colletotrichum</i> spp.)	5.5 – 7*	Make the first application during early bloom and repeat on 7- to 10-day intervals if conditions remain favorable for disease development.  Resistance Management: After 2 applications of Vangard WG, alternate with another fungicide with a different mode of action for 2 applications.

\*5.5 oz product contains 0.26 lb ai cyprodinil/A

\*7 oz product contains 0.33 lb ai cyprodinil/A

**\*\*Complete List of Subgroup 24A Crops:** Aisen; bael fruit; Burmese grape; cat's-eyes; inga; longan; Lychee; madras-thorn; manduro; matisia; mesquite; mongongo, fruit; pawpaw, small-flower; satinleaf; Sierra Leone-tamarind; Spanish lime; velvet tamarind; wampi; white star apple; cultivars, varieties, and hybrids of these commodities.

**Application Instructions:** Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 20 gallons/A spray volume by air.

### Specific Use Restrictions

- 1) Maximum Single Application Rate: Do not exceed the maximum rate listed in the table.
- 2) Make no more than two applications by air.
- 3) Minimum Application Interval: 7 days
- 4) Do not apply more than 28 oz/A of Vangard WG (1.3 lb ai/A of cyprodinil) per year.
- 5) Do not apply more than a maximum total of 4 applications (air plus ground) per year.
- 6) May be applied on the day of harvest (0-day PHI).

Vangard® and the Syngenta logo are trademarks of a Syngenta Group Company

©2022 Syngenta