

**RESTRICTED USE PESTICIDE**

**DUE TO HIGH ACUTE TOXICITY TO HUMANS**

For retail sale and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. Direct supervision for this product requires the certified applicator to review federal and supplemental label instructions with all personnel prior to application, mixing, loading or repair or cleaning of application equipment.



**2(ee) Recommendation**

Use information in accordance with Section 2(ee) of FIFRA

**DUPONT™ LANNATE® LV INSECTICIDE**

EPA Reg. No. 352-384

**FOR USE ON CABBAGE FOR THE CONTROL  
OF SLUGS IN THE STATE OF NEW YORK**

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in any way that will contact workers or other persons; either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe; consult the agency responsible for pesticide regulation. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

DuPont™ LANNATE® LV can be applied at 3 pts/acre when slug populations reach locally determined economic thresholds. Consult the cooperative extension service, professional consultants or other qualified authorities to determine appropriate threshold levels for treatment in your area.

**APPLICATION**

Slugs primarily feed in the early evening, at night or in cool, early morning hours. Slugs are most active when weather conditions are moist and cool. It is recommended to make applications in the early evening through early morning when slugs are active and feeding.

LANNATE® LV should be applied at 5 to 7 day intervals to maintain control. Use sufficient water to obtain thorough, uniform coverage. Since LANNATE® LV is a fast acting contact insecticide, best results follow direct spraying of the target insect.

**NOTE:**

Add a wetting agent to improve coverage.

Do not apply LANNATE® LV within 1 day of harvest.

Do not apply more than 7.2 pounds of methomyl (24 pints of LANNATE® LV) per acre per growing season.

Do not make more than a total of 15 applications of LANNATE® LV per crop; minimum interval between treatments is 2 days.

(Replaces R-1266 041712 04-23-12)

**IMPORTANT NOTICE**

**BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.**

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

**Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically directed by DuPont. User assumes all risks associated with such non-directed use. If terms are not acceptable, return the unopened package at once to Seller for a full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.**

Copies of DuPont product labels are available by calling 1-888-6-DUPONT  
or at <http://cropprotection.dupont.com/>