

SUPPLEMENTAL LABELING

**RESTRICTED USE PESTICIDE
DUE TO ACUTE TOXICITY**
TO BE USED BY CERTIFIED APPLICATORS ONLY – NOT TO BE USED BY UNCERTIFIED PERSONS
WORKING UNDER THE SUPERVISION OF A CERTIFIED APPLICATOR, EXCEPT THAT UNCERTIFIED
PERSONS MAY TRANSPORT CONTAINERS AS PROVIDED UNDER DIRECTIONS FOR USE.

Syngenta Crop Protection, LLC

P. O. Box 18300
Greensboro, North Carolina, 27419-8300
SCP 1652A-S20A 0822

| | | | |
|---------------------|-------|-----------|-----------|
| PARAQUAT DICHLORIDE | GROUP | 22 | HERBICIDE |
|---------------------|-------|-----------|-----------|

**Gramoxone® SL 3.0
HERBICIDE**

EPA Reg. No. 100-1652

A Weed, Grass, and Harvest Aid Desiccant/Defoliant Herbicide

Active Ingredients:

| | |
|----------------------------|---------------|
| Paraquat dichloride* | 43.9% |
| Other Ingredients: | 56.1% |
| Total: | 100.0% |

*CAS No. 4685-14-7

Gramoxone SL 3.0 is formulated as a liquid containing 3.0 pounds paraquat cation per gallon as 4.134 pounds paraquat dichloride per gallon.

Gramoxone SL 3.0 contains alerting agent (odor), emetic, and dye.

KEEP OUT OF REACH OF CHILDREN / MANTENER FUERA DEL ALCANCE DE LOS NIÑOS.



**DANGER / POISON
PELIGRO / VENENO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)
See additional precautionary statements and directions for use inside booklet.

All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Gramoxone SL 3.0 as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



**DIRECTIONS FOR USE
RESTRICTED USE PESTICIDE**

Wheat, ID Only

| Crops (Including all cultivars, varieties, and/or hybrids of these) | | | |
|--|--|---|--|
| Wheat | | | |
| Target Weed | Rate pt/A | Application Timing | Use Directions |
| <p>Suppression:</p> <p>Downy brome</p> <p>Rye</p> | <p><u>Downy Brome</u></p> <p style="text-align: center;">0.5</p> <p><u>Rye</u></p> <p style="text-align: center;">0.33 – 0.5</p> | <p>Postemergence (Directed Spray)</p> | <p>For use in Idaho Only as a salvage spray to be used only when wheat production is in danger of being severely reduced by rye or downy brome (cheatgrass) competition and where reseeding is the only other alternative.</p> <p>Apply as a postemergence broadcast either by ground application in a minimum of 20 gal/A spray volume or by air in a minimum of 5 gal/A spray volume.</p> <p>Apply after wheat plants have developed five tillers or six inches in height in the spring and prior to emergence of the head from the boot.</p> <p>Add a nonionic surfactant containing at least 50% surface active ingredient at 1-2 pt per 100 gal of dilute spray.</p> <p>See Section 4.4.4 of the Gramoxone SL 3.0 label for adjuvant recommendations.</p> |
| <p>Tank Mix Options:</p> <ul style="list-style-type: none"> Gramoxone SL 3.0 can be tank mixed with other herbicides labeled for use on wheat applied by the same methods and at the same timings. Refer to Section 4.4.2 for tank-mix precautions in the Gramoxone SL 3.0 label. | | | |
| <p>Resistance Management:</p> <ul style="list-style-type: none"> Refer to Section 3.1. | | | |
| <p>Precaution:</p> <ul style="list-style-type: none"> Failure to comply with these instructions increases the chance of crop injury and may result in illegal Gramoxone SL 3.0 residue in the grain. Yield reductions may occur at any stage of growth. Yield reductions can be expected in exchange for improved grain quality. | | | |
| USE RESTRICTIONS | | | |
| <ol style="list-style-type: none"> Refer to Section 6.1 for additional product use restrictions. Maximum Single Application Rate: 0.5 pt/A REI: 48 hours Minimum Application Interval: Not Applicable Maximum Annual Rate: 0.5 pt/A <ol style="list-style-type: none"> DO NOT exceed 0.19 lb cation/A/year of paraquat-containing products. DO NOT make more than 1 application per year. DO NOT apply this product through any type of irrigation system. DO NOT graze treated fields or feed treated forage or straw to livestock. Pre-Harvest Interval (PHI): Harvest crop at normal maturity. | | | |

Gramoxone® and the SYNGENTA Logo are Trademarks of a Syngenta Group Company.

©2022 Syngenta

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina, 27419-8300