

---

**SUPPLEMENTAL LABELING**

---

**Syngenta Crop Protection, LLC**  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300

**SCP 1571B-S3 0820**

<b>OXATHIPIPROLIN</b>	<b>GROUP</b>	<b>49</b>	<b>FUNGICIDE</b>
-----------------------	--------------	-----------	------------------

**Orondis® Gold 200**

Fungicide

**This supplemental label expires on 08/17/2023 and must not be used or distributed after this date.**

Active Ingredient:

Oxathiapiprolin*	18.7%
------------------	-------

Other Ingredients:	81.3%
--------------------	-------

Total:	100.0%
--------	--------

\*CAS No. 1003318-67-9

Orondis® Gold 200 is formulated as a suspension concentrate and contains 1.67 pounds of oxathiapiprolin per gallon of product.

**KEEP OUT OF REACH OF CHILDREN.**

**EPA Reg. No. 100-1571**

**All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed. Before using Orondis Gold 200 as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

**syngenta®**

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### Blueberry, lowbush

Crops (including cultivars and or varieties of these)			
Blueberry, lowbush Not for use in CA			
Target Diseases	Rate fl oz/A (lb ai/A)	Application Timing	Use Directions
Phytophthora root rot ( <i>Phytophthora</i> spp.)	4.8 – 9.6 (0.06 - 0.12)	<p><b>Soil Application:</b></p> <p>For <b>New Plantings</b>, make a single soil application at planting. A second application may be used 7-14 days later if needed.</p> <p>For <b>Established Plantings</b>, make a single soil application as soon as plants begin growing in the spring. A second application may be used 7-14 days later if needed.</p>	<p>Apply as a soil drench, or soil directed spray, or through irrigation water (micro-sprinkler, or drip).</p> <p>For effective disease control, ensure that the product solution thoroughly wets the target root zone. If the application method does not move the product to the root zone, and rain is not imminent, then follow with irrigation.</p> <p>See <b>Section 4.5</b> for drip irrigation instructions.</p> <p>Use the higher rate in fields with a history of <i>Phytophthora</i> disease.</p>
<b>Resistance Management:</b> <ul style="list-style-type: none"> <li>Refer to <b>Section 3.2</b>.</li> <li>Make no more than 2 sequential applications of Orondis Gold 200 (or any other FRAC 49-containing product) before rotating to a fungicide with a different mode of action.</li> </ul>			
USE RESTRICTIONS			
<ol style="list-style-type: none"> <li>Refer to <b>Section 6.1</b> for additional product use restrictions.</li> <li><b>Maximum Single Application Rate:</b> <ol style="list-style-type: none"> <li><b>Soil Applications:</b> <b>DO NOT</b> exceed the maximum rate listed in the table for this application type.</li> </ol> </li> <li><b>Maximum Number of Applications per Year:</b> <ol style="list-style-type: none"> <li><b>Soil applications:</b> <b>DO NOT</b> make more than 2 applications at the maximum rate per year.</li> </ol> </li> <li><b>Minimum Application Interval:</b> 7 days</li> <li><b>Maximum Annual Rate:</b> <ol style="list-style-type: none"> <li><b>Soil Applications:</b> 19.2 fl oz/A/year (0.25 lb ai/A/year)</li> <li><b>DO NOT</b> apply more than 0.25 lb ai/A/year of soil-applied oxathiapiprolin-containing products.</li> </ol> </li> <li><b>DO NOT</b> use in nursery production of transplanted crops.</li> <li><b>Pre-harvest Interval (PHI):</b> <ol style="list-style-type: none"> <li><b>Soil Applications:</b> 0 days</li> </ol> </li> </ol>			

## Bushberry subgroup 13-07B, except blueberry, lowbush

Crops (including all cultivars, varieties, and/or hybrids of these)			
Aronia berry		Elderberry	Jostaberry
Blueberry, highbush		European barberry	Juneberry (Saskatoon Berry)
Buffalo currant		Gooseberry	Lingonberry
Chilean guava		Highbush cranberry	Native currant
Currant, black		Honeysuckle, edible	Salal
Currant, red		Huckleberry	Sea buckthorn
Target Disease	Rate fl oz/A (lb ai/A)	Application Timing	Use Directions
Phytophthora root rot ( <i>Phytophthora</i> spp.)	4.8 – 9.6 (0.06 - 0.12)	<p><b>New Plantings:</b> Make first application at planting. Make a second application a minimum of 7-30* days later, coinciding with a period favorable for root rot development.</p> <p><b>Established Plantings:</b> Make first application before plants start to grow in the spring. Make a second application a minimum of 7-30* days later, coinciding with a period favorable for root rot development.</p> <p>*Although the minimum number of days before treatment is 7-30, a minimum of 30 days before treatment is recommended.</p>	<p>Apply as a soil drench, or soil directed spray, or through irrigation water (micro-sprinkler or drip).</p> <p>For effective disease control, ensure that the product solution thoroughly wets the target root zone. If the application method does not move the product to the root zone, and rain is not imminent, then follow with irrigation.</p> <p>See <b>Section 4.1.2</b> for at-planting, or transplant water instructions. See <b>Section 4.5</b> for drip irrigation instructions.</p> <p>Use the higher rate in fields with a history of <i>Phytophthora</i> disease. Orondis Gold 200 will not revitalize plants showing moderate to severe root rot symptoms.</p> <p>Use Orondis Gold 200 in conjunction with good cultural practices to manage diseases.</p>
<b>Resistance Management:</b> <ul style="list-style-type: none"> <li>Refer to <b>Section 3.2</b>.</li> <li>Make no more than 2 sequential applications of Orondis Gold 200 (or any other FRAC 49-containing product) before rotating to a fungicide with a different mode of action.</li> </ul>			
USE RESTRICTIONS			
<ol style="list-style-type: none"> <li>Refer to <b>Section 6.1</b> for additional product use restrictions.</li> <li><b>Maximum Single Application Rate:</b> DO NOT exceed the maximum rate listed in the table for this application type.</li> <li><b>Maximum Number of Applications per Year:</b> DO NOT make more than 2 applications at the maximum rate per year.</li> <li><b>Minimum Application Interval:</b> 7 days</li> <li><b>Maximum Annual Rate:</b> 19.2 fl oz/A/year (0.25 lb ai/A/year) <ol style="list-style-type: none"> <li>DO NOT apply more than 0.25 lb ai/A/year of soil-applied oxathiapiprolin-containing products.</li> </ol> </li> <li>DO NOT use in nursery production of transplanted crops.</li> <li><b>Pre-harvest Interval (PHI):</b> 1 day</li> </ol>			

Orondis® and the Syngenta logo are trademarks of a Syngenta Group Company

©2020 Syngenta

Chron :

Orondis 1571 IR-4 SUPP DEC2015-Version C-Clean – rk – 11/1/16  
000100-01571.20151223C.ORONDIS\_IR-4\_SUPP\_DEC2015-CL.pdf

SCP 1571B-S2 1116 – rk 7/5/17  
FPSL for WA state as requested by Business