



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

Axiom® DF Herbicide

For control of weeds in Christmas Tree Plantations
(Douglas Firs and True Firs)

EPA Reg. No. 264-766

EPA SLN. No. OR-150015

FIFRA Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON

THIS LABEL IS VALID UNTIL DECEMBER 31, 2020 OR UNTIL OTHERWISE AMENDED, WITHDRAWN, CANCELED, OR SUSPENDED.

Axiom® DF Herbicide

ACTIVE INGREDIENTS:

Flufenacet

N-(4-Fluorophenyl)-*N*-(1-methylethyl)-2-[[5-(trifluoromethyl)-1,3,4-thiadiazol-2-yl]-oxy]acetamide 54.4%

Metribuzin:

4-Amino-6-(1,1-dimethylethyl)-3-(methylthio)-1,2,4-triazin-5(4*H*)-one..... 13.6%

OTHER INGREDIENTS: 32.0%
100.0%

STOP - Read the label before use
Keep out of reach of children
CAUTION

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not allow sprays to drift onto adjacent desirable plants. Do not contaminate water when disposing of equipment washwaters or rinsate.

Metribuzin has properties that may result in ground water contamination. Application in areas where soils are permeable or coarse and ground water is near the surface could result in ground water contamination.

Ground Water: Flufenacet is known to leach through soil into ground water under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Surface Water: Flufenacet and metribuzin residues can contaminate surface water through spray drift. Under some conditions, they also have high potential for runoff into surface water (primarily via dissolution in runoff water or erosion), for several to many months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlying tile drainage systems that drain to surface waters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This SLN label and the main EPA-registered label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, and precautions on this SLN label and the main EPA-registered label.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

CHRISTMAS TREE PLANTATIONS	RATE oz/A	WEEDS
Douglas Fir True Fir (including Noble, Grand, Frazier and Nordman)	12 – 20	Controlled: Annual Bluegrass Annual Ryegrass <i>Bromus</i> spp. Fall panicum Rattail fescue Suppressed: False Dandelion Lambsquarters Pigweeds Mustards
REMARKS: Apply the specified dosage per acre as a pre- or very early post-emergent (1-leaf stage) application to weeds. Use the higher rate for heavier weed populations or where longer control is needed. Treat only dormant trees (prior to bud swell) which have gone through one or more growing seasons. Apply in a minimum of 10 gal of water per acre by ground applications only. Axiom DF Herbicide may be applied only one time per growing season. Allow at least 11 months between applications.		

RESTRICTIONS/PRECAUTIONS:

- Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours following application.
- Do not apply to trees that are under stress (drought, insects, disease).
- For improved control of emerged weeds, and/or control of additional weed species, a tank-mix application with another registered herbicide may be used. Consult tank-mix partner label for use directions and precautions.
- Do not apply this product through any type of irrigation system. Do not apply aerially.

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937))*