

**RESTRICTED USE PESTICIDE
DUE TO ACUTE TOXICITY**

**FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS ONLY– NOT TO BE
USED BY UNCERTIFIED PERSONS WORKING UNDER THE SUPERVISION OF A
CERTIFIED APPLICATOR.**

syngenta.

ITERA

Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF COLORADO

**Gramoxone® SL 2.0
For Use as a Harvest Aid in Alfalfa Grown for Seed**

**EPA Reg. No. 100-1431
EPA SLN No. CO-160002**

**This label expires and must not be distributed or used in accordance with this SLN
registration after July 1, 2022**

**KEEP OUT OF REACH OF CHILDREN / MANTENER FUERA DEL ALCANCE DE LOS
NIÑOS.**



**DANGER / POISON
PELIGRO / VENENO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered Gramoxone SL 2.0 label.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL
MAY RESULT IN POOR DESICCATION, ILLEGAL RESIDUES AND/OR CROP INJURY.**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

Specific Crop Use Directions

Crop	Use Pattern	Rate Per Treated Acre	Minimum Spray Volume gal/acre	Preharvest Interval (Days)
Alfalfa Desiccation of alfalfa to facilitate harvest of alfalfa seed	Broadcast	2.5 – 4.0 pts.	Ground: 20-25 gals. Air: 5-10 gals.	See precautions below

ALWAYS ADD ONE OF THE FOLLOWING:

Add a nonionic surfactant containing 75% or more surface-active agent at 0.125% v/v (1 pint per 100 gallons), OR add a nonionic surfactant containing 50-74% surface-active agent at 0.25% v/v (2 pints per 100 gallons) of the finished spray volume for ground applications. For aerial applications, add a nonionic surfactant at 0.25% v/v (2 pints per 100 gallons) of the finished spray volume

Precautions, Restrictions, and Comments:

- DO NOT apply more than two applications per year
- DO NOT harvest until at least 4 days after application.
- DO NOT apply when weather conditions favor drift from treated areas.
- DO NOT apply by ground equipment within 25 ft. or by air within 75 ft. of lakes; reservoirs; rivers; permanent streams, marshes or natural ponds; estuaries; and commercial fish ponds.
- For use only on fields in production of alfalfa seed. Not for use on fields producing alfalfa for livestock feed. No portion of the treated field, including seed, seed screenings, hay forage, or stubble, may be used for human or animal feed. Screenings from alfalfa seed processing are prohibited from feed channels. All Gramoxone SL 2.0 treated alfalfa seed screenings must be removed from the feed market.
- DO NOT cut current year's alfalfa seed crop for hay or forage. DO NOT graze current year's treated alfalfa seed crops.
- Treated alfalfa seed is not to be used for sprouting. All alfalfa seed treated with Gramoxone SL 2.0 is to be tagged at processing plants, "NOT FOR HUMAN CONSUMPTION OR ANIMAL FEED". It shall be the grower's responsibility to notify the processing plants of any seed crop treated with Gramoxone SL 2.0.
- DO NOT apply this product through any type of irrigation system.

©2021 Syngenta

Gramoxone® and the Syngenta logo are trademarks of a Syngenta Group Company

24(c) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Label Code: CO1431111BA0821