

PENTHIOPYRAD	GROUP	7	FUNGICIDE
--------------	-------	---	-----------

FIFRA Section 24(c) Special Local Need (SLN) Label

Corteva Agriscience

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF NEW YORK

Fontelis®

EPA Reg. No. 352-834

SLN No. NY-130003

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2028

FOR PLANT DISEASE CONTROL ON APPLE, PEAR, AND STONE FRUIT IN THE STATE OF NEW YORK

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This state-specific Section 24(c) label and the federal label for this product must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, and precautions on the container label for EPA Reg. No. 352-834.
- Follow all directions on this supplemental label.

Not for sale, sale into, distribution, or use in Nassau or Suffolk Counties, New York.

Refer to the product label for additional Precautions and Restrictions, Mixing Instructions and Application Directions.

PRODUCT INFORMATION

Fontelis® fungicide contains mineral oil in the formulation. Fontelis is a broad-spectrum fungicide, recommended for control of foliar and soil-borne plant diseases, and has preventive, curative, and locally systemic activity. Fontelis must be applied in a regularly scheduled protective spray program in rotation with other fungicides. See directions below for specific crop/disease recommendations

DIRECTIONS FOR USE

Fontelis can be applied with commonly used ground equipment, hose-end, pressurized, greenhouse and hand-held sprayers, air or chemigation equipment, except as otherwise directed, using sufficient water to obtain thorough coverage of plants. Thorough coverage of all foliage is essential for effective disease control. Maintain agitation during mixing and application to assure uniform product suspension.

Application Volumes

- For conventional ground application, apply a minimum of 15 gallons per acre, increasing the spray volume as the plants mature to ensure thorough coverage of foliage.
- For air-assisted ground application, apply a minimum of 10 gallons per acre.
- For aerial application, apply a minimum of 2 gallons per acre (10 gallons per acre for trees and orchards).

CULTIVAR/VARIETAL CROP SAFETY

Not all crops within a crop group, and not all varieties, cultivars or hybrids of crops, have been individually tested for crop safety. It is not possible to evaluate for crop safety all applications of Fontelis on all crops within a crop group, on all varieties, cultivars, or hybrids of those crops, or under all environmental conditions and growing circumstances. To test for crop safety, apply the product in accordance with the label instructions to a small area of the target crop to ensure that a phytotoxic response will not occur, especially where the application is a new use of the product by the applicator.

TANK MIXTURES

Fontelis fungicide contains mineral oil in the formulation. If tank mixing with oil sensitive products (for example, captan and sulfur) read and follow their label restrictions.

Always follow the tank mix instructions of the product label that is most restrictive. The crop safety of all tank mixtures with Fontelis which may include physically compatible pesticides, fertilizers, adjuvants, and/or additives, has not been tested. When considering a tank mixture with Fontelis which is not specifically described on product labeling or in other Corteva product use instructions, it is important to understand crop safety. To test for crop safety prepare a small volume of the intended tank mixture, apply it to an area of the target crop as directed by both this and the tank mix partner product labels, and observe the treated crop to ensure that a phytotoxic response does not occur. Corteva will not be responsible for any crop injury arising from the use of a tank mixture that is not specifically described on Fontelis product labeling or in other Corteva product use instructions. Some materials including oils, surfactants, adjuvants and pesticide formulations when applied individually, sequentially, or in tank mixtures may solubilize the plant cuticle, facilitate penetration into plant tissue, and increase the potential for crop injury.

Always follow the tank mix instructions of the product label that is most restrictive. Consult a Corteva representative or local agricultural authority for more information concerning tank mixtures.

APPLICATION INFORMATION

Mixing Instructions

1. Fill clean spray tank 1/4 - 1/2 full of water.
2. While agitating, add the required amount of Fontelis, continuing agitation until the product is completely dispersed.
3. Continue filling the tank, with agitation, following the sequence listed below in "tank mixing sequence."

Adjuvants

Fontelis fungicide may be used with adjuvants, for example, nonionic surfactants, crop oils, methylated seed oils, and blends at typical agricultural use rates for these adjuvants.

USE RATES AND APPLICATION INSTRUCTIONS

Crop/Crop Group	Target Diseases	Use Rate per Acre (fl oz)	Remarks
Apple; pear	Alternaria leaf spots* (<i>Alternaria</i> spp.) Scab, apple or pear (<i>Venturia inaequalis</i> (apple), <i>Venturia pirina</i> (pear)) Powdery mildew (<i>Podosphaera leucotricha</i>) Rusts* (<i>Gymnosporangium</i> spp.)	16 to 20 fl oz	Begin applications prior to disease development and continue on a 7- to 21-day interval. Use higher rate and shorter interval when disease pressure is high. - Application interval for scab is 7 to 10 days. For apple scab, a reliable disease forecasting system should be used. NOTE: Do NOT tank-mix Fontelis with thinning agents.

Make no more than 2 sequential applications of Fontelis before switching to a fungicide with a different mode of action. Minimum time from application to harvest (PHI) is 28 days. Do not exceed 61 fl oz/acre per year.

Crop/Crop Group	Target Diseases	Use Rate per Acre (fl oz)	Remarks
Stone fruits Apricot; cherry, sweet; cherry, tart; nectarine; peach; plum; plum, Chickasaw; plum, Damson; plum, Japanese; plumcot; prune (fresh)	Alternaria rot (<i>Alternaria</i> spp.) Botrytis rots (<i>Botrytis cinerea</i>) Brown rot blossom blight and fruit rot (<i>Monilinia</i> spp.) Green fruit rot (<i>Sclerotinia sclerotiorum</i>) Powdery mildew (<i>Podosphaera clandestina</i> , <i>Sphaerotheca pannosa</i>) Rust (<i>Tranzschelia discolor</i>) Scab (<i>Cladosporium carpophilum</i>) Shot hole (<i>Wilsonomyces carpophilus</i>)	14 to 20 fl oz	Begin applications prior to disease development and continue on a 7- to 14-day interval. Use higher rate and shorter interval when disease pressure is high.
	Disease Suppression: Cherry leaf spot (<i>Blumeriella jaapii</i>)	14 to 20 fl oz	

Make no more than 2 sequential applications of Fontelis before switching to a fungicide with a different mode of action. Minimum time from application to harvest (PHI) is 0 days. Do not exceed 61 fl oz/acre per year.

IMPORTANT NOTICE:

BEFORE BUYING OR USING FONTELIS FUNGICIDE, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS IN THIS SECTION 24(c) LABEL AND ON THE EPA-REGISTERED LABEL.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. Follow the instructions carefully. This labeling must be in possession of the user at the time of pesticide application.

Read the Limitation of Warranty and Liability on the product label before buying or using this product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the product label.

^{™®} Trademarks of Corteva Agriscience and its affiliated companies

R613-009

Accepted: 10/25/23

Replaces: R-1465.1