

|            |       |   |           |
|------------|-------|---|-----------|
| CLOPYRALID | GROUP | 4 | HERBICIDE |
|------------|-------|---|-----------|

# FIFRA Section 24(c) Special Local Need (SLN) Label

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Stinger<sup>®</sup>

EPA Reg. No. 62719-73

24(c) Special Local Need Registration SLN IN-150002

**Postemergence Broadleaf Weed Control in Blueberry  
(For Distribution and Use Only in the State of Indiana)**

**This label expires and must not be distributed or used in accordance with this SLN registration  
after December 31, 2027**

### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of pesticide application.
- Read the label affixed to the container for Stinger<sup>®</sup> herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- In addition to restrictions and limitations imposed by this Special Local Need labeling, use of Stinger herbicide is subject to all use precautions and limitations imposed by the label affixed to the container for Stinger herbicide.

### Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters. Do not contaminate water used for irrigation or domestic purposes.

### Ground Water Advisory

Clopyralid is a chemical which can travel (seep or leach) through soil and under certain conditions contaminate groundwater which may be used for irrigation or drinking purposes. Users are advised not to apply clopyralid where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an underlying aquifer is shallow, or to soils containing sinkholes over limestone bedrock, severely fractured surfaces, and substrates which would allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

### Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a medium potential for reaching surface water via runoff for several weeks after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as

ponds, streams, and springs will reduce the potential loading of clopyralid from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

## Directions for Use

Refer to the product label for Precautions and Restrictions, Mixing Instructions, and Application Directions.

### Target Broadleaf Weeds, Application Rate, and Use Restrictions:

| Target Weeds  | Stinger<br>(fl oz/acre)   | Use Restrictions  |
|---|---|---|
| beggartick<br>common ragweed<br>giant ragweed<br>wild bean  | 2.67 - 4<br>(1/6 - 1/4 pint/acre)<br>(0.063 – 0.094 lb ae<br>clopyralid/acre) | <ul style="list-style-type: none"> <li>• <b>Preharvest interval:</b> Do not apply within 30 days of harvest.</li> <li>• Do not exceed a total of 10.6 fl oz (0.25 lb ae clopyralid) per acre per year.</li> <li>• Do not apply within 5 hours of expected rainfall or irrigation.</li> <li>• Do not apply by aircraft.</li> <li>• <b>Chemigation:</b> Do not apply through any type of irrigation system.</li> <li>• Do not apply to low growing varieties of blueberry.</li> </ul> |
| annual sowthistle (suppression)<br>aster species<br>Canada thistle (suppression)<br>clover species<br>dandelion<br>joe-pye weed<br>narrow-leaved golden rod | 4 – 5.3<br>(1/4 – 1/3 pint/acre)<br>(0.094 – 0.125 lb ae<br>clopyralid/acre)  |   |

**Application timing:** Blueberry plants are more sensitive to Stinger applied in the spring prior to bloom, before and/or during the crop's annual flush of growth, than after bloom. Do not apply Stinger from one week prior to bloom until one week after bloom. After bloom, apply Stinger up until 30 days prior to harvest. Stinger can be applied after harvest. Determine the rate of Stinger based upon the targeted weed species and whether one or two sprays will be applied during the growing season. **Note:** Stinger is a residual herbicide and applications must be based upon accurate rate per acre calibrations. Applications of Stinger can injure the blueberry plant and significantly reduce yields depending upon rates used, timing of application, and environmental conditions.

**Application Rate:** Apply Stinger at 2.67 to 5.3 fl oz per acre directed to the soil from the blueberry plants to the row middle without contacting the foliage or woody portions of blueberry plants. Apply with ground broadcast equipment, backpack sprayer, or wipe applicator in a total spray volume of a minimum of 10 gallons per acre. Hand-held sprayers may be used for spot applications. Refer to the package label for conversion rates, mixing, and application directions for hand-held sprayers.

Use the lower rate in the rate range for young succulent growth for sensitive weed species. Use the higher rate in the rate range for less sensitive weed species, perennials, and under environmental conditions where target weeds are less susceptible. Make a maximum of two applications with the total usage of Stinger from all types of applications not to exceed 10.6 fl oz per acre or 2/3 pint per acre (0.25 lb ae clopyralid per acre) per year.

**Wipe Treatments:** For wipe treatments, apply a 2% solution of Stinger in water (2.5 fl oz or 75 ml per gallon). Make a maximum of two applications with the total usage of Stinger from all types of applications not to exceed 10.6 fl oz per acre or 2/3 pint per acre (0.25 lb ae clopyralid per acre) per annual growing season. Do not permit Stinger to contact desirable foliage or crop injury will result.

### **Special Conditions and Risks of Use**

**USE OF Stinger HERBICIDE (THE "PRODUCT") ON BLUEBERRY (THE "CROP") MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS. READ AND UNDERSTAND THESE SPECIAL CONDITIONS AND RISKS OF USE BEFORE USING THE PRODUCT ON THE CROP.**

**Corteva Agriscience makes the product available for use in the manner described in this Special Local Need Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury or loss. The decision to use this Product in the manner described in this Special Local Need Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls, and (iv) any other relevant factors.**

**By purchasing the Product for use, or using the Product, in the manner described in this Special Local Need Labeling, you acknowledge and accept that:**

- (1) You assume all risk of Crop injury, Crop yield reduction and Crop loss;
- (2) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this Supplemental labeling;
- (3) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaim all warranties, expressed or implied, including any implied warranty of merchantability;
- (4) Corteva Agriscience disclaims all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to Crop injury, Crop yield reduction and/or Crop loss;
- (5) These Conditions and Risks of Use for Special Local Need supersede any contrary representations or recommendations by Corteva Agriscience, or its respective agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

**If these Special Conditions and Risks of Use are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container. These Special Conditions and Risks of Use for Special Local Need for Special Local Need are required by Corteva Agriscience and not specified by the US EPA or the State of Indiana.**

<sup>TM</sup>® Trademarks of Corteva Agriscience and its affiliated companies

R043-148

Accepted: 04/27/22

Replaces Label: R043-087