



Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF NORTH DAKOTA

Reflex® Herbicide

For Control of Weeds in Dry Beans in North Dakota West of U.S. Highway 281 in the Counties of Benson, Bottineau, Burleigh, Dickey, Eddy, Emmons, Foster, Grant, Kidder, Lamoure, Logan, McHenry, McIntosh, McLean, Mercer, Morton, Oliver, Pierce, Renville, Rolette, Sheridan, Sioux, Stutsman, Towner, Ward and Wells

**EPA Reg. No. 100-993
EPA SLN No. ND-180004**

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2028

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.
- Failure to follow the directions for use and precautions on this label may result in poor pest control, crop injury, or illegal residues.

Application Methods

Apply Reflex Herbicide at 0.75 pt/A as a postemergence broadcast application to dry beans. Apply in a minimum of 15 GPA.

Spray Additives

Add methylated seed oil (MSO) or crop oil concentrate (COC) at 1% v/v to the final spray volume.

Weeds Controlled

Weed	Maximum Number of Leaves At Application
Canola, volunteer	4
Mustard, wild	4
Nightshade, black	2
Pigweed, redroot	4

Rotational Crop Restrictions

Rotational Crops	Minimum Rotational Interval after Reflex Herbicide Application (Months)
Bean, Dry Potato Soybean	0 months
Small Grains, such as Wheat, Barley, Rye	9 months (4)
Edible-podded beans and peas not otherwise specified in this label Pea, Dry	12 months
Corn, Field* Corn, Seed*	18 months (10)
All other crops not listed above	Minimum 18 months and a successful field bioassay (refer to Field Bioassay Instructions section) must be performed before planting rotational crop

*For fields with greater than 4% organic matter, a successful field bioassay (refer to **Field Bioassay Instructions** section) must be performed before planting field corn or seed corn.

Field Bioassay Instructions

Plant several strips of the desired crop variety across the field which has been previously treated with Reflex Herbicide. Plant the strips perpendicular to the direction Reflex Herbicide was applied. The strips should be located so that all the different field conditions are encountered, including differences in soil texture, organic matter, pH, and drainage. If the crop does not show visible symptoms of injury and/or stand reduction, this field can be planted to this crop. If visible injury and/or stand reduction occur, do not plant the field to this crop; plant only a labeled crop.

Use Restrictions

- Do not exceed 0.75 pt/A of Reflex Herbicide in any one year.
- Do not apply a fomesafen containing product to any field more than once every two years.
- Do not make a Reflex Herbicide application after June 20.
- Do not harvest dry beans within 45 days of application.
- Do not graze animals on green forage or stubble.
- Do not utilize hay or straw for animal feed or bedding.
- Avoid overlapping spray swaths, as injury may occur to rotational crops.

©2023 Syngenta

Reflex® and the Syngenta logo are trademarks of a Syngenta Group Company

24(c) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Label Code: ND0993070BA1023