

## SUPPLEMENTAL LABEL

FENPYROXIMATE GROUP 21A INSECTICIDE



EPA Reg. No. 71711-4

For Use On: Caneberry Subgroup (Crop Subgroup 13-07A)

This supplemental label expires December 12, 2022 and must not be used or distributed after this date.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling and the EPA approved container label must be in the possession of the user at the time of application. Read the label affixed to the container for FUJIMITE® Miticide/Insecticide before applying. Use of FUJIMITE Miticide/Insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for FUJIMITE Miticide/Insecticide.

New use directions appear on this supplemental labeling that may be different from those that appear on the container label.

### APPLICATION RATE CHART FOR FUJIMITE MITICIDE/INSECTICIDE

Caneberry Subgroup (Crop Subgroup 13-07A)		
blackberry (including Andean blackberry, arctic blackberry, bingleberry, black satin berry, boysenberry, brombeere, California blackberry, Chesterberry, Cherokee blackberry, Cheyenne blackberry, common blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, evergreen blackberry, Himalayaberry, hullberry, lavacaberry, loganberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, mora, mures deronce, nectarberry, Northern dewberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, Southern dewberry, tayberry, youngberry, zarzamora); raspberry, black and red; wild raspberry; cultivars, varieties, and/or hybrids of these		
Pest	Rate/Acre	Use Directions
Willamette spider mite	1.5 to 2.0 pints (0.08 to 0.105 lb ai)	<ul style="list-style-type: none"><li>• Apply by ground using a minimum of 30 gallons of water per acre.</li><li>• For vines with a heavy canopy or in high pressure situations, higher water volumes are recommended.</li><li>• Allow 14 days between applications.</li><li>• Preharvest Interval (PHI): 1 day</li></ul>
Leafhoppers <sup>1</sup> Mealybugs Mites (see <b>Target Species</b> ) Powdery Mildew*	2.0 pints (0.105 lb ai)	<p>USE RESTRICTIONS</p> <ul style="list-style-type: none"><li>• Do not apply by air.</li><li>• Do not apply through any type of irrigation system.</li><li>• Do not apply more than 4.0 pints (0.210 lb ai) per acre per year.</li><li>• Do not make more than 2 applications per year.</li></ul>
*suppression		
<sup>1</sup> Best control of leafhoppers is achieved by applications when majority of the population is in an immature development stage.		

©2020 Nichino America, Inc. FujiMite and Nichino America logo are registered trademarks of Nichino America, Inc.

**NICHINO**  
**AMERICA**

Nichino America, Inc.  
4550 Linden Hill Road, Suite 501  
Wilmington, DE 19808  
888-740-7700