



ITERA

Section 24(c) Special Local Need Label

ACIBENZOLAR-S-METHYL

GROUP

P01

FUNGICIDE

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MICHIGAN

**Actigard® 50WG Plant Activator
For Protection Against Bacterial Spot on Bell Peppers**

EPA Reg. No. 100-922

EPA SLN No. MI-170003

**This label expires and must not be distributed or used in accordance with this SLN
registration after December 31, 2027**

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered Actigard 50WG Plant Activator label.

For protection against bacterial spot, *Xanthomonas* spp., on bell peppers apply Actigard 50WG at 0.25 to 0.33 oz/A. Actigard 50WG can be applied as a foliar application or drip irrigation. To reduce crop tolerance effects and improve disease protection, do not apply within one week of transplanting. Apply once plants are actively growing and continue as needed on a 7 day interval. Do not apply Actigard once plants begin to bloom.

Actigard can be applied by ground, drip irrigation, or by air. Apply in sufficient water to ensure adequate coverage. A minimum of 20 GPA should be used for ground applications and a minimum of 10 GPA should be used for aerial applications. If disease is present in the production area, another registered product for that disease should be applied at the same timing as the Actigard application (can tank mix if applying Actigard as a foliar spray). Banded applications can be made early when plants are small (7" proportional band) as long as coverage is adequate.

Precautions:

- 1) Actigard 50WG should be applied to healthy, actively growing plants.
- 2) Use the higher rate over time, even if water volume does not increase.

Restrictions:

- 1) Do not apply Actigard 50WG to plants that are stressed due to drought, excessive moisture, cold weather, or herbicide injury, etc.
- 2) Do not apply more than 8 times per crop per season.
- 3) Do not apply on less than 7-day interval.
- 4) Do not apply more than 6 oz (0.1875 lb ai) of Actigard 50WG per acre per season. Allow 14 days between the last application and harvest (14-day PHI).

Precaution: Under certain conditions, this product, when used on peppers, may lead to reductions in yield. Syngenta recommends that the user and/or grower test this product in order to determine its suitability for its intended use in peppers. Syngenta makes this product available to the user, and/or grower solely to the extent that the benefit and utility, in the sole opinion of the user and/or grower, outweigh the extent of potential reductions in yield of peppers following use of Actigard 50WG. The decision to use or not use Actigard 50WG in peppers must be made by each individual user and/or grower on the basis of possible reductions in yield, the severity of disease incidence, the cost of alternate disease control options, if any, and other factors.

©2022 Syngenta

Actigard® and the Syngenta logo are trademarks of a Syngenta Group 'Company

24(c) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Label Code: MI0922060BA1222