

**TIDE**

TEBUCONAZOLE | GROUP 3 | FUNGICIDE

TIDE USA TEBU 3.6F FUNGICIDE**ACTIVE INGREDIENT**

Tebuconazole
 (alpha-[2-(4-chlorophenyl)ethyl]
 alpha-(1,1-dimethylethyl)-1H-
 1,2,4-triazole-1-ethanol).....38.7%

OTHER INGREDIENTS61.3%

TOTAL..... 100.0%

Contains 3.6 pounds Tebuconazole per gallon.

EPA Reg.No. 84229-34

EPA Est. No. 69845-CHN-002

39578-TX-1 67194-MEX-001

5905-GA-01 70815-GA-001

5905-AR-01 88302-IND-2

5905-CA-01 88302-IND-3

5905-IA-01 88302-IND-4

84264-IND-3 95781-IND-1

95348-IND-1 95350-IND-1

**KEEP OUT OF REACH
 OF CHILDREN
 CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand this label, find someone to explain it to you in detail.)

See inside label booklet for First Aid, additional Precautionary Statements and Directions for Use including Storage and Disposal instructions.

Manufactured for:

Tide International USA, Inc.
 21 Hubble, Irvine, CA 92618, USA

Net Contents: 1 Gallon

PEEL BACK BOOK HERE AND RESEAL AFTER OPENING

PROOF

Proof date: 06/17/2024
 Customer: Tide USA
 Job number: TIC-FUN8422934
 Label size: 145 x 165mm
 Leaflet flat size: 275 x 165mm
 Leaflet folded size: 145 x 165mm

Label colors: PANTONE Process Black C,
 PANTONE 2413 C
 PANTONE Process Blue C
 Leaflet "in" colors: Black
 Leaflet "out" colors: PANTONE Process Black C,
 PANTONE 2413 C
 PANTONE Process Blue C



TEBUCONAZOLE GROUP 3 FUNGICIDE

TIDE USA TEBU 3.6F FUNGICIDE

ACTIVE INGREDIENT

Tebuconazole

(alpha-[2-(4-chlorophenyl)ethyl]

alpha-(1,1-dimethylethyl)-1H-

1,2,4-triazole-1-ethanol).....38.7%

OTHER INGREDIENTS61.3%**TOTAL**..... 100.0%

Contains 3.6 pounds Tebuconazole per gallon.

EPA Reg.No. 84229-34**EPA Est. No.** 69845-CHN-002 39578-TX-1 67194-MEX-001 5905-GA-01 70815-GA-001 5905-AR-01 88302-IND-2 5905-CA-01 88302-IND-3 5905-IA-01 88302-IND-4 84264-IND-3 95781-IND-1 95348-IND-1 95350-IND-1

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand this label, find someone to explain it to you in detail.)

See inside label booklet for First Aid, additional Precautionary Statements and Directions for Use including Storage and Disposal instructions.

Manufactured for:

Tide International USA, Inc.
21 Hubble, Irvine, CA 92618, USA**Net Contents: 1 Gallon**

PEEL BACK BOOK HERE AND RESEAL AFTER OPENING

FIRST AID

If swallowed:	<ul style="list-style-type: none">● Call a poison control center or doctor immediately for treatment advice.● Have person sip a glass of water if able to swallow.● DO NOT induce vomiting unless told to do so by the poison control center or doctor.● DO NOT give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none">● Take off contaminated clothing.● Rinse skin immediately with plenty of water for 15-20 minutes.● Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none">● Move person to fresh air.● If person is not breathing, call 911 or an ambulance, then give artificial respiration preferably by mouth-to-mouth, if possible.● Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none">● Hold eye open and rinse slowly and gently with water for 15-20 minutes.● Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.● Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact CHEMTREC at 1-800-424-9300 for emergency medical treatment information.	
NOTE TO PHYSICIAN: No specific antidote. Treat symptomatically.	
Symptoms of Poisoning: The compound does not cause any definite symptoms that would be diagnostic. Contact with eyes may cause irritation.	

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, absorbed through skin, or inhaled. Avoid contact with eyes, skin, or clothing. Avoid breathing spray mist or vapor. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, including barrier laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately, if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when disposing of equipment washwater or rinseate.

Ground Water Advisory: Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams and springs will reduce the potential for contamination of water from rainfall- runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

PHYSICAL/CHEMICAL HAZARDS

DO NOT mix or allow to come in contact with oxidizing agents. A hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers or protected supervisors may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Aerial application Prohibited in New York State.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) listed in the application directions for the crop being treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, including barrier laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Golf Course Turf and Landscape Uses: Keep children and pets out of treated areas until sprays have dried.

INFORMATION FOR AGRICULTURAL USES

SHAKE WELL BEFORE USING

Spray Volume: Apply Tide USA TEBU 3.6F Fungicide with ground or aerial equipment using sufficient volume of spray to provide thorough coverage. Tide USA TEBU 3.6F Fungicide may be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. Check equipment calibration frequently. Continuous agitation is required to keep the material in suspension. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase the spray volume per acre for complete crop coverage. Use the higher rate under conditions of severe disease pressure. Also, see local State Extension Service guidance for application schedules.

Chemigation: Apply Tide USA TEBU 3.6F through irrigation equipment only to crops and diseases for which the chemigation use is specified. Apply Tide USA TEBU 3.6F only through center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move or drip (trickle) irrigation systems. **DO NOT** apply Tide USA TEBU 3.6F through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts. **DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments if needed.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, for example, a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension. Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. Pesticide may be applied continuously for the duration of the water application.

Mixing: Add specified amount of Tide USA TEBU 3.6F Fungicide into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the Tide USA TEBU 3.6F Fungicide must be thoroughly dispersed prior to the addition of other materials.

NOTE: DO NOT tank mix with products containing a prohibition against tank mixing. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Compatibility: To determine the compatibility of Tide USA TEBU 3.6F Fungicide with other products, the following procedure must be followed: Pour the specified proportions of the product into a suitable container of water, mix thoroughly and allow to stand at least five minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible.

OBSERVE THE FOLLOWING RESTRICTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS INCLUDING LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.

Apply only during alternate years in fields adjacent to aquatic areas listed on this label.

DO NOT apply by ground or air within 100 feet of aquatic areas listed on this label.

DO NOT cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.

MANDATORY SPRAY DRIFT REQUIREMENTS

Aerial Applications:

- **DO NOT** release spray at a height greater than 10 ft above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- For all applications, applicators are required to use a medium or coarser spray droplet size (ASABE S572.1).
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- Applicators must use ½ swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Ground Applications:

- Apply with the nozzle height specified by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- For all applications, applicators are required to use a medium or coarser spray droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Boom-less Ground Applications:

- Applicators are required to use a medium or coarser droplet size (ASABE S572.1) for all applications.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure specified for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- Adjust Nozzles - Follow nozzle manufacturer's directions for setting up nozzles. To reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT - Ground Boom

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift. When applying aurally to crops, **DO NOT** release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Boom-less Ground Applications:

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications:

Take precautions to minimize spray drift.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

DO NOT make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on this label as soon as practical after last application. Any crop not specified on this label may be planted into treated areas 120 days after last application.

RESISTANCE MANAGEMENT

For resistance management, Tide USA TEBU 3.6F Fungicide contains a Group 3 fungicide. Any fungal/bacterial population may contain individuals naturally resistant to Tide USA TEBU 3.6F Fungicide and other Group 3 fungicides/bactericides. A gradual or total loss of pest control may occur over time if these fungicides/bactericides are used repeatedly in the same fields. Follow appropriate resistance-management strategies.

To reduce selection pressure for resistant pests:

- Use Tide USA TEBU 3.6F Fungicide in rotation with classes of fungicides with different modes of action.
- Use Tide USA TEBU 3.6F Fungicide as part of a disease management program that includes cultural and biological control where possible.
- Tide USA TEBU 3.6F Fungicide is in the Group 3 class of fungicides. The mode of action for tebuconazole, the active ingredient in this product, is as a demethylation inhibitor of sterol biosynthesis (DMI) which disrupts membrane synthesis by blocking demethylation. Resistance can develop when products with the same mode of action are used repeatedly.
- Consult your State or local agricultural pest control advisor(s) for disease control strategies established for your area.

To delay fungicide/bactericide resistance, take the following steps:

- Rotate the use of Tide USA TEBU 3.6F Fungicide or other Group 3 fungicides/bactericides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide/bactericides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide/bactericide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.

- Where possible, make use of predictive disease models to effectively time fungicide/bactericide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal/bacterial populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM guidance for specific crops and pathogens.
- For more information or to report suspected resistance contact Tide International USA, Inc. You can also contact your pesticide distributor or university extension specialist to report resistance.

DISEASE CONTROL IN CROPS

Crop	Diseases	Rate in fl. oz. of Tide USA TEBU 3.6F/Acre (lbs. ai/Acre)	Use Directions
Almond	Brown rot blossom blight (<i>Monilinia laxa</i> , <i>M. fructicola</i>)	8 (0.22)	Begin application at pink bud. If the bloom period is extended, and/or severe disease conditions exist, make a second application at full bloom. If conditions remain favorable for disease, make another application at petal fall. Make applications on a 7-14 day interval. Begin applications when conditions are favorable for disease but before infection. Apply Tide USA TEBU 3.6F Fungicide in a minimum spray volume of 15 gallons per acre by air or 50 gallons per acre by ground. Reduce the application interval for varieties that are highly susceptible to the indicated diseases or when severe disease conditions exist. The use of ground application after petal fall is preferred because of difficulty in penetrating the canopy and obtaining thorough coverage of the foliage and fruit by air.

Application Directions: Tide USA TEBU 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide. It may be applied in a tank mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy. Tank mixing Tide USA TEBU 3.6F Fungicide with other DMI fungicides is not advised.

RESTRICTIONS:

- **DO NOT** apply more than 8 fl. oz. (0.22 lb. ai)/acre per application.
- **DO NOT** apply more than 32 fl. oz. (0.9 lb. ai)/acre per year.
- **DO NOT** cut cover crops in treated areas for feed, or allow livestock to graze treated areas.
- **DO NOT** make more than 4 applications per year.
- Minimum retreatment interval (RTI) = 7 days
- Restricted-entry interval (REI) = 12 hours
- Pre-harvest interval (PHI) = 35 days

Crop	Diseases	Rate in fl. oz. of Tide USA TEBU 3.6F/Acre (lbs. ai/Acre)	Use Directions
Asparagus	Rusts (<i>Puccinia</i> spp.)	4 – 6 (0.11 – 0.17)	Apply as a foliar spray to the developing ferns after harvest of spears is completed. Apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development. Apply the specified amount (0.11 to 0.17 lb. of active ingredient per acre) in alternation with another effective fungicide. Under conditions of severe rust pressure, use the higher rate. Repeat applications on a 14- day interval as necessary to maintain control of rust.
<p>Application Directions: Applications may be made using ground or aerial application equipment. For optimum disease control, tank mix Tide USA TEBU 3.6F Fungicide with the lowest labeled rate of a spray surfactant. Tide USA TEBU 3.6F Fungicide is a sterol demethylation inhibitor (DMI) fungicide (Group 3). Alternating Tide USA TEBU 3.6F Fungicide with other DMI fungicides may lead to resistance.</p> <p>RESTRICTIONS:</p> <ul style="list-style-type: none"> ● DO NOT apply more than 6 fl. oz. (0.17 lb ai)/acre per application. ● DO NOT apply more than 18 fl. oz. (0.51 lb. ai)/acre per year. ● DO NOT make more than three foliar applications per year. ● DO NOT apply to harvestable spears. ● A 50 feet spray drift buffer zone is required for all aerial applications. ● Minimum retreatment interval (RTI) = 14 days ● Restricted-entry interval (REI) = 12 hours ● Pre-harvest interval (PHI) =100 days (California); 180 days (all other states) 			
Barley	Rusts (<i>Puccinia</i> spp.) Head Blight (<i>Fusarium</i> spp.)	4 (0.11)	Apply Tide USA TEBU 3.6F Fungicide in a minimum of 10 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by air. Straw cut after harvest may be fed or used for bedding. Observe barley fields closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.
<p>Application Timing of Tide USA TEBU 3.6F Fungicide for Optimum Control: Rusts: Apply Tide USA TEBU 3.6F Fungicide at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of Tide USA TEBU 3.6F Fungicide for Fusarium head blight suppression is when stem heads have fully emerged (Feekes 10.5) on 50 % of the plants.</p>			
<p>Application Directions: For optimum disease control, tank mix Tide USA TEBU 3.6F Fungicide with the lowest labeled rate of a spray surfactant. Tide USA TEBU 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tide USA TEBU 3.6F Fungicide will be resistant to weathering. Tide USA TEBU 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>RESTRICTIONS:</p> <ul style="list-style-type: none"> ● DO NOT apply more than 4 fl. oz. (0.11 lb. ai)/acre per application. ● DO NOT apply more than 4 fl. oz. (0.11 lb. ai)/acre per year. ● DO NOT make more than one application per year. ● Grazing livestock or feeding of green forage is permitted 6 or more days after the last application of Tide USA TEBU 3.6F Fungicide. ● Restricted-entry interval (REI) = 12 hours ● Pre-harvest Interval (PHI) = 30 days 			

Crop	Diseases	Rate in fl. oz. of Tide USA TEBU 3.6F/Acre (lbs. ai/Acre)	Use Directions
Beans (fresh & dry, except succulent shelled)	Rust <i>(Uromyces appendiculatus)</i>	4 – 6 (0.11 – 0.17)	Apply Tide USA TEBU 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 14-day intervals, or as necessary to maintain control.
<p>Application Directions: For optimum disease control, tank mix the lowest labeled rate of a spray surfactant with Tide USA TEBU 3.6F Fungicide. Tide USA TEBU 3.6F Fungicide must have two to four hours of drying time on bean foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tide USA TEBU 3.6F Fungicide will be resistant to weathering. Tide USA TEBU 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>RESTRICTIONS:</p> <ul style="list-style-type: none"> ● DO NOT apply more than 6 fl. oz. (0.17 lb. ai)/acre per application. ● DO NOT apply more than 24 fl. oz. (0.68 lb. ai)/acre per year (fresh beans); DO NOT apply more than 12 fl. oz. (0.34 lb. ai)/acre per year (dry beans). ● DO NOT make more than 6 applications per year when applying at the low rate (fresh beans); DO NOT make more than 3 applications per year when applying at the low rate (dry beans). ● Restricted-entry interval (REI) = 12 hours ● Minimum retreatment interval (RTI) = 14 days ● Pre-harvest interval (PHI) = 7 days (fresh beans); 14 days (dry beans) 			
Corn* (sweet corn, field corn, field corn grown for seed, and popcorn) *Not approved for use in New York.	Rust <i>(Puccinia spp.)</i> Northern Leaf Blight <i>(Helminthosporium turcicum)</i> Southern Leaf Blight <i>(Helminthosporium maydis)</i> Northern Leaf Spot <i>(Helminthosporium carbonum)</i> Gray Leaf Spot <i>(Cercospora zeaemaydis)</i>	4 – 6 (0.11 – 0.17)	Apply Tide USA TEBU 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for disease development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control.
<p>Application Directions: For optimum disease control, tank-mix the lowest labeled rate of a spray surfactant with Tide USA TEBU 3.6F Fungicide. Tide USA TEBU 3.6F Fungicide must have two to four hours of drying time on corn foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tide USA TEBU 3.6F Fungicide will be resistant to weathering. Tide USA TEBU 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>RESTRICTIONS:</p> <ul style="list-style-type: none"> ● DO NOT apply more than 6 fl. oz. (0.17 lb. ai)/acre per application. ● DO NOT apply more than 24 fl. oz. (0.68 lb. ai)/acre per year. ● DO NOT make more than 6 applications per year when applying at the low rate. ● Minimum retreatment interval (RTI) = 7 days ● Restricted-entry interval (REI) for sweet corn = 19 days ● Pre-harvest interval (PHI) for sweet corn = 7 days (ears or forage); 49 days (fodder) ● Restricted-entry interval (REI) for all corn except sweet corn = 12 hours ● Pre-harvest interval (PHI) for field, seed or popcorn = 21 days (forage); 36 days (grain or fodder) 			

Crop	Diseases	Rate in fl. oz. of Tide USA TEBU 3.6F/Acre (lbs. ai/Acre)	Use Directions
<p>COTTON</p> <p>COTTONSEED SUBGROUP 20C*</p> <p>Cottonseed: Cultivars, varieties and/or hybrids of these</p> <p>*Not approved for use in California.</p>	<p>Southwestern Cotton Rust (<i>Puccinia cacabata</i>)</p>	<p>6 – 8 (0.17 – 0.22)</p>	<p>Apply Tide USA TEBU 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control.</p>
<p>Application Directions: For optimum disease control, tank-mix the lowest labeled rate of a spray surfactant with Tide USA TEBU 3.6F Fungicide. Tide USA TEBU 3.6F Fungicide must have two to four hours of drying time on cotton foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tide USA TEBU 3.6F Fungicide will be resistant to weathering. Tide USA TEBU 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>RESTRICTIONS:</p> <ul style="list-style-type: none"> ● DO NOT apply more than 8 fl. oz. (0.22 lb. ai)/acre per application. ● DO NOT apply more than 24 fl. oz. (0.68 lb. ai)/acre per year. ● DO NOT make more than 4 applications per year when applying at the low rate. ● Minimum retreatment interval (RTI) = 7 days ● Restricted-entry interval (REI) = 12 hours ● Pre-harvest interval (PHI) = 30 days 			
<p>CUCURBIT VEGETABLES GROUP:</p> <p>Chayote, Chinese Waxgourd, Citron Melon, Cucumber, Gherkin, Edible Gourd (includes Hyotan, Cucuzza, Hechima, and Chinese Okra), Momordica spp. (includes Balsam Apple, Balsam Pear, Bitter Melon and Chinese Cucumber), Muskmelon (includes Cantaloupe, Casaba, Crenshaw Melon, Golden Pershaw Melon, Honeydew Melon,</p>	<p>Powdery Mildew (<i>Sphaerotheca fuliginea/Podosphaera xanthii</i>) (<i>Erysiphe cichoracearum</i>)</p>	<p>4 – 6 (0.11 – 0.17)</p>	<p>Apply specified dosage in a protective spray schedule to foliage and fruit. Repeat applications at 10- to 14-day intervals.</p>
	<p>Gummy Stem Blight – suppression (<i>Didymella bryoniae</i>) (watermelon, squash, pumpkin, and melons only)</p>	<p>8 (0.22)</p>	<p>Apply specified dosage in a protective spray schedule to foliage and fruit. Repeat applications at 10- to 14-day intervals.</p>

Crop	Diseases	Rate in fl. oz. of Tide USA TEBU 3.6F/Acre (lbs. ai/Acre)	Use Directions
[cont'd] Honey Balls, Mango Melon, Persian Melon, Pineapple Melon, Santa Clause Melon and Snake Melon), Pumpkin Summer Squash (includes Crookneck Squash, Scallop Squash, Straightneck Squash, Vegetable Marrow and Zucchini), Winter Squash (includes Butternut Squash, Calabaza, Hubbard Squash, Acorn Squash and Spaghetti Squash), Watermelon	Powdery Mildew (<i>Sphaerotheca fuliginea</i> / <i>Podosphaera xanthii</i>) (<i>Erysiphe cichoracearum</i>)	4 – 6 (0.11 – 0.17)	Apply specified dosage in a protective spray schedule to foliage and fruit. Repeat applications at 10- to 14-day intervals.
	Gummy Stem Blight – suppression (<i>Didymella bryonae</i>) (watermelon, squash, pumpkin, and melons only)	8 (0.22)	Apply specified dosage in a protective spray schedule to foliage and fruit. Repeat applications at 10- to 14-day intervals.

Application Directions: For optimum disease control, tank-mix the lowest labeled rate of a spray surfactant with Tide USA TEBU 3.6F Fungicide. Tide USA TEBU 3.6F Fungicide must have two to four hours of drying time for the active ingredient to move systematically into plant tissue before rain or irrigation occurs. After this period of time, Tide USA TEBU 3.6F Fungicide will be resistant to weathering. Tide USA TEBU 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

RESTRICTIONS:

- **DO NOT** apply more than 8 fl. oz. (0.22 lb. ai)/acre per application.
- **DO NOT** apply more than 24 fl. oz. (0.68 lb. ai)/acre per year.
- **DO NOT** make more than 6 applications per year when applying at the low rate.
- Minimum retreatment interval (RTI) = 10 days
- Restricted-entry interval (REI) = 12 hours
- Pre-harvest interval (PHI) = 7 days

Crop	Diseases	Rate in fl. oz. of Tide USA TEBU 3.6F/Acre (lbs. ai/Acre)	Use Directions
<p>BULB VEGETABLE CROP, SUBGROUP 3-07A</p> <p>Daylily Bulb*† Fritillaria Bulb*† Dry Bulb Onion Garlic Great-headed Garlic Bulb Serpent Garlic Bulb*† Lily Bulb*† Chinese Onion Bulb*† Pearl Onion*† Potato Onion Bulb*† Shallot Cultivars, varieties, and/or hybrids of these*†</p> <p>*Not approved for use in New York. †Not approved for use in California.</p>	<p>White Rot <i>(Sclerotium cepivorum)</i></p> <p>Rust <i>(Puccinia allii, Puccinia porri)</i></p> <p>Purple Blotch <i>(Alternaria porri)</i></p>	<p>White rot: 20.5 fl. oz. (0.58 lb. ai) per acre applied in a 4 to 6 inch band over/into each furrow</p> <p>4 – 6 (0.11 – 0.17)</p>	<p>White Rot: For the control of white rot, make one application in the furrow at the time of planting. Make the in-furrow application at the rate of 20.5 fl. oz. Tide USA TEBU 3.6F Fungicide per acre. Apply the entire per acre rate in a 4 to 6 inch band over/into each furrow. Additional control may be obtained by including two foliar applications at 4 – 6 fl. oz./acre. Application by chemigation is permitted for control of white rot.</p> <p>Rust: For the control of rust, make foliar applications at the rate of 4 - 6 fl. oz./acre per application. Repeat at an interval of 10 - 14 days. Apply Tide USA TEBU 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for rust development.</p>

Application Directions: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest specified rate of a spray surfactant may be tank-mixed with Tide USA TEBU 3.6F Fungicide. Tide USA TEBU 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tide USA TEBU 3.6F Fungicide will be resistant to weathering. Tide USA TEBU 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

RESTRICTIONS:

- **DO NOT** apply more than 6 fl. oz. (0.17 lb. ai)/acre per application unless applying for White rot in which case **DO NOT** apply more than 20.5 fl. oz. (0.58 lb. ai)/acre per application.
- **DO NOT** apply more than 32.5 fl. oz. (0.91 lb. ai)/acre per year if an in-furrow treatment is made. If not applied as an in-furrow treatment, then **DO NOT** apply more than 12 fl. oz. (0.34 lb. ai)/acre per year as a foliar spray.
- **DO NOT** make more than 1 in-furrow application per year. If an in-furrow application is made, up to 2 additional foliar applications per year may be made. If no in-furrow application is made, **DO NOT** make more than 3 applications per year when applying at the low rate.
- Minimum retreatment interval (RTI) = 10 days
- Restricted-entry interval (REI) = 12 hours
- Pre-harvest interval (PHI) = 7 days

Crop	Diseases	Rate in fl. oz. of Tide USA TEBU 3.6F/Acre (lbs. ai/Acre)	Use Directions
<p>BULB VEGETABLE CROP, SUBGROUP 3-07B</p> <p>Chive*† Chinese Chive*† Elegans Hosta*† Fritillaria*† Kurrat*† Lady's Leek*† Wild Leek*† Beltsville Bunching Onions*† Fresh Onion*† Green Onion Green Eschalots Shallot Fresh Leaves*† Japanese Bunching Onion Leek Scallion Spring Onion Macrostem Onion*† Tree Onion Tops*† Welsh Onion Tops*† Cultivars, varieties, and/or hybrids of these*†</p> <p>*Not approved for use in New York. †Not approved for use in California.</p>	<p>White Rot (<i>Sclerotium cepivorum</i>) (suppression only)</p> <p>Rust (<i>Puccinia allii</i>, <i>Puccinia porri</i>)</p> <p>Purple Blotch (<i>Alternaria porri</i>)</p>	<p>4 – 6 (0.11-0.17)</p>	<p>For the control of diseases, make foliar applications using an interval of 10 - 14 days. Apply Tide USA TEBU 3.6F Fungicide in a protective spray schedule or when weather conditions are favorable for rust development.</p>

Application Directions: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest specified rate of a spray surfactant may be tank-mixed with Tide USA TEBU 3.6F Fungicide. Tide USA TEBU 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tide USA TEBU 3.6F Fungicide will be resistant to weathering. Tide USA TEBU 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

RESTRICTIONS:

- DO NOT apply more than 6 fl. oz. (0.17 lb. ai)/acre per application.
- DO NOT apply more than 24 fl. oz. (0.68 lb. ai)/acre per year.
- DO NOT make more than 6 applications per year when applying at the low rate.
- Minimum retreatment interval (RTI) = 10 days
- Restricted-entry interval (REI) = 12 hours
- Pre-harvest interval (PHI) = 7 days

Crop	Diseases	Rate in fl. oz. of Tide USA TEBU 3.6F/Acre (lbs. ai/Acre)	Use Directions
<p>FILBERT (HAZELNUT)*</p> <p>*Not approved for use in California</p>	<p>Eastern filbert blight (<i>Anisogramma anomala</i>)</p>	<p>8 (0.22)</p>	<p>Apply Tide USA TEBU 3.6F Fungicide in a preventive spray schedule when conditions are favorable for disease (including budbreak when the first green tissue is visible). Continue applications of Tide USA TEBU 3.6F Fungicide at 7- to 14-day intervals if weather conditions are conducive to disease development. Use shorter spray intervals during budbreak and rapid shoot elongation. The expanding shoot tip must be protected. Reduce the application interval for varieties that are highly susceptible to the indicated disease or when severe disease conditions exist.</p> <p>Apply Tide USA TEBU 3.6F Fungicide in sufficient spray volume for thorough coverage. Tank mixing the lowest labeled rate of a spray surfactant with Tide USA TEBU 3.6F Fungicide may improve spray coverage and penetration of the active ingredient into plant tissue. The use of ground application is preferred because of the difficulty in penetrating the canopy and obtaining thorough coverage of the foliage and stems by air.</p>
<p>Application Directions: Tide USA TEBU 3.6F Fungicide is a sterol demethylation inhibitor (DMI) fungicide. It may be applied in tank mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy. Tank mixing Tide USA TEBU 3.6F Fungicide with other DMI fungicides is not advised.</p> <p>RESTRICTIONS:</p> <ul style="list-style-type: none"> ● DO NOT apply more than 8 fl. oz. (0.22 lb. ai)/acre per application. ● DO NOT apply more than 32 fl. oz. (0.9 lb. ai)/acre per year. ● DO NOT make more than 4 applications per year. ● DO NOT cut cover crops in treated areas for feed, or allow livestock to graze treated areas. ● Minimum retreatment interval (RTI) = 7 days ● Restricted-entry interval (REI) = 12 hours ● Pre-harvest interval (PHI) = 35 days 			
<p>FRUITING VEGETABLES GROUP (EXCEPT OKRA)*</p> <p>African eggplant Bush tomato Bell pepper Cocona</p>	<p>Early blight (<i>Alternaria solani</i>)</p>	<p>8 (0.22)</p>	<p>Make applications on 7-day intervals.</p>

Crop	Diseases	Rate in fl. oz. of Tide USA TEBU 3.6F/Acre (lbs. ai/Acre)	Use Directions
[cont'd] Currant tomato Eggplant Garden huckleberry Goji berry Ground cherry Martynia Naranjila Pea eggplant Pepino Nonbell Pepper Roselle Scarlet eggplant Sunberry Tomatillo Tomato Tree tomato Cultivars, varieties, and/or hybrids of these *Not registered for this use in California			
<p>Application Directions: For optimum results, use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest labeled rate of a spray surfactant may be tank-mixed with Tide USA TEBU 3.6F Fungicide. Tide USA TEBU 3.6F Fungicide must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tide USA TEBU 3.6F Fungicide will be resistant to weathering. Tide USA TEBU 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>RESTRICTIONS:</p> <ul style="list-style-type: none"> ● DO NOT apply more than 8 fl. oz. (0.22 lb. ai)/acre per application. ● DO NOT apply more than 48 fl. oz. (1.35 lbs. ai)/acre per year. ● DO NOT make more than 6 applications per year. ● Minimum retreatment interval (RTI) = 7 days ● Restricted-entry interval (REI) = 12 hours ● Pre-harvest interval (PHI) = 7 days 			
Okra	Cercospora Leaf Spot (<i>Cercospora</i> spp.)	4 – 6 (0.11 – .17)	Apply specified dosage of Tide USA TEBU 3.6F Fungicide in a preventative spray program. Use the highest rate when disease conditions are favorable and in areas where high disease pressure is expected. Applications may be repeated at 14-day intervals in order to maintain control of the disease. Apply specified dosage as a foliar spray in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air.

Crop	Diseases	Rate in fl. oz. of Tide USA TEBU 3.6F/Acre (lbs. ai/Acre)	Use Directions
<p>[cont'd]</p> <p>Application Directions: For optimum disease control, tank-mix the lowest labeled rate of spray surfactant with Tide USA TEBU 3.6F Fungicide. Tide USA TEBU 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tide USA TEBU 3.6F Fungicide will be resistant to weathering. Tide USA TEBU 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>RESTRICTIONS:</p> <ul style="list-style-type: none"> ● DO NOT apply more than 6 fl. oz. (0.17 lb. ai)/acre per application. ● DO NOT apply more than 24 fl. oz. (0.68 lb. ai)/acre per year. ● DO NOT make more than 6 applications per year when applying at the low rate. ● Minimum retreatment interval (RTI) = 14 days ● Restricted-entry interval (REI) = 12 hours ● Pre-harvest interval (PHI) = 3 days 			
Hops	Powdery Mildew (<i>Sphaerotheca humuli/Sphaerotheca maculaha</i> s)	4 – 8 (0.11 – 0.22)	Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 10- to 14-day intervals. Increase the spray volume and the application rate within specified rate range as vine growth increases during the season.
<p>Application Directions: For optimum disease control, tank-mix the lowest labeled rate of a spray surfactant with Tide USA TEBU 3.6F Fungicide. Tide USA TEBU 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tide USA TEBU 3.6F Fungicide will be resistant to weathering. Tide USA TEBU 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>RESTRICTIONS:</p> <ul style="list-style-type: none"> ● DO NOT apply more than 8 fl. oz. (0.22 lb. ai)/acre per application. ● DO NOT apply more than 32 fl. oz. (0.9 lb. ai)/acre per year. ● DO NOT make more than 8 applications per year when applying at the low rate. ● Minimum retreatment interval (RTI) = 10 days ● Restricted-entry interval (REI) = 12 hours ● Pre-harvest interval (PHI) = 14 days 			
LEAFY BRASSICA GREENS SUBGROUP 4-16B (except watercress) Arugula†, Chinese Broccoli†, Broccoli Raab, Abyssinian Cabbage†, Chinese Cabbage (Bok Choy), Seakale Cabbage†, Collards, Garden Cress†,	Cercospora Leaf Spot (<i>Cercospora brassicicola</i>) Powdery Mildew (<i>Erysiphe cruciferarum</i>) Alternaria Leaf Spot (<i>Alternaria brassicicola</i>)	3 – 4 (0.08 – 0.11)	

Crop	Diseases	Rate in fl. oz. of Tide USA TEBU 3.6F/Acre (lbs. ai/Acre)	Use Directions
[cont'd] Upland Cress†, Hanover Salad†, Kale, Maca Leaves†, Mizuna, Mustard Greens, Radish Leaves†, Rape Greens, Wild Rocket†, Shepherd's Purset†, Turnip Greens, Cultivars, varieties, and hybrids of these commodities† †Not approved for use in California.	Cercospora Leaf Spot <i>(Cercospora brassicicola)</i> Powdery Mildew <i>(Erysiphe cruciferarum)</i> Alternaria Leaf Spot <i>(Alternaria brassicicola)</i>	3 – 4 (0.08 – 0.11)	

Application Directions: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest specified rate of a spray surfactant may be tank-mixed with Tide USA TEBU 3.6F Fungicide. Tide USA TEBU 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tide USA TEBU 3.6F Fungicide will be resistant to weathering. Tide USA TEBU 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).

RESTRICTIONS:

- **DO NOT** apply more than 4 fl. oz. (0.11 lb. ai)/acre per application.
- **DO NOT** apply more than 16 fl. oz. (0.45 lb. ai)/acre per year.
- **DO NOT** make more than 5 applications per year when applying at the low rate.
- Minimum retreatment interval (RTI) = 10 days
- Restricted-entry interval (REI) = 12 hours
- Pre-harvest interval (PHI) = 7 days
- Application to turnip greens is limited to east of the Rockies.

Garden Beet roots and tops (leaves)	Cercospora Leaf Spot <i>(Cercospora beticola)</i>	3 – 7.2 (0.08 - 0.20)	Make applications on 14-day intervals.
<p>Application Directions: For optimum results, use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest labeled rate of a spray surfactant may be tank-mixed with Tide USA TEBU 3.6F Fungicide. Tide USA TEBU 3.6F Fungicide must have two to four hours of drying time on beet foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tide USA TEBU 3.6F Fungicide will be resistant to weathering. Tide USA TEBU 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>RESTRICTIONS:</p> <ul style="list-style-type: none"> ● DO NOT apply more than 7.2 fl. oz. (0.20 lb. ai)/acre per application. ● DO NOT apply more than 28.8 fl. oz. (0.81 lb. ai)/acre per year. ● DO NOT make more than 9 applications per year when applying at the low rate. ● Minimum retreatment interval (RTI) = 14 days ● Restricted-entry interval (REI) = 12 hours ● Pre-harvest interval (PHI) = 7 days 			

Crop	Diseases	Rate in fl. oz. of Tide USA TEBU 3.6F/Acre (lbs. ai/Acre)	Use Directions
Peanut	<p>SOILBORNE:</p> <p><i>Cylindrocladium</i></p> <p>Black Rot (suppression)</p> <p>Rhizoctonia Limb Rot</p> <p>Rhizoctonia Pod Rot (Virginia and North Carolina only)</p> <p>Sclerotium Stem and Pod Rot (White Mold, Southern Blight, Southern Stem Rot)</p> <p>FOLIAR:</p> <p>Early Leaf Spot</p> <p>Late Leaf Spot</p> <p>Leaf Rust</p> <p>Pepper Spot (<i>Leptosphaerulina</i>)</p> <p>Web Blotch (<i>Phoma</i>)</p>	7.2 (0.20)	<p>FOUR-APPLICATION SPRAY PROGRAM: Apply the specified rate in a preventive spray schedule. See table below for proper timing of applications. Applications of chlorothalonil must be made prior to and following applications of Tide USA TEBU 3.6F Fungicide to discourage development of resistant strains of fungi. For optimum control of foliar diseases including leaf rust, web blotch, and pepper spot, tank mix the lowest label specified rate of a spray surfactant with Tide USA TEBU 3.6F Fungicide.</p> <p>LEAF SPOT ADVISORY SCHEDULE: For control of soilborne diseases in an advisory schedule, apply Tide USA TEBU 3.6F Fungicide in the first advisory spray in July and continue Tide USA TEBU 3.6F Fungicide applications at 14-day intervals. When applying Tide USA TEBU 3.6F Fungicide after August 15, tank mix with chlorothalonil for resistance management purposes.</p>

Application Directions: For optimum control of the specified soilborne diseases, four consecutive applications of Tide USA TEBU 3.6F Fungicide must be made at 14-day intervals. Tide USA TEBU 3.6F Fungicide is a sterol demethylation inhibitor (DMI) fungicide. Chlorothalonil may be tank mixed at the rate of 12 ounces of active ingredient with Tide USA TEBU 3.6F Fungicide as a leaf spot resistance management strategy. A spray surfactant is not necessary when Tide USA TEBU 3.6F Fungicide is tank mixed with chlorothalonil. Mixing or alternating Tide USA TEBU 3.6F Fungicide with other DMI fungicides may lead to resistance.

Tide USA TEBU 3.6F Fungicide must be carried by rainfall or irrigation into the root and pod zone for control of root and pod rots caused by *Sclerotium rolfsii* and *Rhizoctonia solani*. Drought conditions will decrease the effectiveness of Tide USA TEBU 3.6F Fungicide against the root and pod rots.

Use Tide USA TEBU 3.6F Fungicide in conjunction with cultural practices that are known to reduce the severity of soilborne diseases, including proper crop rotation practices.

RESTRICTIONS:

- **DO NOT** feed hay or threshings or allow livestock to graze in treated areas.
- **DO NOT** apply more than 7.2 fl. oz. (0.20 lb. ai)/acre per application.
- **DO NOT** apply more than 28.8 fl. oz. (0.81 lb. ai)/acre per year.
- **DO NOT** make more than 4 applications per year.
- Minimum retreatment interval (RTI) = 14 days
- Restricted-entry interval (REI) = 12 hours
- Pre-harvest interval (PHI) = 14 days

Application Timing of Tide USA TEBU 3.6F Fungicide for Optimum Control of White Mold and Rhizoctonia Limb and Pod Rot

Spraying Program	Tide USA TEBU 3.6F Application No.	Chlorothalonil Application No.
7 applications	3, 4, 5 and 6	1, 2, and 7

Crop	Diseases	Rate in fl. oz. of Tide USA TEBU 3.6F/Acre (lbs. ai/Acre)	Use Directions
Pecan	Brown Leaf Spot <i>(Sirosporium diffusium)</i> Downy Spot <i>(Mycosphaerella caryigena)</i> Liver Spot <i>(Gnomonia caryae)</i> Scab <i>(Cladosporium caryigenum)</i> Vein Spot <i>(Gnomonia nerviseda)</i> Zonate Leaf Spot <i>(Grovesinia pyramidalis)</i>	4 – 8 (0.11 – 0.22)	<p>Apply Tide USA TEBU 3.6F Fungicide in a preventative spray schedule beginning at early bud break (young leaves unfolding), and continue applications at 10- to 14-day intervals through the pollination period. Tide USA TEBU 3.6F Fungicide must be applied at 4 fl. oz./acre in a tank-mix with the specified rate of Super-Tin® in cover sprays. Follow label directions for the use of Super-Tin.</p> <p>Apply Tide USA TEBU 3.6F Fungicide in a spray volume of 15 or more gallons per acre by air or 50 or more gallons per acre by ground. Apply 7 - 8 fl. oz./acre of Tide USA TEBU 3.6F Fungicide to fullsize mature trees, and 4 - 6 fl. oz./acre to smaller trees. Apply the high rate to varieties that are highly susceptible to the indicated diseases, or when severe disease conditions exist. The lowest labeled rate of a surfactant may be added to the spray solution for optimum control of the indicated diseases.</p>

Application Directions: For optimum disease control, tank-mix the lowest labeled rate of a spray surfactant with Tide USA TEBU 3.6F Fungicide. Tide USA TEBU 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tide USA TEBU 3.6F Fungicide will be resistant to weathering. Tide USA TEBU 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy.

RESTRICTIONS:

- **DO NOT** add a surfactant to the spray solution when tank-mixing Tide USA TEBU 3.6F Fungicide with Super-Tin.
- **DO NOT** cut cover crops in treated areas for feed or allow livestock to graze treated areas.
- **DO NOT** apply more than 8 fl. oz. (0.22 lb. ai)/acre per application.
- **DO NOT** apply more than 32 fl. oz. (0.9 lb. ai)/acre per year.
- **DO NOT** make more than 8 applications per year when applying at the low rate.
- Minimum retreatment interval (RTI) = 10 days
- Restricted-entry interval (REI) = 12 hours
- Pre-harvest interval (PHI) = **DO NOT** apply after shucks begin to split

Crop	Diseases	Rate in fl. oz. of Tide USA TEBU 3.6F/Acre (lbs. ai/Acre)	Use Directions
Pistachio* *Not approved for use in California	Botryosphaeria panicle and shoot blight <i>(Botryosphaeria dothidea)</i>	8 (0.22)	Begin application at 25 to 50% bloom and repeat again 10 to 14 days later to protect young flower clusters and fruit. Make two additional applications of Tide USA TEBU 3.6F 14 days apart beginning 49 days before harvest.* Apply Tide USA TEBU 3.6F in a minimum spray volume of 15 gallons per acre by air or 50 gallons per acre by ground. Reduce the application interval for varieties that are highly susceptible to the indicated diseases or when severe disease conditions exist. The use of ground application after flowering is preferred because of difficulty in penetrating the canopy and obtaining thorough coverage of the foliage and fruit by air. Control of insect vectors and a thorough pruning program to remove plant tissue infected by Botryosphaeria are critical for optimum control of this disease.
<p>Application Directions: Tide USA TEBU 3.6F is a demethylation inhibitor (DMI) fungicide. It may be applied in a tank mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy. Tank mixing Tide USA TEBU 3.6F with other DMI fungicides is not advised.</p> <p>RESTRICTIONS:</p> <ul style="list-style-type: none"> ● DO NOT cut cover crops in treated areas for feed or allow livestock to graze treated areas. ● DO NOT apply more than 8 fl. oz. (0.22 lb. ai)/acre per application. ● DO NOT apply more than 32 fl. oz. (0.9 lb. ai)/acre per year. ● DO NOT make more than 4 applications per year. ● Minimum retreatment interval (RTI) = 10 days ● Restricted-entry interval (REI) = 12 hours ● Pre-harvest interval (PHI) = 35 days 			
Soybean	Rust <i>(Phakopsora pachyrhizi)</i> Powdery Mildew <i>(Microsphaera diffusa)</i>	3 – 4 (0.08 – 0.11)	Apply Tide USA TEBU 3.6F Fungicide as a broadcast foliar spray as a preventative spray or at first visible symptoms of disease. Repeat applications on a 10- to 14-day spray interval if environmental conditions are favorable for continued disease development. Use of the higher rates and shorter spray intervals are advised when disease pressure is severe. The lowest label specified rate of a spray surfactant may be tank-mixed with Tide USA TEBU 3.6F Fungicide. Tide USA TEBU 3.6F Fungicide must be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment.

Crop	Diseases	Rate in fl. oz. of Tide USA TEBU 3.6F/Acre (lbs. ai/Acre)	Use Directions
RESTRICTIONS: <ul style="list-style-type: none"> ● DO NOT apply more than 4 fl. oz. (0.11 lb. ai)/acre per application. ● DO NOT apply more than 12 fl. oz. (0.34 lb. ai)/acre per year. ● DO NOT make more than three applications per year. ● Minimum retreatment interval (RTI) = 10 days ● Restricted-entry interval (REI) = 12 hours ● Pre-harvest interval (PHI) = 21 days 			
Sunflower	Rust (<i>Puccinia helianthi</i>)	4 – 6 (0.11 – 0.17)	Apply specified dosage of Tide USA TEBU 3.6F Fungicide at the earliest sign of infection (rust pustules developing) or when weather conditions are favorable for rust development. Apply higher rate to highly susceptible varieties and/or under severe disease conditions. Application may be repeated at 14 days if necessary to maintain control of the disease. Apply specified dosage in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air.
Application Directions: For optimum disease control, tank-mix the lowest labeled rate of a spray surfactant with Tide USA TEBU 3.6F Fungicide. Contact your state extension service for a list of approved surfactants. Tide USA TEBU 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tide USA TEBU 3.6F Fungicide will be resistant to weathering. Tide USA TEBU 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). RESTRICTIONS: <ul style="list-style-type: none"> ● DO NOT apply more than 6 fl. oz. (0.17 lb. ai)/acre per application. ● DO NOT apply more than 16 fl. oz. (0.45 lb. ai)/acre per year. ● DO NOT make more than 4 applications per year when applying at the low rate. ● Minimum retreatment interval (RTI) = 14 days ● Restricted-entry interval (REI) = 12 hours ● Pre-harvest interval (PHI) = 50 days 			
Tomato (Greenhouse)* *Not approved for use in California	Powdery mildew (<i>Oidiopsis sicula</i> , <i>Oidium neolycopersici</i> , <i>Oidium lycopersici</i>)	8 (0.22)	A total of 6 applications can be made at minimum application intervals of 7 days, with the last application on the day of harvest. For ground applications, apply a minimum of 10 gallons of spray volume per acre, increasing the spray volume as plants mature to ensure thorough coverage of the foliage.
RESTRICTIONS: <ul style="list-style-type: none"> ● DO NOT apply more than 8 fl. oz. (0.22 lb. ai)/acre per application. ● DO NOT apply more than 48 fl. oz. (1.35 lbs. ai)/acre per year. ● DO NOT exceed 6 applications per year. ● Minimum retreatment interval (RTI) = 7 days ● Restricted-entry interval (REI) = 12 hours. If harvesting is initiated before 12 hours, PPE is required. ● Pre-harvest interval (PHI) = 0 days 			

Crop	Diseases	Rate in fl. oz. of Tide USA TEBU 3.6F/Acre (lbs. ai/Acre)	Use Directions
<p>TREE NUT GROUP 14-12* (See separate use direction sections for Pecan, Almond, Pistachio and Filbert)</p> <p>African nut-tree, Beechnut, Brazil nut, Brazilian pine, Bunya, Bur oak, Butternut, Cajou nut, Candlenut, Cashew, Chestnut, Chinquapin, Coconut, Coquito nut, Dika nut, Ginkgo, Guiana chestnut, Heartnut, Hickory nut, Japanese horse-chestnut, Macadamia nut, Mongongo nut, Monkey-pot, Monkey puzzle nut, Okari nut, Pachira nut, Peach palm nut, Pequi, Pili nut, Pine nut, Sapucaia nut, Tropical almond, Walnut, black, Walnut, English, Yellowhorn, Cultivars, varieties, and/or hybrids of these</p> <p>*Not approved for use in California</p>	<p>Brown rot blossom blight (<i>Monilinia taxa</i>, <i>M. fructicola</i>)</p> <p>Botryosphaeria panicle and shoot blight (<i>Botryosphaeria dothidea</i>)</p> <p>Eastern filbert blight (<i>Anisogramma anomala</i>)</p>	<p>8 (0.22)</p>	<p>Begin applications when conditions are favorable for disease but before infection. Apply on a 7- to 14-day spray schedule.</p>

Application Directions: Tide USA TEBU 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide. It may be applied in a tank mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy. Tank mixing Tide USA TEBU 3.6F Fungicide with other DMI fungicides is not advised.

RESTRICTIONS:

- **DO NOT** apply more than 8 fl. oz. (0.22 lb. ai)/acre per application.
- **DO NOT** apply more than 32 fl. oz. (0.9 lb. ai)/acre per year.
- **DO NOT** make more than 4 applications per year.
- **DO NOT** cut cover crops in treated areas for feed, or allow livestock to graze treated areas.
- Minimum retreatment interval (RTI) = 7 days
- Restricted-entry interval (REI) = 12 hours
- Pre-harvest interval (PHI) = 35 days

Crop	Diseases	Rate in fl. oz. of Tide USA TEBU 3.6F/Acre (lbs. ai/Acre)	Use Directions
<p>TROPICAL AND SUBTROPICAL, SMALL FRUIT, INEDIBLE PEEL, SUBGROUP 24A*</p> <p>Aisen, Bael Fruit, Burmese Grape, Cat's-Eyes, Inga, Longan, Lychee, Madras Thorn, Manduro, Matisia, Mesquite, Mongongo Fruit, Pawpaw, Small-Flower, Satinleaf, Sierra Leone Tamarind, Spanish Lime, Velvet Tamarind, Wampi, White Star Apple, Cultivars, varieties, and hybrids of these, commodities</p> <p>*Not approved for use in California or New York</p>	<p>Anthracnose (<i>Colletotrichum gloeosporioides</i>)</p>	<p>4 – 6 (0.11 – 0.17)</p>	<p>Begin first application of Tide USA TEBU 3.6F Fungicide as panicle emerges. Spray up to 6 fl. oz./A every 10 days thereafter for a total of 8 sprays. Apply specified dosage in a minimum of 50 gallons of spray solution per acre by ground only.</p>
<p>RESTRICTIONS:</p> <ul style="list-style-type: none"> ● DO NOT apply more than 6 fl. oz. (0.17 lb. ai)/acre per application. ● DO NOT apply more than 48 fl. oz. (1.35 lbs. ai)/acre per year. ● DO NOT make more than 12 applications per year when applying at the low rate. ● Minimum retreatment interval (RTI) = 10 days ● Restricted-entry interval (REI) = 12 hours, if harvesting is initiated before 12 hours, PPE is required. ● Pre-harvest interval (PHI) = 0 days 			
<p>Turnip* (Application is limited to east of the Rockies)</p> <p>*Not approved for use in New York.</p>	<p>Cercospora Leaf Spot (<i>Cercospora brassicicola</i>)</p>	<p>4 – 7.2 (0.11 – 0.20)</p>	<p>Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 12- to 14-day intervals.</p>
<p>Application Directions: For optimum disease control, tank-mix the lowest labeled rate of a spray surfactant with Tide USA TEBU 3.6F Fungicide. Tide USA TEBU 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tide USA TEBU 3.6F Fungicide will be resistant to weathering. Tide USA TEBU 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>RESTRICTIONS:</p> <ul style="list-style-type: none"> ● DO NOT apply more than 7.2 fl. oz. (0.20 lb. ai)/acre per application. ● DO NOT apply more than 28.8 fl. oz. (0.81 lb. ai)/acre per year. ● DO NOT make more than 7 applications per year when applying at the low rate. ● Minimum retreatment interval (RTI) = 12 days ● Restricted-entry interval (REI) = 12 hours ● Pre-harvest interval (PHI) = 7 days 			

Crop	Diseases	Rate in fl. oz. of Tide USA TEBU 3.6F/Acre (lbs. ai/Acre)	Use Directions
Watercress* *Not approved for use in California	Cercospora leaf spot <i>(Cercospora armoraciae)</i>	7.0 (0.20)	Apply as a foliar broadcast or chemigation (via overhead irrigation) application. For the first harvest, a total of 2 foliar broadcast or chemigation (via overhead irrigation) applications can be made at a minimum application interval of 7 days. The last application must be made 3 days before the first harvest. For the second harvest, a total of 2 foliar broadcast or chemigation (via overhead irrigation) applications can be made at a minimum application interval of 7 days.
RESTRICTIONS: <ul style="list-style-type: none"> ● DO NOT apply more than 7 fl. oz. (0.2 lb. ai)/acre per application. ● DO NOT apply more than 28 fl. oz. (0.79 lb. ai)/acre per year. ● DO NOT make more than 4 applications per year. ● DO NOT apply with handheld mechanical sprayers. ● Production fields must be drained of water 24 hours prior to application and water must not be reapplied to the field for a minimum of 24 hours following the application. ● Restricted-entry interval (REI) = 12 hours ● Pre-harvest interval (PHI) = 3 days ● Minimum retreatment interval (RTI) = 7 days 			
Wheat	Rusts leaf, stem, and stripe <i>(Puccinia spp.)</i> Head blight or scab <i>(Fusarium spp.) — Suppression</i>	4.0 (0.11)	Notes: Wheat fields must be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. Straw may be fed or used for bedding. Apply Tide USA TEBU 3.6F Fungicide in a minimum of 10 gallons of spray solution per acre by ground, or in a minimum of 5 gallons of spray solution per acre by air. Application timing: Rusts: Apply Tide USA TEBU 3.6F Fungicide at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of Tide USA TEBU 3.6F Fungicide for Fusarium head blight suppression is the beginning of flowering on main stem heads (Feekes 10.51).
Application Directions: For optimum disease control, tank-mix the lowest specified rate of a spray surfactant with Tide USA TEBU 3.6F Fungicide. Tide USA TEBU 3.6F Fungicide must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Tide USA TEBU 3.6F Fungicide will be resistant to weathering. Tide USA TEBU 3.6F Fungicide is a demethylation inhibitor (DMI) fungicide (Group 3). RESTRICTIONS: <ul style="list-style-type: none"> ● DO NOT apply more than 4 fl. oz. (0.11 lb. ai)/acre per application. ● DO NOT apply more than 4 fl. oz. (0.11 lb. ai)/acre per year. ● DO NOT make more than 1 application per year. ● DO NOT allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with Tide USA TEBU 3.6F Fungicide. ● Restricted-entry interval (REI) = 12 hours ● Pre-harvest interval (PHI) = 30 days 			

Crop	Diseases	Rate in fl. oz. of Tide USA TEBU 3.6F/Acre (lbs. ai/Acre)	Use Directions
Grasses Grown for seed	Rusts (<i>Puccinia</i> spp.)	4 – 8 (0.11 – 0.22)	Apply the specified rate of Tide USA TEBU 3.6F Fungicide as soon as weather conditions are favorable for rust development or when first rust pustules are present. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl. oz./A and apply at shorter spray intervals.
	Powdery mildew	4 – 8 (0.11 – 0.22)	Apply the specified rate of Tide USA TEBU 3.6F Fungicide when powdery mildew first appears on the leaves. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl. oz./A and apply at shorter spray intervals.
<p>Application Directions: Apply the specified rate of Tide USA TEBU 3.6F Fungicide in a minimum of 20 gallons of water per acre with aircraft. Thorough coverage is important for optimum disease control. For optimum benefit, tank-mix the lowest specified rate of a spray surfactant with Tide USA TEBU 3.6F Fungicide. Chaff, screenings and straw from treated areas may be used for feed purposes. Regrowth may be grazed starting 17 days after last application.</p> <p>RESTRICTIONS:</p> <ul style="list-style-type: none"> ● DO NOT apply more than 8 fl. oz. (0.22 lb. ai)/acre per application. ● DO NOT apply more than 16 fl. oz. (0.45 lb. ai)/acre per year. ● DO NOT make more than 4 applications per year when applying at the low rate. ● DO NOT forage, cut green crop, or use seed for feed purposes. ● DO NOT graze regrowth within 17 days of last application. ● Minimum retreatment interval (RTI) = 14 days ● Restricted-entry interval (REI) = 12 hours ● Pre-harvest interval (PHI) = 4 days. 			
SEED TREATMENT Corn* (Sweet, Field, Field Corn Grown for Seed, and Popcorn)	Soilborne and Seedborne Fusarium Soilborne and Seedborne Head Smut	0.071 (0.002) 0.27 – 0.54 (0.008 – 0.015)	Apply as a seed treatment using standard slurry or mist-type seed treatment equipment. Uniform application of seed is necessary to ensure seed safety and best disease protection. Seed must be sound and well cured prior to treatment. Product must be diluted with sufficient water to ensure complete seed coverage. Consult a seed treatment specialist regarding slurry rates advised for the crop to be treated with Tide USA TEBU 3.6F Fungicide. The length of control will vary depending on the rate used.
<p>TEBUCONAZOLE Seed Bag Label Requirements:</p> <p>The Federal Seed Act requires that containers containing treated seeds shall be labeled with the following statements:</p> <ul style="list-style-type: none"> ● This seed has been treated with Tebuconazole 3.6F Fungicide, a fungicide containing tebuconazole. ● DO NOT use treated seed for feed, food, or oil purposes. 			

Crop	Diseases	Rate in fl. oz. of Tide USA TEBU 3.6F/Acre (lbs. ai/Acre)	Use Directions
<p>[cont'd]</p> <p>The US Environmental Protection Agency requires the following statements on containers containing seed treated with tebuconazole:</p> <ul style="list-style-type: none"> ● Store treated seed away from food and feedstuffs. ● DO NOT allow children, pets or livestock to have access to treated seeds. ● Wear long pants, long-sleeved shirt, and protective gloves when handling treated seed. ● Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting. ● Dispose of all excess treated seed by burying seed away from bodies of water. ● DO NOT contaminate bodies of water when disposing of planting equipment wash water. ● Dispose of seed packaging or containers in accordance with local requirements. ● Excess treated seed may be used for ethanol production if <ul style="list-style-type: none"> 1) by-products are not used for livestock feed and 2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice. <p>RESTRICTIONS:</p> <p>When using formulations that do not contain dye, to comply with 40 CFR 153.155, all seed treated with an economic poison must be colored with an EPA approved dye including one of the dyes listed in 40 CFR Section 180.910 or 180.950 to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals.</p>			

DISEASE CONTROL IN GOLF COURSE TURF, FIELD, NURSERY AND CONTAINER ORNAMENTALS AND COMMERCIAL AND RESIDENTIAL LANDSCAPES

Chemigation: **DO NOT** apply this product through any type of irrigation system.

OBSERVE THE FOLLOWING RESTRICTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS INCLUDING LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.

- **DO NOT** apply within 100 feet of aquatic areas listed above.
- **DO NOT** cultivate within 10 feet of an aquatic area to allow growth of a vegetation filter strip.
- See Spray Drift Management section for additional information.

Spray Volume: For best results, apply this product in 66-132 gallons of water per acre for turf using ground based equipment. For ornamentals, 50-300 gallons of finished spray per acre are advised depending upon equipment, plant species and plant growth stage at time of application. For the most effective results, check equipment calibration regularly. When using lower spray volumes, be sure to maintain uniform application and full crop coverage so as to ensure effective control. Increase spray volume to ensure proper application, if required.

Compatibility Test for Mix Components:

Before mixing components, always perform a compatibility jar test. For 66 gallons per acre spray volume, use 5 cups of water in a clear, clean mixing jar. For other spray volumes adjust accordingly. Only use water from the intended source at the source temperature. Add components in the sequence indicated below in Mixing Order using 3 teaspoons for each pound of dry product or 1 1/2 teaspoons for each pint of liquid product of specified label rate per acre. Always cap the jar and invert 10 cycles between component additions. When the components have all been added to the jar and fully mixed, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution must not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent and use the compatibility agent as directed on its label.

Mixing:

Continuous agitation is required during mixing. When mixing this product and water, use the specified application rates as listed for each crop on this label. Before combining any other substances with the mixture, ensure that this product is completely dispersed in the mixture.

Mixing Procedure:

1. Water. Add three-quarters of the required volume to a thoroughly clean sprayer tank.
2. Agitation. Start agitation and maintain constant agitation throughout mixing and application.
3. Inductor. If an inductor is used, rinse it thoroughly after each component has been added.
4. Products in PVA Bags. Place any product contained in water soluble PVA bags into the mixing tank. Wait until all water soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5. Water Dispersible Products. Including dry flowables (DF), wettable powders (WP), suspension concentrates (SC) or suspo-emulsions (SE).
6. Water-soluble products.
7. Emulsifiable concentrates (including oil concentrates when applicable).
8. Water soluble additives (including AMS or UAN when applicable).
9. Remaining quantity of water.

Pump Style Sprayers

1. Add the appropriate amounts of concentrate and water to the sprayer tank.
2. Close the sprayer, shake well and pressurize.
3. Adjust nozzle to a coarse spray pattern and apply.
4. Occasionally re-pressurize the sprayer if needed to maintain a good spray pattern.

DISEASE CONTROL IN GOLF COURSE TURF

TURF USE RESTRICTIONS

- For use on golf course turf only.
- **DO NOT** use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e., elementary, middle and high schools), campgrounds, churches, and theme parks.
- Not for residential use.
- Not for use on turf being grown for sale or commercial use as sod.
- **DO NOT** use clippings for animal feed.
- **DO NOT** exceed 3.6 fl. oz. of this product (0.10 lb. ai) per 1000 sq. ft. per year.
- **DO NOT** apply more than 6 applications per year.
- In New York State **DO NOT** apply more than 3 applications of 0.6 fl. oz. of Tide USA TEBU 3.6F Fungicide (0.017 lb. ai) not to exceed 1.8 fl. oz. of product (0.05 lb. ai) per 1,000 sq. ft. per year.
- In New York State, **DO NOT** exceed annual application rate of 2.2 lb. ai/acre/yr.

Product Information

For use on all golf turf applications of cool season and warm season grasses (including Bentgrasses, Bluegrasses, Fescues, Ryegrasses, St. Augustine grasses, and Zoysia) or their mixtures. This product is not phytotoxic to any of the above mentioned grasses when used in accordance with the label.

Note: Bermudagrass can be sensitive to this product under certain conditions. **DO NOT** apply consecutive applications during or just after dormancy break. Avoid applications when the temperatures are expected to exceed 85 degrees F.

Use this product for the prevention and control of the diseases mentioned in the table below. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Preventative treatments can be applied using 28 day intervals as indicated. When treating golf greens, always treat aprons and approaches. Spray uniformly over the area to be treated with properly calibrated equipment.

Apply the specified amount of this product in sufficient water for thorough coverage. A volume of 66 – 132 gallons per acre (1.5-3.0 gallons per 1,000 sq ft) is advised. Apply using properly calibrated low volume, hand held, mechanical or motorized ground broadcast equipment. Application to small areas may be made with low-pressure hand wand or backpack equipment. Maintain constant agitation during application.

Depending on the disease, water this product into the crown and active root zone for best results. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. For best results, use spray mixture the same day it is prepared.

Golf Course Turf Disease Control

DISEASE	RATE of Tide USA TEBU 3.6 F Fungicide in Fl. Oz./1000 Sq. Ft. (Lbs. ai/1000 Sq. Ft.)		Application Directions	Use Restrictions
	California	Other States		
Dollar Spot (<i>Sclerotinia homoeocarpa</i>)	0.6 (0.017)	0.6 to 1.1 (0.017 to 0.031)	For prevention, begin applications when conditions are favorable for disease development. Alternate with another fungicide with a different mode of action. A second application may be made after 21 days. California – maintain a retreatment interval of 28 days.	DO NOT make two consecutive applications of this product. In California, DO NOT apply more than 0.6 fl. oz. (0.017 lb. ai)/1000 sq. ft. per application. In other states, DO NOT apply more than 1.1 fl. oz. (0.031 lb. ai)/1000 sq. ft. per application. In California, minimum retreatment interval = 28 days. In other states, minimum retreatment interval = 21 days.
Copper Spot (<i>Gloeocercospora sorghi</i>)				
Powdery Mildew (<i>Erysiphe graminis</i>)				
Corticium Red Thread (<i>Laetisaria fuciformis</i>)				
Rusts (<i>Puccinia</i> spp.)				
Brown Patch/Rhizoctonia Blight, Large Patch (<i>Rhizoctonia solani</i>)	0.6 (0.017)	0.6 to 1.1 (0.017 to 0.031)	For prevention, begin applications when conditions are favorable for disease development. DO NOT make two consecutive applications of this product. Alternate with another fungicide with a different mode of action. A second application may be made after 21 days. California – maintain a retreatment interval of 28 days.	DO NOT make two consecutive applications of this product. In California, DO NOT apply more than 0.6 fl. oz. (0.017 lb. ai)/1000 sq. ft. per application. In other states, DO NOT apply more than 1.1 fl. oz. (0.031 lb. ai)/1000 sq. ft. per application. In California, minimum retreatment interval = 28 days. In other states, minimum retreatment interval = 21 days.
Brown Ring Patch (<i>R. circinata</i>)				
Anthracnose-Basal and Foliar (<i>Colletotrichum cereal</i>)	0.6 (0.017)	0.6 to 1.1 (0.017 to 0.031)		
Red Thread (<i>Laetisaria fuciformis</i>)				
Pink Patch (<i>Limonomyces rosipellis</i>)				
Bermuda Grass decline (<i>Gaeumannomyces graminis</i> var. <i>graminis</i>)	0.6 (0.017)	0.6 to 1.1 (0.017 to 0.031)	Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent on the depth of the root zone. For prevention, begin applications two or four weeks prior to the historical appearance of disease symptoms. Initiate cultural control practices at the same time the fungicide is applied. Refer to your local County Extension Service for this information. Apply subsequent applications at 21 day intervals. California – maintain a retreatment interval of 28 days.	In California, DO NOT apply more than 0.6 fl. oz. (0.017 lb. ai)/1000 sq. ft. per application. In other states, DO NOT apply more than 1.1 fl. oz. (0.031 lb. ai)/1000 sq. ft. per application. In California, minimum retreatment interval = 28 days. In other states, minimum retreatment interval = 21 days.

DISEASE	RATE of Tide USA TEBU 3.6 F Fungicide in Fl. Oz./1000 Sq. Ft. (Lbs. ai/1000 Sq. Ft.)		Application Directions	Use Restrictions
	California	Other States		
Take All Patch <i>(Gaeumannomyces graminis)</i>	0.6 (0.017)	0.6 to 1.1 (0.017 to 0.031)	For prevention, apply in the fall when soil temperature reaches 55-65°F and again in the spring under similar soil temperature conditions. Applications in both fall and spring may be necessary. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone. Subsequent applications at 21 days intervals may be necessary for both fall and spring applications. California – maintain a retreatment interval of 28 days.	In California, DO NOT apply more than 0.6 fl. oz. (0.017 lb. ai)/1000 sq. ft. per application. In other states, DO NOT apply more than 1.1 fl. oz. (0.031 lb. ai)/1000 sq. ft. per application. In California, minimum retreatment interval = 28 days. In other states, minimum retreatment interval = 21 days.
Gray Leaf Spot <i>(Pyricularia grisea)</i>	0.6 (0.017)	0.6 to 1.1 (0.017 to 0.031)	Apply when conditions are favorable for disease development at 21 day intervals. This product can be tank mixed with a registered contact fungicide at the label rate. California – maintain a retreatment interval of 28 days.	In California, DO NOT apply more than 0.6 fl. oz. (0.017 lb. ai)/1000 sq. ft. per application. In other states, DO NOT apply more than 1.1 fl. oz. (0.031 lb. ai)/1000 sq. ft. per application. In California, minimum retreatment interval = 28 days. In other states, minimum retreatment interval = 21 days.
Stripe Smut <i>(Ustilago striiformis)</i>	0.6 (0.017)	0.6 to 1.1 (0.017 to 0.031)	Make a single application to historical disease areas in spring as grass growth begins.	In California, DO NOT apply more than 0.6 fl. oz. (0.017 lb. ai)/1000 sq. ft. per application. In other states, DO NOT apply more than 1.1 fl. oz. (0.031 lb. ai)/1000 sq. ft. per application.
Spring Dead Spot <i>(Leptosphaeria korrae, L. narmari, Ophiosphaerella herpotricha, Gaeumannomyces graminis)</i> Necrotic Ring Spot <i>(Leptosphaeria korrae)</i>	0.6 (0.017)	0.6 to 1.1 (0.017 to 0.031)	For prevention, apply in fall when soil temperature reaches 55-60 °F and again in spring under similar soil temp conditions or after dormancy break. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.	In California, DO NOT apply more than 0.6 fl. oz. (0.017 lb. ai)/1000 sq. ft. per application. In other states, DO NOT apply more than 1.1 fl. oz. (0.031 lb. ai)/1000 sq. ft. per application.

DISEASE	RATE of Tide USA TEBU 3.6 F Fungicide in Fl. Oz./1000 Sq. Ft. (Lbs. ai/1000 Sq. Ft.)		Application Directions	Use Restrictions
	California	Other States		
Fusarium Patch <i>(Fusarium roseum)</i>	0.6 (0.017)	0.6 to 1.1 (0.017 to 0.031)	Apply first application in mid-June or 14-28 days prior to time this blight normally becomes evident. Make applications at no less than 21 day intervals. California – maintain a retreatment interval of 28 days.	In California, DO NOT apply more than 0.6 fl. oz. (0.017 lb. ai)/1000 sq. ft. per application. In other states, DO NOT apply more than 1.1 fl. oz. (0.031 lb. ai)/1000 sq. ft. per application. In California, minimum retreatment interval = 28 days. In other states, minimum retreatment interval = 21 days.
Summer Patch <i>(Magnaporthe poae)</i>	0.6 (0.017)	0.6 to 1.1 (0.017 to 0.031)	Apply beginning in the spring. Alternate with another fungicide with a different mode of action. Second and third applications may be made at 28 day intervals. See local university guidance for suggested timing. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.	DO NOT make two consecutive applications of this product. In California, DO NOT apply more than 0.6 fl. oz. (0.017 lb. ai)/1000 sq. ft. per application. In other states, DO NOT apply more than 1.1 fl. oz. (0.031 lb. ai)/1000 sq. ft. per application. In California, minimum retreatment interval = 28 days. In other states, minimum retreatment interval = 21 days.
Zoysia Patch, Large Patch of zoysia <i>(Rhizoctonia solani)</i>	0.6 (0.017)	0.6 to 1.1 (0.017 to 0.031)	Make first application in early fall (mid-September to mid-October) prior to development of disease symptoms. A second application in early spring may be necessary in areas where disease pressure is known to be heavy.	In California, DO NOT apply more than 0.6 fl. oz. (0.017 lb. ai)/1000 sq. ft. per application. In other states, DO NOT apply more than 1.1 fl. oz. (0.031 lb. ai)/1000 sq. ft. per application.
Gray Snow Mold/Typhula Blight <i>(Typhula incarnate)</i> Pink Snow Mold/ Microdochium Patch <i>(Microdochium nivale)</i>	0.6 (0.017)	0.6 to 1.1 (0.017 to 0.031)	Apply in the fall, before anticipated turf dormancy and before the first snow cover. If turf breaks dormancy during winter months, a second application may be made. It is advised that this product be tank-mixed with other registered snow mold products for best season long results.	In California, DO NOT apply more than 0.6 fl. oz. (0.017 lb. ai)/1000 sq. ft. per application. In other states, DO NOT apply more than 1.1 fl. oz. (0.031 lb. ai)/1000 sq. ft. per application. DO NOT apply over snow cover, or when turf is dormant.

DISEASE	RATE of Tide USA TEBU 3.6 F Fungicide in Fl. Oz./1000 Sq. Ft. (Lbs. ai/1000 Sq. Ft.)		Application Directions	Use Restrictions
	California	Other States		
Fairy Ring <i>(Chlorophyllum (Lepiota), Lycoperdon, Marasmius spp.)</i>	N/A	0.6 to 1.1 (0.017 to 0.031)	For Cool Season turf make preventative applications in the spring when soil temperatures reach 55-60°F. Make applications at no less than 21 days intervals. Immediately after fungicide is applied, the area must be irrigated with sufficient water to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent on the depth of the root zone. Alternate with another fungicide containing a different mode of action. Use a wetting agent under hydrophobic soil conditions.	DO NOT apply more than 1.1 fl. oz. (0.031 lb. ai)/1000 sq. ft. per application. Minimum retreatment interval = 21 days. DO NOT use a wetting agent under hydrophilic soil conditions. Dormancy breaking Warm Season turf, DO NOT make two consecutive applications of this product or another fungicide containing the active ingredient Tebuconazole. Not for use in California for control of Fairy Ring.
<p>NOTE: Apply the specified amount of this product in 1.5 to 3.0 gallons of water per 1000 sq. ft. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. DO NOT use clippings for animal feed. DO NOT exceed 3.6 fl. oz of this product (0.10 lb. ai) per 1000 sq. ft. per year. DO NOT exceed 6 applications per year except in New York. In New York State, DO NOT apply more than 3 applications of 0.6 fl. oz. (0.017 lb. ai) not to exceed 1.8 fl. oz. of this product (0.05 lb. ai) per 1,000 sq. ft. per year (equivalent to 2.2 lbs. ai/acre/year).</p>				

DISEASE CONTROL IN FIELD, NURSERY AND CONTAINER ORNAMENTALS AND COMMERCIAL and RESIDENTIAL LANDSCAPES

ORNAMENTAL USE RESTRICTIONS

- For use on ornamental plants only, not for woodlands or forest management.
- Intended only for use by professional applicators.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply more than 10 fl. oz. (0.28 lb. ai) per acre in a single application.
- **DO NOT** apply more than 0.31 gallons (40 fl. oz.) of this product (equal to 1.13 lbs. of tebuconazole) per acre per year; except to Leatherleaf fern (Florida only) where the maximum annual application rate is 5 pints (80 fl. oz.) of this product (equal to 2.25 lbs. of tebuconazole) per acre per year.
- **DO NOT** make more than 4 applications per year at highest rate, except to Leatherleaf fern (Florida only) where the maximum number of applications per year is 8 at the highest rate.
- Minimum retreatment interval: 14 days; except for applications to prevent Petal Blight where applications may be made 2-3 times per week into the flowers as they open and develop color and except to Leatherleaf fern (Florida only) where the minimum retreatment interval is 12 days.
- **DO NOT** apply to bearing fruit trees or vegetables.

Use this product in a preventative and curative disease control program for the listed plant types and disease in the table below. Optimum disease management is obtained when this product is used in conjunction with sound disease management practices.

Apply material with properly calibrated hand held, mechanical or motorized spray equipment. Begin applications when disease first appears and repeat at 14-21 day intervals during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease. Under heavy disease pressure, use the highest rate and the shortest interval.

For hand held, mechanical, or motorized applications, mix as directed below and apply as a full coverage spray to drip for the prevention and control of the diseases listed below. Choose a finished spray volume appropriate for the size of the plants and amount of foliage, which will provide thorough coverage throughout the canopy. Allow sprays to dry before overhead irrigation is applied.

Apply this product at rates of 4 – 10 fl oz per acre in 100 gallons of water. Spray volume may range from 50 up to 300 gallons of finished spray per acre depending upon equipment, plant species and plant growth stage at time of application.

Note: The "Directions for Use" of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product is not advised for use on African Violets, Begonias, Boston Fern, and Geraniums.

ORNAMENTALS DISEASE CONTROL

PLANTS	DISEASE	APPLICATION	
		To Prevent Disease	To Treat Existing Disease
Roses	Black Spot Powdery Mildew Rust	Apply every 14-21 days during the growing season, starting when leaves first appear.	Apply every 14 days for a total of 3 applications beginning at the first sign of disease.
Flowers	Leaf Spot Powdery Mildew Rust Southern Blight	Apply at least 3 times per year, 14-21 days apart, beginning with Spring bud break. Rotation or Tank mixing with barrier protectant fungicides is advised for resistance management.	
Crabapples (Ornamental), Dogwoods and Other Landscape (Ornamental) Trees	Anthraxnose Leaf Spot Powdery Mildew Rust Scab		
Azaleas, Camellias, Rhododendrons and Other Landscape (Ornamental) Shrubs	Anthraxnose Black Spot Leaf Spot Petals Blight	Petals Blight – Apply 2-3 times per week into the flowers as they open and develop color.	
Ground Covers and Vines	Powdery Mildew Rust Southern Blight		

HOW MUCH TO USE FOR SMALL PLANTINGS: ADD 1 TEASPOON TO 2.5 GALLONS OF WATER.

Restricted-entry interval (REI) = 12 hours

PLANTS	DISEASE	RATE IN FL. OZ. OF TIDE USA TEBU 3.6 F FUNGICIDE /ACRE	REMARKS
Leatherleaf Fern (Florida only)	Anthracnose (suppression)	5 – 10	<p>Make the first application before anthracnose symptoms are present and continue at 12- to 14-day intervals.</p> <p>Apply Tide USA TEBU 3.6 F Fungicide in a minimum of 5 gallons of spray solution per acre using ground equipment or chemigation.</p> <p>DO NOT apply more than 5 pints (80 fl oz; 2.25 lb AII/A per year) of Tide USA TEBU 3.6 F Fungicide per acre per year.</p>
<p>USE LIMITATIONS: Tide USA TEBU 3.6 F Fungicide can cause phytotoxicity to Leatherleaf fern under certain environmental conditions. Applications in temperatures less than 70°F can cause phytotoxicity in the form of leaf burning and/or yellowing. Application followed by temperatures falling below 55°F can cause similar symptoms. Before using this product on Leatherleaf Fern, read the CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND DISCLAIMER section in its entirety</p>			

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, secure, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If container is leaking, invert to prevent leakage. If the container is leaking or material is spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. **DO NOT** walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticide below. In spill or leak incidents, keep unauthorized people away.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

NONREFILLABLE CONTAINERS

Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying.

(Nonrefillable container ≤ 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND DISCLAIMER

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Tide International USA, Inc. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TIDE INTERNATIONAL USA, INC. MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. NO AGENT OF TIDE INTERNATIONAL USA, INC. IS AUTHORIZED TO MAKE ANY WARRANTIES BEYOND THOSE CONTAINED HEREIN OR TO MODIFY THE WARRANTIES CONTAINED HEREIN. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, TIDE INTERNATIONAL USA, INC. DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT TIDE INTERNATIONAL USA, INC.'S ELECTION, THE REPLACEMENT OF PRODUCT.

Label Code No.: TIC-FUN8422934

EPA 20231212

Created: 06/17/2024