
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 953A-S11 0425

| | | | |
|--------------------|--------------|-----------|------------------|
| CYPRODINIL | GROUP | 9 | FUNGICIDE |
| FLUDIOXONIL | GROUP | 12 | FUNGICIDE |

Switch® 62.5WG

Fungicide

This supplemental label expires on 4/17/2028 and must not be used or distributed after this date.

Active Ingredients:

| | |
|---------------------|--------|
| Cyprodinil* | 37.5% |
| Fludioxonil** | 25.0% |
| <hr/> | |
| Other Ingredients: | 37.5% |
| Total: | 100.0% |

*CAS No. 121552-61-2

**CAS No. 131341-86-1

Switch 62.5WG is a water-dispersible granule containing 37.5% cyprodinil and 25% fludioxonil.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

EPA Reg. No. 100-953

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Switch 62.5WG as permitted according to this Supplemental Labeling, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of federal law to use this product in a manner inconsistent with its labeling.

syngenta

DIRECTIONS FOR USE

| Crop | Disease | Product Rate oz/Acre | Use Directions |
|--|---|-------------------------|--|
| Cranberry (Not registered for use in California) | Yellow rot (<i>Botrytis cinerea</i>) | 11-14* | Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development Resistance Management: After 2 applications of Switch 62.5WG, alternate with another fungicide with a different mode of action for 2 applications. |
| | Powdery mildew (<i>Erysiphe vaccinii</i>) | | |
| | Bitter Rot (<i>Colletotrichum</i> spp.) | | |
| | Alternaria Fruit Rot (<i>Alternaria</i> spp.) | | |
| | Viscid Rot (<i>Phomopsis</i> spp.) | | |
| *11 oz product/A contains 0.26 lb cyprodinil/A and 0.17 lb fludioxonil/A. *14 oz product/A contains 0.33 lb cyprodinil/A and 0.22 lb fludioxonil/A. | | | |
| Application Instructions: Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 5 gallons/A spray volume by air. Make no more than two applications by air. The use of an adjuvant at the specified rates may improve disease control. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy. | | | |
| Specific Use Restrictions <ol style="list-style-type: none"> 1) Maximum Single Application Rate: DO NOT exceed the maximum rate listed in the table above. 2) DO NOT apply more than 4 applications per year at the highest rate mentioned in the table above. 3) Minimum Application Interval: 7 days 4) Applications can be made to a dry bed only. No direct applications to water. 5) DO NOT allow release of irrigation or flood water for at least 7 days after the last application. 6) The use of handheld misters or foggers is prohibited. 7) DO NOT make more than two applications by air. 8) DO NOT apply more than 56 oz/A of Switch 62.5WG per year (1.3 lb cyprodinil and 0.9 lb fludioxonil). 9) DO NOT apply more than 1.3 lb ai/A of cyprodinil-containing products and 0.9 lb ai/A of fludioxonil-containing products per year. 10) DO NOT apply within 30 days of harvest (30-day PHI). | | | |

Switch® 62.5WG and the Syngenta logo are trademarks of a Syngenta Group Company