

Bayer CropScience LP P.O. Box 12014 2 T.W. Alexander Drive Research Triangle Park, North Carolina 27709 1-866-99BAYER (1-866-992-2937)

SIVANTO™ prime

For Suppression of Chilli Thrips (Scirtothrips dorsalis) in Strawberry, Lowbush Blueberry and Highbush Blueberry.

EPA Reg. No. 264-1141

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF FLORIDA

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

BLUEBERRY (LOWBUSH); STRAWBERRY - FOLIAR	
Pest Suppressed	Product Rate (fl oz/A)
Chilli Thrips (Scirtothrips dorsalis)	14.0
Foliar Application Restrictions:	
Pre-Harvest Interval (PHI): 0 day	

Minimum interval between applications: 10 days

Minimum application volumes: 10 gallons/Acre (Ground); 3 gallons/Acre (Aerial)

Maximum number of applications per year: 2

Maximum SIVANTO prime allowed per year: 28.0 fluid ounces/Acre (0.365 lb Al/Acre).

Product Rate (fl oz/A)
14.0
_

Foliar Application Restrictions: Pre-Harvest Interval (PHI): 3 days

Minimum interval between applications: 7 days

Minimum application volumes: 25 gallons/Acre (Ground); 3 gallons/Acre (Aerial)

Maximum number of applications per year: 2

Maximum SIVANTO prime allowed per year: 28.0 fluid ounces/Acre (0.365 lb Al/Acre).

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577. For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

> As with any crop-protection product, always read and follow label directions. For additional information call toll-free 1-866-99BAYER (1-866-992-2937).