



Arysta LifeScience

SUPPLEMENTAL LABELING

GROUP 11 FUNGICIDE

EVITO[®] 480 SC

Supplemental Label to Add **DIRECTIONS FOR USE** on **BARLEY**.

This supplemental label expires April 30, 2019 and must not be used or distributed after this date.

For Agricultural Use Only

INGREDIENTS: _____ % BY WT.

ACTIVE INGREDIENT:

Fluoxastrobin: [(1*E*)-[2-[[6-(2-Chlorophenoxy)-5-fluoro-4-pyrimidinyl]oxy]phenyl] (5,6-dihydro-1,4,2-dioxazin-3-yl) methanone-O-methyloxime].....40.3%

OTHER INGREDIENTS:59.7%

TOTAL:100.0%

This product contains 3.98 pounds of fluoxastrobin per gallon (478 g per liter).

KEEP OUT OF REACH OF CHILDREN CAUTION / PRECAUCIÓN

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR **EVITO[®] 480 SC** BEFORE APPLYING. USE OF **EVITO 480 SC** ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS, RESTRICTIONS, AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR **EVITO 480 SC (EPA Reg. No. 66330-64)**.

102517V001Su

BARLEY

Disease Control	Product Rate to Use	Application Directions
Barley Stripe <i>(Drechslera graminea = Pyrenophora graminea)</i>	2.0-4.0 fl oz/A (0.063-0.12 lb ai/A)	For optimum results, begin applications preventively and continue as needed on a 14- to 21-day interval. Use the higher specified rates and shorter interval when disease pressure is high.
Net Blotch <i>(Pyrenophora feres)</i>		For early-season disease suppression you can apply product at 1.0-2.0 fl oz/A.

Leaf Rust <i>(Puccinia recondita f. sp. tritici)</i> Stripe Rust <i>(Puccinia striiformis)</i> Stem Rust <i>(Puccinia graminis)</i> Scald <i>(Rhynchosporium secalis)</i> Septoria Leaf and Glume Blotch <i>(Septoria tritici, Septoria nodorum)</i> Spot Blotch <i>(Cochliobolus sativus)</i> Tan Spot <i>(Pyrenophora tritici-repentis)</i>		An adjuvant may be added to the tank.
Powdery Mildew <i>(Erysiphe graminis)</i> Stagonospora Blotch <i>(Stagonospora nodorum)</i>	2.5-4.0 fl oz/A (0.079-0.12 lb ai/A)	
RESTRICTIONS: <ul style="list-style-type: none"> • Do not apply more than 4.0 fl oz (0.12 lb ai/A) of product per acre per single application. • Do not apply more than 0.24 lb ai of fluoxastrobin per acre per year in barley. • There is a maximum number of 2 applications per year for barley, and a minimum interval of 14 days between applications. • Do not apply later than Feekes growth stage 10.5. • Do not apply product within 40 days of harvest for barley. • Do not apply product within 7 days of harvest for forage and hay. 		

Produced For:
ARYSTA LIFESCIENCE NORTH AMERICA, LLC
15401 Weston Parkway, Suite 150
Cary, NC 27513

Arysta LifeScience and the Arysta LifeScience logo are registered trademarks of Arysta LifeScience Corporation.

EVITO and the EVITO logo are registered trademarks of an Arysta LifeScience Group Company.

©2017 Arysta LifeScience Group Company.