
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC

P. O. Box 18300

Greensboro, North Carolina 27419-8300

SCP 1476A-S2A 1220

BENZOVINDIFLUPYR	GROUP	7	FUNGICIDE
DIFENOCONAZOLE	GROUP	3	FUNGICIDE

Aprovia® Top Fungicide

SOLATENOL® Technology*

This supplemental label expires on 12/30/2023 and must not be used or distributed after this date.

Active Ingredients:

Difenoconazole** 10.95%

Benzovindiflupyr*** 7.30%

Other Ingredients: 81.75%

Total: 100.00%

*Technology denotes the active ingredient, Benzovindiflupyr.

**CAS No. 119446-68-3

***CAS No. 1072957-71-1

Aprovia Top Fungicide is formulated as an emulsifiable concentrate containing 0.97 lb ai of difenoconazole active ingredient and 0.65 lb ai of benzovindiflupyr active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN.**WARNING/AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1476

All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Aprovia Top Fungicide as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR USE

Blueberries

Crops (Including all cultivars, varieties, and/or hybrids of these)			
Blueberry (lowbush only)			
Target Disease	Rate (fl oz/A)	Application Timing	Use Directions
Blueberry leaf rust (<i>Thekopsora minima</i>) Septoria leaf spot (<i>Septoria</i> spp.)	13.5	Apply at first sign of diseases.	Apply by ground or by air. A second application can be made after 10-14 days. Apply in a minimum spray volume of 20 gallons per acre. See Section 4.4.5.
Resistance Management:			
<ul style="list-style-type: none"> • Refer to Section 3.1. 			
USE RESTRICTIONS			
<ol style="list-style-type: none"> 1. Refer to Section 6.1 for additional product use restrictions. 2. Maximum Single Application Rate: 13.5 fl oz/A 3. Minimum Application Interval: 10 days 4. Maximum Annual Rate: 27 fl oz/A/year <ol style="list-style-type: none"> a. DO NOT exceed 0.136 lb ai/A/year of benzovindiflupyr-containing products. b. DO NOT exceed 0.46 lb ai/A/year of difenoconazole-containing products 5. DO NOT exceed 2 applications per year. 6. DO NOT apply by air in New York State. 7. Pre-Harvest Interval (PHI): 7 day 			

Ginseng

Crops (Including all cultivars, varieties, and/or hybrids of these) Not for use in California			
Ginseng			
Target Disease	Rate (fl oz/A)	Application Timing	Use Directions
Ginseng Alternaria blight (<i>A. panax</i>) Powdery mildew (<i>Erysiphe spp.</i>)	13.5	For foliar disease, make an application at the onset of disease or when conditions are conducive for disease.	Apply by ground. For ground applications, use a minimum of 50 gal/A of water. See Section 4.4.5.
Resistance Management:			
<ul style="list-style-type: none"> Refer to Section 3.1. For resistance management, make no more than 2 applications before alternating to another fungicide with a non-Group 7 mode of action. 			
USE RESTRICTIONS			
<ol style="list-style-type: none"> Refer to Section 6.1 for additional product use restrictions. Maximum Single Application Rate: 13.5 fl oz/A Minimum Application Interval: 14 days Maximum Annual Rate: 54 fl oz/A/year <ol style="list-style-type: none"> DO NOT exceed 0.272 lb ai/A/year of benzovindiflupyr-containing products. DO NOT exceed 0.46 lb ai/A/year of difenoconazole-containing products. DO NOT exceed 4 applications per year. Pre-Harvest Interval (PHI): 15 days 			

©2021 Syngenta

Aprovia® Top, SOLATENOL® and the Syngenta logo are trademarks of a Syngenta Group Company