



TECH-GRO™



B-17 BORIC ACID SPRAY™



GUARANTEED ANALYSIS

Boron (B).....	17.0%
Derived from: Boric Acid.	

TECH-GRO B-17 is a soluble, fine granular product formulated with improved dispersion and wetting characteristics to provide superior foliar boron performance.

**ADEQUATE AGITATION SHOULD BE MAINTAINED AT ALL TIMES
DURING MIXING AND APPLICATION.**

WARNING: This product contains Boron (B) and is to be used only where Boron is known to be beneficial and on the crops recommended on this label. Any other use may result in injury to crops.

MADE IN USA
NUTRIENT TECHNOLOGIES, INC.
1092 E. KAMM AVE., DINUBA CA 93618
(559)595-8090

TECH-GRO™ B-17 Boric Acid Spray™



GENERAL RECOMMENDATIONS

B-17 BORIC ACID SPRAY at a rate of 5.9 lbs. provides the equivalent of 1 lb. of boron. B-17 contains no sodium and is mildly acidic.

DILUTION: Apply the suggested rates of B-17 BORIC ACID SPRAY in the following amounts of water per acre. Minimum dilution rate - 1 lb. per 5 gallons of water per suggested rate.

Field & Row Crops: By air 3-20 gals.
By ground 10 gals. minimum.
Tree Crops: By air 10 gals. minimum.
By ground concentrate 25-100 gals.
By ground dilute 100-500 gals.
In oil sprays, use 1-5 lbs. per 100 gallons.

COMPATIBILITY: This product has been designed to be compatible with most commonly used insecticides, fungicides and nutritional products. However, a compatibility test is recommended for combinations that have not been used previously. Consult your NUTRI-TECH representative for detailed compatibility information.

FOR ALL CROPS- DO NOT EXCEED THE RATE OF 6.5 LBS. PER ACRE PER YEAR OF B-17 IN FOLIAR SPRAYS, 12 LBS. PER ACRE IN SOIL APPLICATIONS. SUGGESTED RATES MAY BE APPLIED IN SPLIT APPLICATIONS.

Information regarding the contents and levels of metals in this product is available on internet at: <http://aapfco.org/metals.htm>



WARNING

Suspected of damaging fertility
or the unborn child

Wear appropriate personal protective equipment.

Avoid breathing dust.

Do not ingest this product.

Dispose of unused contents, container, and other contaminated wastes in accordance with local, state, and federal regulations.

If exposed or concerned: Get medical advice or attention.

SUGGESTED USES (RATES PER ACRE)

Alfalfa, Hay, & Other Forage Crops - Apply 0.5-3 lbs. over spring growth, repeat after each cutting.
Almonds, Peaches, Plums, Cherries, & Other Stone Fruits - Apply 1-5 lbs. pre-bloom and at 3-4 week intervals as needed post-bloom. (SEE BELOW)
Apples, Pears - Apply 1-6.25 lbs. at 2-4 week intervals starting pre-bloom. (SEE BELOW)
Barley, Oats, Rice, Wheat & Other Small Grains - Apply 0.5-3 lbs. at tillering, repeat at early boot if necessary.
Beans, Peas, & related Legumes - Apply 0.5-3 lbs. at early bud, repeat after bloom if necessary.
Cabbage, Cauliflower, Broccoli, Brussels Sprouts & Other Cole Crops - Apply 0.5-3 lbs. as needed after thinning or transplant.
Carrots, Onions, Radishes, Beets & Other Vegetable Root Crops - Apply 0.5-3 lbs. when top growth is 4-8" high, repeat as necessary.
Citrus, Avocados - Apply 1-5 lbs. in spring, summer, and fall sprays.
Corn, Field & Sweet - Apply 0.5-3 lbs. when 10-20" high, repeat in 2-4 weeks as necessary.
Cotton - Apply 0.5-3 lbs. at first square, repeat in 2-4 weeks as necessary.
Cucumbers, Melons, Squash & Other Cucurbits - Apply 0.5-3 lbs. pre-bloom, repeat in 2-4 weeks as necessary.
Grapes - Apply 0.5-3 lbs. dormant or as needed pre-bloom to post-bloom.
Hops - Apply 0.5-3 lbs. as needed in cover sprays.
Lettuce, Celery, Spinach & Other Green Leafy Vegetables - Apply 0.5-3 lbs. as needed after thinning and transplant.
Mint - Apply 0.5-3 lbs. over new growth, repeat as needed.
Peanuts - Apply 0.5-3 lbs. at early bloom and just prior to pegging.
Pecans, & Other Deciduous Nut Trees - Apply 1-5 lbs. at 3-4 week intervals starting at bud break.
Olives - Apply 0.5-5 lbs. pre-bloom or post-bloom, repeat 3-4 weeks as necessary.
Ornamentals, Trees & Shrubs - Apply 0.5-3 lbs. at 4-6 week intervals as needed.
Potatoes - Apply 0.5-3 lbs. at tuber initiation, repeat in 2-4 weeks as necessary.
Raspberries, Blackberries, Blueberries, Cranberries & Other Small Fruit - Apply 0.5-3 lbs. pre-bloom, repeat in 2-4 weeks as necessary.
Soybeans, Seed Crops - Apply 0.5-3 lbs. pre-bloom, repeat in 2-4 weeks as necessary.
Sugarbeets - Apply 0.5-3 lbs. at 4th-6th leaf, repeat in 2-4 weeks as necessary.
Strawberries - Apply 0.5-3 lbs. pre-bloom, repeat in 2-4 weeks as necessary.
Tomatoes, Peppers, Eggplant & Okra - Apply 0.5-3 lbs. pre-bloom, repeat in 2-4 weeks as necessary.
Walnuts - Apply 1-5 lbs. in blight sprays.

For apples, pears, and all deciduous stone fruits, including almonds, B-17 may be applied in delayed dormant programs and post harvest sprays as per local recommendations.

SOIL APPLICATIONS:

For all crops: 3-12 lbs per acre as needed.

NOTICE: Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used according to directions under normal conditions on the crops specified. Because of widely varying use conditions, it is impossible to eliminate all risks even when the label directions are followed. Therefore, the manufacturer makes no other warranties, expressed or implied. The buyer assumes all risks associated with the use of this product. In the event of damage resulting from a breach of warranty, the buyer agrees to accept a refund of the purchase price as full discharge of the manufacturer's liability.

MADE IN USA
NUTRIENT TECHNOLOGIES, INC.
1092 E. KAMM AVE., DINUBA CA 93618
(559)595-8090