

Supplemental Labeling



Corteva Agriscience LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Delegate[®] WG

EPA Reg. No. 62719-541

Tropical Tree Fruits¹

¹Acerola, atemoya, avocado, biriba, black sapote, canistel, cherimoya, custard apple, feijoa, guava, ilama, jaboticaba, longan, lychee, mamey sapote, mango, papaya, passionfruit, pitaya (dragon fruit), pulasan, rambutan, sapodilla, soursop, Spanish lime, star apple, starfruit, sugar apple, ti leaves, wax jambu, white sapote

This label expires and must not be distributed or used in accordance with this supplemental registration after March 26, 2024.

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Delegate[®] WG insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Delegate WG according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Delegate WG.

Directions for Use

Pests and Application Rates:

| Pests | Delegate WG (oz/acre) |
|---|-----------------------|
| katydids ¹ lepidopterous larvae, including: avocado leafroller citrus peelminer cutworms fruit tree leafroller light brown apple moth navel orangeworm orange tortrix western tussock moth thrips ² | 4 – 7 |

¹Katydid: Control of nymphs only, suppression of adults.

²Control of thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants section under Mixing Directions.

Application Timing: Treat when pests appear, or in accordance with local economic thresholds. Consult your Corteva Agriscience representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate: The amount of Delegate WG applied per acre will depend upon tree size and pest pressure. Use a lower rate in the rate range for light infestations and/or small trees and a higher rate in the rate range for heavy infestations and/or large trees.

Resistance Management: Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after two consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your Corteva Agriscience representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area. Do not apply to tropical tree fruits grown in greenhouses and nurseries.

Restrictions except for pitaya (dragon fruit):

- **Preharvest Interval:** Do not apply within 1 day of harvest.
- Do not apply more than a total of 14 oz of Delegate WG (0.219 lb ai spinetoram) per acre per year.
- **Maximum Number of Applications:** Do not make more than three applications per calendar year.
- **Minimum Treatment Interval:** Do not apply treatments less than 4 days apart.

Restrictions for pitaya (dragon fruit):

- **Preharvest Interval:** Do not apply within 1 day of harvest.
- Do not apply more than a total of 24 oz of Delegate WG (0.3752 lb ai spinetoram) per acre per year.
- **Maximum Number of Applications:** Do not make more than four applications per calendar year.
- **Minimum Treatment Interval:** Do not apply treatments less than 4 days apart.

™®Trademarks of Corteva Agriscience and its affiliated companies

R359-041

EPA accepted 04/15/2021

Initial printing