

RESTRICTED USE PESTICIDE**Due to Toxicity to Fish and Aquatic Organisms****For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.****PRODUCT INFORMATION BULLETIN****FIFRA Section 2(ee) Recommendation****Expiration Date: April 2022**

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF CALIFORNIA, CONNECTICUT, DELAWARE, IDAHO, INDIANA, MASSACHUSETTS, MARYLAND, MAINE, MICHIGAN, NEVADA, NEW JERSEY, OHIO, OREGON, PENNSYLVANIA, RHODE ISLAND, UTAH, VERMONT, VIRGINIA, WASHINGTON AND WISCONSIN

GROUP**3****INSECTICIDE****SULTRUS™**

EPA Reg. No. 5905-599

For Use on Pome Fruit to control Spotted Wing Drosophila

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. This product bulletin should be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling.

Pome Fruit

Includes all members of Crop Group 11: Apple, Crabapple, Loquat, Mayhaw, Pear, Oriental pear, and Quince

PEST CONTROLLED	Rate fluid ounces/Acre	Rate lb. AI/Acre	Method
Spotted Wing Drosophila (adult)	2.4 – 2.8	0.019 – 0.022	Apply as a foliar spray, targeting adult growth stage. Reapply as necessary but no closer than a 14-day interval.
Notes and Restrictions: Pre-Harvest Interval: 7 days Maximum SULTRUS allowed per 14-day interval: 2.8 fluid ounces/Acre (0.022 lb AI/Acre) Maximum SULTRUS allowed per crop season: 2.8 fluid ounces/Acre (0.022 lb AI/Acre)			

**HELENA AGRI-ENTERPRISES, LLC
225 SCHILLING BOULEVARD, SUITE 300
COLLIERVILLE, TN 38017**