

# FIFRA Section 24(c) Special Local Need (SLN) Label

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Stinger®

EPA Reg. No. 62719-73

24(c) Special Local Need Registration EPA SLN No. IN-220002

### Postemergence Broadleaf Weed Control in Strawberry (For Distribution and Use Only in the State of Indiana)

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2027

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Stinger® herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- In addition to restrictions and limitations imposed by this Special Local Need labeling, use of Stinger is subject to all use precautions and limitations imposed by the label affixed to the container for Stinger.

#### Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters. Do not contaminate water used for irrigation or domestic purposes.

#### Ground Water Advisory

Clopyralid is a chemical which can travel (seep or leach) through soil and under certain conditions contaminate groundwater which may be used for irrigation or drinking purposes. Users are advised not to apply clopyralid where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an underlying aquifer is shallow, or to soils containing sinkholes over limestone bedrock, severely fractured surfaces, and substrates which would allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

#### Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a medium potential for reaching surface water via runoff for several weeks after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of clopyralid from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

#### Directions for Use

Refer to product label for General Use Precautions, Restrictions, Mixing Directions, and Application Directions.

## Target Broadleaf Weeds, Application Rate, and Use Restrictions

Target Broadleaf Weeds	Stinger (pint/acre)	Use Restrictions
artichoke, Jerusalem clovers cocklebur, common curly dock dandelion groundsel jimsonweed lettuce, prickly nightshade ragweed, common ragweed, giant smartweed (suppression) sorrel, red sowthistle, annual (suppression) sunflower thistle, bull (suppression) thistle, Canada (suppression) vetch	<p><b>Post-harvest application:</b> 1/3 to 2/3</p> <p><b>Spring application:</b> 1/3</p>	<ul style="list-style-type: none"> <li>• Make 1 to 2 applications per year not to exceed 2/3 pint per acre per year. Make only 1 application in the spring.</li> <li>• <b>Do not</b> use this product with a surfactant on strawberries.</li> <li>• <b>Do not</b> tank mix this product with other herbicides registered for use on strawberry.</li> <li>• <b>Spring applications:</b> Do not apply this product within 30 days of harvest.</li> </ul>

**Note:** Some minor leaf cupping may occur. Do not use if unwilling to accept minor leaf injury.

### Broadcast Applications

Apply uniformly with ground equipment in 20 to 75 gallons of water per acre.

### Post-Harvest Application

Make a ground application after harvest at 1/3 to 2/3 pint per acre. For control of Canada thistle from after harvest to early fall, apply this product after the majority of basal leaves have emerged but prior to bud stage.

### Spring Application

If needed, make a ground application in the spring no later than 30 days prior to harvest at 1/3 pint per acre.

### Spot Applications

Refer to the product label for Spot Treatments with Hand Held Sprayers.

### Recommendations:

- For difficult-to-control weeds, best results will be achieved with a post-harvest application followed by a spring application.
- This product may be applied to either established or newly planted strawberries. **Note:** If newly planted, **do not** make an application in spring directly after planting; instead, make an application once plants are established and, if needed, a spring application the following year.

### Other Use Restrictions:

- **Do not** apply by aircraft.
- **Chemigation:** Do not apply Stinger through any type of irrigation system.
- **Do not** apply within 6 to 8 hours of expected rainfall or irrigation.
- **Do not** compost treated vegetation if compost will be used on sensitive plants.
- Refer to product label for crop rotation intervals.

## **Special Conditions and Risks of Use**

**USE OF Stinger HERBICIDE (THE “PRODUCT”) ON STRAWBERRY (THE “CROP”) MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS. READ AND UNDERSTAND THESE SPECIAL CONDITIONS AND RISKS OF USE BEFORE USING THE PRODUCT ON THE CROP.**

**Corteva Agriscience makes the Product available for use in the manner described in this Special Local Need Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury or loss. The decision to use this Product in the manner described in this Special Local Need Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls, and (iv) any other relevant factors.**

**By purchasing the Product for use, or using the Product, in the manner described in this Special Local Need Labeling, you acknowledge and accept that:**

- (1) you assume all risk of Crop injury, Crop yield reduction and Crop loss;
- (2) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this Special Local Need Labeling;
- (3) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaims all warranties, express or implied, including any implied warranties of fitness for a particular purpose or merchantability;
- (4) Corteva Agriscience disclaims all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to Crop injury, Crop yield reduction and/or Crop loss;
- (5) these Special Conditions and Risks of Use supersede any contrary representations or recommendations by Corteva Agriscience, its agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

**If these Special Conditions and Risks of Use are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container. These Special Conditions and Risks of Use are required by Corteva Agriscience and not specified by the U.S. EPA or the State of Indiana.**

<sup>TM</sup><sup>®</sup> Trademarks of Corteva Agriscience and its affiliated companies

R043-147  
Accepted: 04/27/22  
Initial Printing