



FIFRA 2(ee) Recommendation

EPA Reg. No. 279-9615

**FOR DISTRIBUTION AND USE
ONLY IN CA, FL, GA, IN, MI, NC,
NJ, OR and WA**

**THIS RECOMMENDATION, WHICH CONTAINS
ADDITIONAL DIRECTIONS FOR USE, IS
MADE AS PERMITTED UNDER FIFRA
SECTION 2(ee) AND HAS NOT BEEN
SUBMITTED TO OR APPROVED BY THE US
EPA.**

This recommendation for EXIREL® insect control is valid until December 31, 2030, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

FOR CONTROL OF OBLIQUEBANDED LEAFROLLER IN BLUEBERRIES

Directions for Use

Pests Controlled, Application Rates, Method, Timing, and Resistance Management:

EXIREL® insect control is recommended for control of obliquebanded leafroller (*Choristoneura rosaceana*) in blueberries.

Apply EXIREL insect control at 10 – 20.5 fl. oz./A (0.065 – 0.133 LB AI/a) as a foliar spray.

Thorough coverage is essential to achieve best results. Select a spray volume appropriate for the size of plants and density of fruit and foliage.

Obliquebanded leafroller (*Choristoneura rosaceana*):

For overwintering larvae, apply in the spring at first sign of active feeding.

For summer generations, apply just prior to or at the beginning of egg hatch. Leafroller feeding stops after ingestion of treated foliage, however, during periods of cold weather when leafrollers are inactive, it may take several days to achieve complete control.



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104

Resistance Management for Obliquebanded Leafroller:

For resistance management purposes, only apply EXIREL insect control (or other Group 28 insecticides) to one generation of obliquebanded leafroller per year. Application(s) to other generations of obliquebanded leafroller should be with an effective product with a different mode of action (i.e. a product with a different IRAC group number).

See the RESISTANCE MANAGEMENT section of the EXIREL insect control Federal label for additional guidance regarding insecticide resistance management.

See 'Tank Mixtures and Crop Safety' section on the EXIREL insect control label for more information and for guidance regarding crop response for tested tank mixtures.

USE RESTRICTIONS

Do not apply more than 0.4 lb a.i. of cyantraniliprole-containing products per acre per calendar year.

The minimum retreatment interval is 5 days.

The preharvest interval is 3 days.

The field re-entry interval is 12 hours.

R-4461 021126 07-07-20

FMC, the FMC logo and EXIREL are Trademarks of FMC Corporation or an affiliate.
©2020 FMC Corporation



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104