

Product Bulletin



Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Fontelis®

EPA Reg. No. 352-834

Special 2(ee) Recommendation†

(For Distribution and Use Only in the State of California)

For Suppression of Hull Rot Caused by *Rhizopus Stolonifer* In Almonds – Fontelis Plus FRAC Group 3, Group 11 or Group 19 Fungicides for Distribution and Sale Only in the State of California

Refer to product label for Fontelis for general use precautions, mixing directions, application directions, and restrictions.

ATTENTION

†This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Fontelis® fungicide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Fontelis according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Fontelis fungicide.

Directions for Use

Use Rate and Application Timing

For suppression of hull rot, apply Fontelis at 20 fluid ounces/acre with one of the following: a FRAC Group 3 fungicide such as tebuconazole at 0.225 lb ai/Acre or difenoconazole at 0.114 lb ai/Acre; a FRAC Group 11 fungicide such as azoxystrobin at 0.244 lb ai/Acre; or a FRAC Group 19 fungicide such as polyoxin-D at 0.044 lb ai/Acre (0.7 oz ai/Acre).

For best results, application timing should be at 1-5% hull split.

Precautions

Read and follow all label instructions on timing, precautions and warning for the companion product before using this tank mixture. Follow the most restrictive labeling. See the full Federal Fontelis label for all applicable use directions and restrictions pertaining to almond.

™® Trademarks of Corteva Agriscience and its affiliated companies

R613-028

Issued: 05/12/23

Replaces: R613-008