

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**DANGER CORROSIVE:** Do not enter an enclosed area without proper respiratory protection. Causes irreversible eye damage and skin burns. May be fatal if inhaled or absorbed through skin. Harmful if swallowed. Do not breathe vapors or spray mist. Do not get in eyes, on skin, or on clothing. Wear goggles and face shield and rubber gloves when handling. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash before reuse.

**Physical or Chemical Hazards:** **Strong oxidizing agent.** Corrosive. Mix only with water at ambient (room) temperature. Product must be diluted in accordance with label directions prior to use. This product is not combustible; however, at temperatures exceeding 156°F, decomposition occurs releasing oxygen. The oxygen released could initiate combustion. **Environmental Hazards:** This pesticide is toxic to birds, fish and aquatic invertebrates. Caution should be used when applying indoors because pets may be at risk. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of the National Pollution Discharge System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product into sewer systems without previously notifying the local sewage plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

Storage and Disposal

**Storage:** Never return this product to the original container after it has been removed. Avoid all contaminants, especially dirt, caustic, reducing agents, and metals. Contamination and impurities will reduce shelf life and can induce decomposition. In case of decomposition, isolate container, douse container with cool water and dilute this product with large volumes of water. Avoid damage to containers. Keep container closed at all times when not in use. **Keep container out of direct sunlight.** To maintain product quality, store at temperatures below 86°F. **Procedure for Leak or Spill:** Stop leak if this can be done without risk. Shut off ignition sources: no flames, smoking, flares, or spark producing tools. Keep combustible and organic materials away. Flush spilled material with large quantities of water. Undiluted material should not enter confined spaces. **Pesticide Disposal:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or Hazardous Waste representative at the nearest EPA Regional Office for guidance. If material has been spilled, an acceptable method of disposal is to dilute with at least 20 volumes of water followed by discharge into suitable treatment system in accordance with all Local, State and Federal environmental laws, rules, regulations, standards, and other requirements. Because acceptable methods of disposal may vary by location, regulatory agencies should be contacted prior to disposal. Micro Blast, which is to be discarded, should be disposed of as hazardous waste after contacting the appropriate Local, State or Federal agency to determine proper procedures. **Container Disposal:** Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Offer for recycling if available. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Directions For Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SANITIZATION

**Sanitizing Food Contact Surfaces:** Sanitize with a concentration of 0.7-2.0 fl. oz. of this product dissolved in 10 gallons of water (93-260 ppm active peroxyacetic acid). Use immersion, coarse spray or circulation techniques as appropriate to the equipment. All surfaces should be exposed tot sanitizing solution for a period of at least 60 seconds or more if specified by a governing code. Drain thoroughly and allow to air dry. Do not rinse.

**Sanitization of Conveyors and Equipment for Meat, Poultry, Seafood, Fruit, Nuts and Vegetables:** This product is effective against the gram negative and gram positive organisms Staphylococcus aureus and Escherichia coli. For use in the static or continuous sanitizing, washing or rinsing of conveyors, slicers, saws and equipment, apply a solution of this product using a recommended 0.7-2.0 fl. oz. per 10 gallons of water (93-260 ppm active peroxyacetic acid). Apply sanitizer solution to the return portion of the conveyor or equipment using coarse spray or similar means of wetting surfaces, so as to prevent puddling. Allow sanitizer to thoroughly wet surface for a minimum of 60 seconds contact time. No rinse is needed.

**Final Bottle or Container Rinse:** This product may be used as a final sanitizer rinse for returnable and non-returnable bottles or containers at 93-260 ppm active peroxyacetic acid (0.7-2.0 fl. oz. Micro Blast dissolved in 10 gallons of water). The container should be drained as much as is practical prior to filling operations.

**Combination Disinfection and Cleaning:** Micro Blast is effective against Staphylococcus aureus and Salmonella choleraesuis at 1.0 oz of this product per 10 gallons of water (130 ppm active peroxyacetic acid) in hard water (400 ppm as CaCO3) and 5% organic soil on hard nonporous surfaces. For heavily soiled areas a pre-cleaning step is required. Apply solution with a mop, cloth, sponge, brush, or by soaking, spraying, or immersion so as to wet all surfaces thoroughly. Allow to remain wet for 10 minutes, then remove excess solution and entrapped soil with a clean wet mop, cloth, wet vacuum pickup or by draining. Prepare a fresh solution daily or when it becomes soiled or diluted.

**REVERSE OSMOSIS (RO) NANO, AND ULTRA FILTRATION CLEANING:** Micro Blast may be used in the sanitization of nano filtration (NF) and ultra filtration (UF) and reverse osmosis (RO) membranes and their associated piping systems. This product is not for use in kidney dialysis equipment. Do not use the intermittent or continuous dosing methods for nano or ultra-filtration food or drinking water applications. This product may not totally eliminate all vegetative microorganisms in RO or NF or UF membranes and their associated piping systems due to their construction or assembly, but can be relied upon to reduce the number of microorganisms to acceptable levels when used as directed. Prior to using this product check with membrane

Micro Blast  
(ANTIMICROBIAL SOLUTION)

FOR USE IN ORGANIC PRODUCTION:

**Micro Blast is a peroxyacetic acid-based microbiocide developed for Equipment Sanitizing and Disinfection, and Bacteria, Slime and Odor Control in: Fruit and Vegetable Process Water, and Bacteria and Algae Control in Recirculating and Agricultural Water Systems.**

ACTIVE INGREDIENT:

Peroxyacetic Acid 15.0%  
Hydrogen Peroxide 22.0%

INERT INGREDIENTS

TOTAL 100.0%

EPA Registration No: 63838-2-81811 EPA Est. No. 40270-AZ; 63838-CA-01; 27706-ID; 60156-IL-01

Before Using This Product, Please Read This Entire Label Carefully.

KEEP OUT OF REACH OF CHILDREN  
DANGER  
FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.  
IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.  
IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses of present after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.  
IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth. Call a poison control center or doctor for treatment advice.  
NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

manufacturer to confirm compatibility of membranes with various types or concentrations of peroxyacetic acid solutions.

**Batch Sanitation of NF, UF and RO Systems:** Isolate incompatible equipment such as carbon filters and ion exchangers. Clean system with an appropriate cleaner and follow with RO permeate water or potable water. Remove mineral deposits if necessary with an acidic cleaner and rinse as before. Fill entire system with water and add up to 0.5% of this product by volume. This will equal 680 ppm peroxyacetic acid and 1000 ppm hydrogen peroxide. Recirculate the sanitizing solution through the piping and membrane system at 20°C for 10 minutes minimum, or up to 4 hrs, depending on the severity of the cleaning to be done. Open and close process valves and solenoids to be sure all parts are in contact with the solution. Rinse the system with RO permeate or potable water until residual peroxygen conc. is below 1 ppm  
**Continuous or Intermittent Addition:** For continuous addition (dosing) for RO systems, use 2-5 ppm of active peroxyacetic acid, which equals 1.5-3.7 fl. oz. of this product per 1000 gallons of process water. For occasional intermittent feed, do not exceed 93 ppm active peroxyacetic acid, which equals 0.7 fl. oz. of this product per 10 gallons of feed water. Continuous or intermittent dosing of this product is not allowed for use in NF or UF systems for on-line food or drinking water applications.

**BIOFOULING CONTROL IN PULP, PAPER AND PAPERBOARD MILL AND WATER SYSTEMS:** (not for use in California) For use in the manufacture of paper and paperboard intended for food or non food contact. Micro Blast can be used to control bacteria and fungi in paper, paperboard or non-woven process water and influent water systems. Suitable dosing points include but are not limited to: stock chests, pulpers, the white water loop, white water storage systems and influent water streams.

**Influent Water Systems:** Micro Blast should be fed continuously to incoming fresh water streams (nonpotable use only) at dosages ranging from 0.11- 2.0 lbs (1.5-27 fl. oz) of this product per 1000 gallons of raw or process water (2.0-36 ppm peroxyacetic acid). Adjust dosage as necessary to maintain microbiological control. **Mill Process Waters:** Intermittent Feed - Micro Blast may be fed intermittently (for example:

2-3 hours per 8 hour shift) at dosages ranging from 0.5 lbs to 1.2 lb (7-16 fl. oz.) of this product per ton (dry basis) of pulp or paper produced. This dosage is equivalent to 37-90 ppm peroxyacetic acid. Repeat as necessary when the peroxyacetic acid concentration reaches less than 2 ppm.  
**Continuous Feed** – Micro Blast should be fed continuously at dosages ranging from 0.11-1.2 lbs (1.5- 16 fl. oz) of this product per ton (dry basis) of pulp or paper produced. This dosage is equivalent to 8.0-90 ppm peroxyacetic acid. **Shock (slug) Dose** - Micro Blast may be used to shock dose systems requiring a high level of biofouling control. Use rates ranging from 1-8 lbs (13.5-108 fl. oz.) of this product per ton (dry basis) of pulp or paper produced may be necessary. This dosage is equivalent to 75-600 ppm peroxyacetic acid. Shock dose every 1-3 hrs as necessary until biofouling control is evident. Thereafter, revert to continuous or intermittent feed methods.

CONTROL OF SLIME FORMING BACTERIA AND BIOFOULING IN COOLING WATER SYSTEMS (COOLING TOWERS, EVAPORATIVE CONDENSERS, AIR WASHERS) AND ORNAMENTAL OR RECREATIONAL WATER FEATURES.

Severely fouled systems should be cleaned before adding the Micro Blast solution. Micro Blast should be added in the water system directly, and not mixed with any other chemicals or additives. Never add this product into any feeding device, such as shot feeders, filter housings, by-pass feeders, or miscellaneous piping of any kind, because dangerous acute decomposition can occur. Discontinue the use of chlorine or bromine products prior to using Micro Blast. Contamination with other chemicals could result in product decomposition. Add Micro Blast only to water at a point in the system where uniform mixing and even distribution will occur. For shock (slug) treatment for moderately to severely fouled systems add 5-20 fl. oz. of this product per 1000 gallons of process water (7-27 ppm peroxyacetic acid). Repeat as necessary until microbiological control is evident. Thereafter, to maintain control use 1.5- 7.5 fl oz of Micro Blast per 1000 gallons of process water (2-10 ppm of peroxyacetic acid) as a continuous treatment method. Continuous dosing methods usually require 1.5-5 fl. oz. per 1000 gallons of water (2-7 ppm peroxyacetic acid) to achieve adequate results. Intermittent dosing treatment usually require dose cycles of a minimum once per day, up to 6 times per 24 hours. Recommended rates for intermittent dose cycles are 5-10 fl. oz of Micro Blast per 1000 gallons of process water (7-14 ppm peroxyacetic acid).

**AIR WASHERS:** This product may be used to control bacteria and biofouling in industrial air washing/scrubbing systems. The air washer must have operational and effective mist elimination systems. Prior to use of this product, heavily fouled systems must be pre-cleaned using the appropriate cleaner. Continuous dosing methods will require 2-7 ppm and intermittent dosing methods require 7-14 ppm (as peroxyacetic acid), as described in the previous 2 paragraphs, depending on the type of system and the level of microbiological control desired.

**TREATMENT OF FRUIT AND VEGETABLE PROCESS WATER SYSTEMS:** Micro Blast can be used in water or ice that contacts raw or fresh, post-harvest or further processed fruits and vegetables for the control of bacteria and fungi.

**Batch, Continuous or Spray System Processes:** Fill vessel containing fruits and vegetables with known amount of water. Ensure that water is circulating in vessel if using the submersion method. Add this product to no more than 533 ppm (wt/wt) total product (80 ppm residual peroxyacetic acid) to the use solution. This can be accomplished by initially adding 53.3 grams (47.3 ml) of this product per 100 liters of water, or 1.0 fl. oz. per 16.4 gallons of water. The fruits and vegetables can be continuously sprayed (using coarse spray) or submerged (dipped) in the resulting solution. Periodic or continuous addition of this product to maintain the required concentration may be added as necessary. Contact time of 60 seconds is recommended to insure efficacy. A potable water rinse is not required.

**AGRICULTURAL OR HORTICULTURAL USES:** There is a Restricted-Entry-Interval of zero (0) hours after the use of this product. This product should never be mixed or combined with any other pesticide or fertilizer. Upon soil contact this diluted product decomposes rapidly to oxygen, carbon dioxide and water. This product may be harmful to fish if exposed on a continuous basis at concentrations of 1 ppm or more of active peroxyacetic acid. Meter this product into pressurized pipes using a plastic or stainless steel injection/backflow device installed far enough upstream from the equipment to insure thorough mixing. For open flowing bodies of water, apply this product as far upstream as possible to allow adequate mixing prior to the flow entering any larger body of water. If open pouring of this product is required, pour product as close to the surface of the water as possible to reduce odor exposure.

**Treatment of Agricultural or Irrigation Water Systems** (sand filters, humidification systems, storage tanks, ponds, reservoirs, canals): For the control of sulfides, odor, slime and algae in water systems, apply this product at 2-10 ppm active peroxyacetic acid. This feed rate equals 15-75 fl. oz per 10,000 gallons of water. Repeat dose as necessary to maintain control, which will vary with seasonal conditions. For prevention of algae, some systems may require continuous low level dosing during warm sunny periods (2-5 ppm peroxyacetic acid). Drip Irrigation Systems: To clean slime and algae from drip system filters, tapes and emitters, meter this product at the rate of 7.5-15 fl. oz per 1000 gallons of water (10-20 ppm peroxyacetic acid). When required during normal irrigation cycles, use this product at the recommended dose for a minimum of 30 minutes. Thereafter, the irrigation cycle should be discontinued and the line should not be flushed.

D.O.T. and Haz Mat Information: UN3109, Organic Peroxide, Type F, Liquid (peroxyacetic acid), 5.2, (8), PGII

Manufactured By:

Meras Engineering, 601 Van Ness Ave, #E3-725, San Francisco, CA 94102  
Rev: #033-CalFed-v5-12-09-8x14

Lot #: (may show elsewhere) \_\_\_\_\_

Net Contents: \_\_\_\_\_gal/lbs

