

**RESTRICTED USE PESTICIDE
DUE TO ACUTE TOXICITY
FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS– NOT TO BE USED
BY UNCERTIFIED PERSONS WORKING UNDER THE SUPERVISION OF A CERTIFIED
APPLICATOR.**

syngenta.

FLORA

Section 24(c) Special Local Need Label

PARAQUAT DICHLORIDE

GROUP

22

HERBICIDE

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF NORTH CAROLINA

**Gramoxone® SL 3.0
For Control of Weeds in Clary Sage**

**EPA Reg. No. 100-1652
EPA SLN No. NC-220003**

**This label expires and must not be distributed or used in accordance with this SLN
registration after December 31, 2027**

**KEEP OUT OF REACH OF CHILDREN / MANTENER FUERA DEL ALCANCE DE LOS
NIÑOS.**



**DANGER / POISON
PELIGRO / VENENO**

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and main EPA-registered label must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the main EPA-registered label.

Gramoxone SL 3.0 – Fall Postemergence

Crop	Rate Per Acre	Use Pattern	Spray Volume Per Acre	Precautions and Comments	
Clary Sage	2-24 fl oz/A, depending on crop size	Fall, Postemergence	Ground: Apply in a minimum of 10 gallons, \geq 15 gallons will provide more thorough coverage of weeds. Air: 5 gallons	<u>Size of Sage (diameter)</u>	<u>Application Rate (fl oz/A)</u>
				2 inches 4 inches 6 inches 12 inches	2-4 4-8 8-16 12-24
				<ul style="list-style-type: none"> • Gramoxone SL 3.0 will injure clary sage when applied postemergence. Grower and/or applicator will accept responsibility for crop injury or crop kill. • During periods of high temperature, high humidity, and/or if Gramoxone SL 3.0 is being tank-mixed with other products the application may be made without NIS. Weed control may be reduced if applied without NIS. • Allow a minimum of 10 days between applications. 	

Gramoxone SL 3.0 – Dormant Season

Crop	Rate Per Acre	Use Pattern	Spray Volume Per Acre	Precautions and Comments
Clary Sage	12-32 fl oz/A	Dormant Season	Ground: Apply in a minimum of 10 gallons, \geq 15 gallons will provide more thorough coverage of weeds. Air: 5 gallons	<ul style="list-style-type: none"> • For suppression of weeds such as Italian ryegrass, prickly lettuce, groundsel, chickweed, downy brome and bluegrass. • Apply when crop is dormant before spring growth begins and when weeds are less than 6" tall.

Add the Following

Add a nonionic surfactant (NIS) containing 80% or more surface-active agent at a minimum of 0.25% v/v (1 quart per 100 gallons) of the finished spray volume for ground applications. For aerial applications, add a nonionic surfactant at 0.25% v/v (1 qt per 100 gallons) of the finished spray volume. See Fall Postemergence section of this label for exceptions to addition of non-ionic surfactant.

Precautions, Restrictions, and Comments

- DO NOT use the treated Clary Sage crop for food or feed.\
- DO NOT allow livestock to graze in treated areas.
- DO NOT feed livestock hay or threshings from treated fields.
- DO NOT exceed 3.0 pts./A in clary sage.
- DO NOT exceed 1.125 lb ai/A/year of paraquat containing products for all combined uses.
- DO NOT apply when weather conditions favor drift from treated areas.
- This pesticide may only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, non-target plants) is minimal (i.e., when the wind is blowing away from the sensitive area. Make aerial or ground applications when the wind velocity favors on target product deposition, (approximately 3-10 mph). Do not apply when wind velocity exceeds 10 mph.
- For aerial applications, the spray boom should be mounted on the aircraft to minimize drift caused by wingtip or rotor vortices. Boom length must not exceed 65% of wingspan for airplanes or 75% of the rotor diameter for helicopters.
- Use nozzles and pressure that will produce medium to coarse size droplets.
- Follow spray volumes listed. Spray volumes should be increased as necessary to obtain complete coverage of the target weed or plant without runoff from the foliage.
- DO NOT apply this product through any type of irrigation system.
- DO NOT enter treated field within 24 hours of application.

©2022 Syngenta

Gramoxone® and the Syngenta logo are trademarks of a Syngenta Group Company

24(c) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Label Code: NC1652059AA0722