



OxiDate® 5.0

FIFRA SECTION 2(ee) USE RECOMMENDATION

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency.

ISSUE DATE:

June 12, 2023

EXPIRATION DATE:

December 31, 2026

**FOR USE AND
DISTRIBUTION ONLY
IN THE STATE OF:**

**CO, CT, DE, IA, ID,
IL, IN, KS, MA, MD,
ME, MI, MN, MO,
MT, NC, ND, NE, NH,
NJ, NM, NV, NY, OH,
OK, OR, PA, RI, SC,
SD, UT, VA, VT, WA,
WI, WV, WY**

EPA REGISTRATION NO. 70299-28

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Crop: Legume Vegetables (succulent or dried) Including, but not limited to: bean (*Lupinus* spp., *Phaseolus* spp., *Vigna* spp.), broad bean, chickpea, dry beans, lima beans, peas, edible podded legume vegetables, lentil, pea (*Pisum* spp.)

Disease: Sclerotinia White Mold

Apply OxiDate 5.0 at dilution rates of 1:256-1:100. Begin applications of OxiDate 5.0 when conditions favor the development of disease, or at first sign/symptom of disease, then continue use as needed. Under moderate to severe disease pressure/conditions, reduce intervals and increase rates of application until control is achieved.

Application Rates and Directions: OxiDate 5.0 works immediately on contact for control of plant diseases. Good coverage and wetting of the foliage is required. Apply OxiDate 5.0 in solution quantities (gallons per acre) great enough to ensure uniform coverage of upper and lower foliage, stems, branches and stalks. Final spray solution volume will depend on crop type, canopy size and/or growth stage. For increased coverage and penetration of spray, use a compatible non-ionic wetting agent/surfactant. For drift reduction and to aid spray deposition, the use of a proper adjuvant is recommended. Before widespread use, run a plant sensitivity test when considering use of spray concentrations greater than 0.39% v/v (1:256) by following instructions under **Plant Sensitivity Testing**.

See **Application Rates and Directions Chart** for disease list and additional instructions.

See **Solution Preparation, Compatibility, and Plant Sensitivity Testing** for additional information and instructions.

DILUTION RATE CHART

Amount of OxiDate 5.0 per Acre						
Dilution Rate of OxiDate 5.0	Spray Volume (Gallons/Acre)					
	5	15	20	50	100	500
1:100 (1.0% v/v)	6.4 fl. oz.	19.2 fl. oz.	26 fl. oz.	64 fl. oz.	1.0 gal.	5.0 gal.
1:256 (0.39% v/v)	2.5 fl. oz.	7.5 fl. oz.	10 fl. oz.	25 fl. oz.	50 fl. oz.	250 fl. oz.



BioSafe Systems, LLC

22 Meadow Street
East Hartford, CT 06108

www.BioSafeSystems.com

1-888-273-3088

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use which are recommended by BioSafe Systems, LLC. which may not appear on the package label. Read and carefully observe the precautionary statements plus all the other information appearing on the product label.

This product bulletin must be in the possession of the user at the time of pesticide application.