

**SUPPLEMENTAL LABELING**

**RESTRICTED USE PESTICIDE  
DUE TO ACUTE TOXICITY**  
TO BE USED BY CERTIFIED APPLICATORS ONLY – NOT TO BE USED BY UNCERTIFIED PERSONS  
WORKING UNDER THE SUPERVISION OF A CERTIFIED APPLICATOR, EXCEPT THAT UNCERTIFIED  
PERSONS MAY TRANSPORT CONTAINERS AS PROVIDED UNDER DIRECTIONS FOR USE.

**Syngenta Crop Protection, LLC**  
P. O. Box 18300  
Greensboro, North Carolina, 27419-8300  
SCP 1652A-S4A 0822

PARAQUAT DICHLORIDE	GROUP	<b>22</b>	HERBICIDE
---------------------	-------	-----------	-----------

**Gramoxone® SL 3.0**  
**HERBICIDE**

EPA Reg. No. 100-1652

A Weed, Grass, and Harvest Aid Desiccant/Defoliant Herbicide

**Active Ingredients:**

Paraquat dichloride* .....	43.9%
Other Ingredients:	56.1%
<b>Total:</b>	<b>100.0%</b>

\*CAS No. 4685-14-7

Gramoxone SL 3.0 is formulated as a liquid containing 3.0 pounds paraquat cation per gallon as 4.134 pounds paraquat dichloride per gallon.  
Gramoxone SL 3.0 contains alerting agent (odor), emetic, and dye.

**KEEP OUT OF REACH OF CHILDREN / MANTENER FUERA DEL ALCANCE DE LOS NIÑOS.**



**DANGER / POISON**  
**PELIGRO / VENENO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)  
See additional precautionary statements and directions for use inside booklet.

All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Gramoxone SL 3.0 as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



**DIRECTIONS FOR USE  
RESTRICTED USE PESTICIDE**

Cucurbits, DE Only

<b>Crops (Including all cultivars, varieties, and/or hybrids of these)</b>			
Cucumber Melons		Pumpkin Squash	
<b>Target Weed</b>	<b>Rate pt/A</b>	<b>Application Timing</b>	<b>Use Directions</b>
Refer to <b>Section 7.0.</b>	1.3	Postemergence (Directed Spray/Shield Spray)	<p>For use in <b>Delaware Only</b> for control or suppression of emerged weeds between rows after crop establishment.</p> <p>Apply as a directed/shielded spray application by air in a minimum of 20 gal/A of spray volume.</p> <p>Use precision directed spray equipment adjusted to prevent spray contact with crop plants.</p> <p>Do not exceed 30 psi nozzle pressure or spray under conditions which may cause excessive drift.</p> <p>Apply when weeds are 6" or less in height and actively growing.</p> <p>Add either: 1) A nonionic surfactant containing 75% or more surface-active agent at 0.25% v/v (2 pt per 100 gal), of the finished spray volume for ground applications, <b>OR</b> 2) Crop Oil Concentrate (non-phytotoxic) containing 15-20% approved emulsifier, at 1.0% v/v (1 gal per 100 gal) of the finished spray volume.</p> <p>See <b>Section 4.4.4</b> of the Gramoxone SL 3.0 for adjuvant recommendations.</p>
<b>Tank Mix Options:</b>			
<ul style="list-style-type: none"> <li>Gramoxone SL 3.0 can be tank mixed with other herbicides labeled for use on cucurbits applied by the same methods and at the same timings. Refer to <b>Section 4.4.2</b> for tank-mix precautions in the Gramoxone SL 3.0 label.</li> </ul>			
<b>Resistance Management:</b>			
<ul style="list-style-type: none"> <li>Refer to <b>Section 3.1.</b></li> </ul>			
<b>Precaution:</b>			
<ul style="list-style-type: none"> <li>Do not allow spray to contact green stems or foliage, as injury may result.</li> </ul>			
<b>USE RESTRICTIONS</b>			
<ol style="list-style-type: none"> <li>Refer to <b>Section 6.1</b> for additional product use restrictions.</li> <li><b>Maximum Single Application Rate:</b> 1.3 pt/A</li> <li><b>REI:</b> 48 hours</li> <li><b>Minimum Application Interval:</b> 7 days</li> </ol>			

5. **Maximum Annual Rate:** 3.9 pt/A
  - a. **DO NOT** exceed 1.5 lb cation/A/year of paraquat-containing products.
6. **DO NOT** apply after bloom.
7. **DO NOT** make more than 3 applications per year.
8. **DO NOT** use on soils with less than 5% clay.
9. **DO NOT** apply this product through any type of irrigation system.
10. **DO NOT** allow animals to graze in treated areas.
11. **Pre-Harvest Interval (PHI):** Harvest at normal crop maturity.

Gramoxone®, and the SYNGENTA Logo are Trademarks of a Syngenta Group Company.

©2022 Syngenta

For non-emergency (e.g., current product information), call  
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, North Carolina, 27419-8300