

FIFRA Section 2(ee) Recommendation

For control of black rot (*Xanthomonas campestris* pv. *campestris*) in Brassica in all States **except** HI, NY, and TX

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the US Environmental Protection Agency.

All applicable directions, restrictions, and precautions on the EPA-registered labels must be followed. Please contact your Certis USA regional sales manager to determine the specific requirements for FIFRA §2(ee) recommendations in your respective state.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Crop: Brassica (Cabbage)

Application Rate: 4.5 oz/100 gal in sufficient water to provide uniform coverage

Apply LifeGard® WG 3-5 days before suspected infection event. Reapply every 7-14 days to maintain control as needed. Thorough spray coverage of plant foliage will result in best performance.

Crop-Specific Restrictions:

Do not apply to plants that are stressed due to drought, excessive moisture, excessive hot or cold temperatures, herbicide injury, or other environmental stresses.

Refer to EPA-approved label for complete list of **USE RESTRICTIONS**.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the precautionary statements plus all other information appearing on the product labels. **For Utah and Washington, this recommendation is not required to be in the possession of the user at the time of pesticide application. For all other states, this recommendation must be in the possession of the user at the time of pesticide application.**

