Drexel Simazine 4L is a state restricted use pesticide and is to be distributed only by licensed pesticide dealers. Only certified applicators or persons under the direct supervision of a certified applicator may apply Drexel Simazine 4L.

SECTION 24(c) SPECIAL LOCAL NEED LABEL FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON



Drexel. Simazine 4L

Herbicide

EPA Reg. No. 19713-60 EPA SLN No. WA-090015

This label for Drexel Simazine 4L expires and must not be distributed in or used in accordance with this SLN registration after December 31, 2025.

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements and precautions on the EPA registered label.

DIRECTIONS FOR USE

Alfalfa Grown for Seed Only / Annual weeds (such as Clasping pepperweed, Downy brome, Flixweed, Mustards, Prickly lettuce, and seedling Shepherdspurse)

Make a single application of 1.6 to 3.2 pints (0.8 to 1.6 lb. active ingredient) of Simazine 4L per acre. Use lower rates on course soils and higher rates on fine soils. Apply as a broadcast spray in a minimum 20 gallons of water per acre by ground application only. Apply between Fall (after harvest) and early Spring (before Spring growth starts). Do not apply if the ground is frozen. Alfalfa may be actively growing, semi-dormant or dormant when the application is made.

Restrictions / Precautions

- Do not enter or allow worker entry into treated areas during the Restricted Entry Interval (REI) of 12 hours.
- Do not apply Simazine 4L to Sand, Loamy sand or gravelly areas of the field. Do not apply to areas of the field where water will accumulate.
- Make only one application of Simazine 4L per year.
- For use under this SLN label, do not apply Simazine 4L through any type of irrigation system.
- · Aerial application is prohibited.
- Apply Simazine 4L only to Alfalfa grown for seed that has been established for at least one year.
- Land treated with Simazine 4L should not be planted to any crop except Corn until the following year as injury may occur. After the last harvest of the treated crop, plow and thoroughly till the soil in the Fall or Spring to minimize possible injury to Spring-seeded rotational crops. Avoid rotating to crops susceptible to injury from Simazine.

- This pesticide is toxic to aquatic invertebrates. Simazine 4L should not be used in accordance with this SLN label where impact on listed threatened or endangered species is likely. You may refer to the WSDA Natural Resources Assessment Section website (agr.wa.gov/departments/land-and-water/natural-resources/endangered-species-program) for additional information related to pesticide use and endangered species protection. Information from EPA about no-spray buffers zones is available at agr.wa.gov/departments/land-and-water/natural-resources/endangered-species-program/buffers. Pesticide applicators may use the Washington Department of Fish and Wildlife's SalmonScape mapping tool at apps.wdfw.wa.gov/salmonscape/ to determine if listed salmonid species occur in or near a proposed application site. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.
- Product may not be mixed/loaded or used within 50 feet of all wells, including abandoned wells, drainage wells and sink holes.
- Product must not be mixed or loaded within 50 feet of intermittent streams or rivers, natural or impounded lakes and reservoirs. Product must not be applied within 66 feet of the points where field surface water runoff enters perennial or intermittent streams and rivers or within 200 feet of natural or impounded lakes and reservoirs. If this product is applied to highly erodible land, the 66 foot buffer or setback from runoff points must be planted to crop or seeded with grass or other suitable crop.
- All Alfalfa seed screenings shall be disposed of in such a way that they cannot be distributed or used for human food or animal feed. The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records to the WSDA immediately upon request. Conditioner disposal records shall consist of documentation of on-farm disposal, disposal at a controlled dumpsite, incinerator, composter or other equivalent disposal site and shall include the lot numbers, amount of material disposed of, the growers(s), and the date of disposal.
- No portion of the Alfalfa plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes.
- Alfalfa seed shall bear a tag or container label which forbids the use of the seed for human consumption or animal feed.
- Alfalfa seed may not be distributed for human consumption or animal feed.



The DREXEL logo is a registered trademark of Drexel Chemical Company.

WSDA Container Disposal Guidance

Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA Waste Pesticide Program web site at agr.wa.gov/departments/pesticides-and-fertilizers/pesticides/waste-pesticide. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.



The DREXEL logo is a registered trademark of Drexel Chemical Company.