

RESTRICTED USE PESTICIDE
DUE TO TOXICITY TO NON-TARGET INVERTEBRATES, MAMMALS, AND AQUATIC ORGANISMS
FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 1351A-S9 0720

ABAMECTIN	GROUP	6	INSECTICIDE
-----------	-------	----------	-------------

Agri-Mek® SC Miticide/Insecticide

This supplemental label expires on 7/21/2023 and must not be used or distributed after this date.

Active Ingredient:	
Abamectin ¹ :	8.0%*
Other Ingredients:	92.0%
Total:	100.0%

¹CAS No. 71751-41-2

*Agri-Mek® SC Miticide/Insecticide is formulated as a suspension concentrate and contains 0.7 lb abamectin per gallon.

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

EPA Reg. No. 100-1351

All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Agri-Mek® SC Miticide/Insecticide as permitted according to this Supplemental Labeling, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

syngenta

DIRECTIONS FOR USE

Dried Shelled Pea and Bean (except Soybean) Crop Subgroup 6C

Crops (including all dried cultivars, varieties, and/or hybrids of these)			
Broad bean (dry) Chickpea Guar Lablab bean (hyacinth bean) Lentil <i>Lupinus</i> spp. Grain lupin Sweet lupin White lupin White sweet lupin	<i>Phaseolus</i> spp. Field bean Kidney bean Lima bean (dry) Navy bean Pinto bean Tepary bean Pigeon pea <i>Pisum</i> spp. Field pea	<i>Vigna</i> spp. Adzuki bean Blackeyed pea Catjang Cowpea Crowder pea Moth bean Mung bean Rice bean Southern pea Urd bean	
Target Pest	Rate (fl oz/A)	Application Timing	Use Directions
<i>Liriomyza</i> leafminers pider Mites	1.75 - 3.5	Apply when adult leafminer flies or spider mites are first observed and repeat application, if needed, to maintain control.	<p>Thorough coverage of the crop canopy is essential for optimum results. Inadequate coverage can result in reduced control.</p> <p>Apply this product diluted in a minimum volume of 10 gal/A by ground or 5 gal/A by air.</p> <p>With aerial application, the resulting control of leafminers and spider mites could be less than with ground application.</p> <p>Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.</p>
USE RESTRICTIONS			
1) Maximum Single Application Rate: 3.5 fl oz Agri-Mek SC/A (0.019 lb ai/A) 2) Minimum Application Interval: 6 days 3) Maximum Annual Rate: 10.5 fl oz Agri-Mek SC/A/calendar year a. DO NOT exceed 0.057 lb ai/A/calendar year of abamectin-containing products including all application types (seed treatment, soil, foliar). b. DO NOT make more than 2 sequential applications of Agri-Mek SC or any other foliarly applied abamectin-containing product. 4) DO NOT allow livestock to graze pea or bean forage. 5) DO NOT harvest pea or bean forage or hay for use as livestock feed. For use on cowpeas that are grown only for dry seed. 6) Pre-Harvest Interval (PHI): 7 days			

Edible-Podded Legume Vegetables Crop Subgroup 6A

Crops (including all cultivars, varieties, and/or hybrids of these)			
Jackbean <i>Phaseolus</i> spp. Runner bean Snap bean Waxbean	<i>Pisum</i> spp. Dwarf pea Edible-podded pea Snow pea Sugar snap pea Pigeon pea Soybean (immature seed)	Sword bean <i>Vigna</i> spp. Asparagus bean Chinese longbean Moth bean Yardlong bean	
Target Pest	Rate (fl oz/A)	Application Timing	Use Directions
<i>Liriomyza</i> leafminers Spider Mites	1.75 - 3.5	Apply when adult leafminer flies or spider mites are first observed and repeat application, if needed, to maintain control.	<p>Thorough coverage of the crop canopy is essential for optimum results. Inadequate coverage can result in reduced control.</p> <p>Apply this product diluted in a minimum volume of 10 gal/A by ground or 5 gal/A by air.</p> <p>With aerial application, the resulting control of leafminers and spider mites could be less than with ground application.</p> <p>Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.</p>
USE RESTRICTIONS			
<ol style="list-style-type: none"> 1) Maximum Single Application Rate: 3.5 fl oz Agri-Mek SC/A (0.019 lb ai/A) 2) Minimum Application Interval: 6 days 3) Maximum Annual Rate: 10.5 fl oz Agri-Mek SC/A/calendar year <ol style="list-style-type: none"> a. DO NOT exceed 0.057 lb ai/A/calendar year of abamectin-containing products including all application types (seed treatment, soil, foliar). b. DO NOT make more than 2 sequential applications of Agri-Mek SC or any other foliarly applied abamectin-containing product. 4) DO NOT allow livestock to graze pea or bean forage. 5) DO NOT harvest pea or bean forage or hay for use as livestock feed. 6) Pre-Harvest Interval (PHI): 7 days 			

Succulent Shelled Pea & Bean Crop Subgroup 6B (except cowpea)

Crops (including all cultivars, varieties, and/or hybrids of these)			
<i>Phaseolus</i> spp. Lima bean (green) Broad bean (succulent)	<i>Vigna</i> spp. Blackeyed pea Southern pea	<i>Pisum</i> spp. English pea Garden pea Green pea Pigeon pea	
Target Pest	Rate (fl oz/A)	Application Timing	Use Directions
<i>Liriomyza</i> leafminers Spider Mites	1.75 - 3.5	Apply when adult leafminer flies or spider mites are first observed and repeat application, if needed, to maintain control.	<p>Thorough coverage of the crop canopy is essential for optimum results. Inadequate coverage can result in reduced control.</p> <p>Apply this product diluted in a minimum volume of 10 gal/A by ground or 5 gal/A by air.</p> <p>With aerial application, the resulting control of leafminers and spider mites could be less than with ground application.</p> <p>Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.</p>
USE RESTRICTIONS			
<ol style="list-style-type: none"> 1) Maximum Single Application Rate: 3.5 fl oz Agri-Mek SC/A (0.019 lb ai/A) 2) Minimum Application Interval: 6 days 3) Maximum Annual Rate: 10.5 fl oz Agri-Mek SC/A/calendar year <ol style="list-style-type: none"> a. DO NOT exceed 0.057 lb ai/A/calendar year of abamectin-containing products including all application types (seed treatment, soil, foliar). b. DO NOT make more than 2 sequential applications of Agri-Mek SC or any other foliarly applied abamectin-containing product. 4) DO NOT allow livestock to graze pea or bean forage. 5) DO NOT harvest pea or bean forage or hay for use as livestock feed. 6) Pre-Harvest Interval (PHI): 7 days 			

Tropical and Subtropical, Small Fruit, Inedible Peel, Crop Subgroup 24A

Crops (Including all cultivars, varieties, and/or hybrids of these)			
Aisen	Madras-thorn	Satinleaf	
Bael fruit	Mandruro	Sierra-Leone-tamarind	
Burmese grape	Matisia	Spanish lime	
Cat's eyes	Mesquite	Velvet tamarind	
Inga	Mongongo, fruit	Wampi	
Longan	Pawpaw, small-flower	White apple star	
Lychee			
Target Pest	Rate (fl oz/A)	Application Timing	Use Directions
Twospotted spider mite <i>Liriomyza</i> leafminers Thrips	2.25 - 4.25	<p>Mites: Apply when mites first appear during spring, summer, and/or fall.</p> <p>Leafminers: Apply to protect new growth during spring, summer, or fall.</p> <p>Thrips: Applications targeted for thrips will only control the current generation and must be correctly timed. Apply when economic thresholds have been reached (after egg hatch has begun – preferably early to mid-hatch).</p>	<p>Apply this product diluted in a minimum volume of 50 gal/A with conventional ground application equipment. When using an electro-static sprayer, less than 50 gal/A may be used; however, do not use less than 5 gal/A.</p> <p>Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.</p>
USE RESTRICTIONS			
<ol style="list-style-type: none"> 1) Maximum Single Application Rate: 4.25 fl oz Agri-Mek SC/A (0.023 lb ai/A) 2) Minimum Application Interval: 30 days 3) Maximum Annual Rate: 8.5 fl oz Agri-Mek SC/A/calendar year <ol style="list-style-type: none"> a. DO NOT exceed 0.046 lb ai/A/calendar year of abamectin-containing products including all application types (seed treatment, soil, foliar). b. DO NOT make more than 2 applications of Agri-Mek SC or any other foliarly applied abamectin-containing product per year. 4) DO NOT apply by air. 5) Pre-Harvest Interval (PHI): 14 days 			

Agri-Mek® and the SYNGENTA Logo are Trademarks of a Syngenta Group Company

©2020 Syngenta