

PULL HERE TO OPEN ►

RESTRICTED USE PESTICIDE DUE TO ACUTE INHALATION TOXICITY

FOR SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

THIAMETHOXAM

GROUP 4A

INSECTICIDE

ABAMECTIN

GROUP 6

INSECTICIDE

Escanee QR
para Español



 Avicta[®] Duo COT202

syngenta[®]

Nematicide/Insecticide

A Seed Treatment Product to Protect Cotton Seedlings from Early Season Nematode and Insect Damage

For use only in Syngenta-certified cottonseed treatment facilities and only in seed treatment equipment with closed transfer and application systems. Not for use in hopper box, planter box, slurry box, or other farmer-applied applications.

Active Ingredients:

Abamectin (CAS No. 71751-41-2) 11.2%

Thiamethoxam (CAS No. 153719-23-4) 28.0%

Other Ingredients:

60.8%

Total:

100.0%

Avicta Duo COT202 is a flowable concentrate containing 1.11 pounds of abamectin and 2.78 pounds of thiamethoxam per gallon.

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use in booklet.

EPA Reg. No. 100-1538 EPA Est. 100-NE-001

SCP 1538A-L1C 1025

4247976

Product ID

55887

15 gallons
Net Contents



| FIRST AID | |
|---|--|
| If inhaled | <ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. |
| If swallowed | <ul style="list-style-type: none"> Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. | |
| SYNGENTA HOTLINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372 | |

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING/AVISO

May be fatal if inhaled. Do not breathe vapor. May be fatal if swallowed. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

This product must be used only in seed treatment equipment with closed transfer application systems.

The following personal protective equipment and clothing are required when using this product:

Personal Protective Equipment (PPE)

Seed Treatment Workers

Workers involved with treating the seed (e.g. connecting and disconnecting hoses and transfer pumps, mixing, equipment calibration, etc.) must wear:

- Chemical-resistant gloves made of: barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton™ ≥ 14 mils
- Protective eyewear
- Coveralls worn over long-sleeved shirt and long pants
- Chemical-resistant apron
- Shoes and socks
- A minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.

continued...

PRECAUTIONARY STATEMENTS (continued)

Workers Filling Seed Containers/Baggers and Bag Sewers must wear:

- Chemical-resistant gloves made of: barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils
- Protective eyewear
- Coveralls worn over long-sleeved shirt and long pants
- Shoes and socks
- A minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.

Workers involved with the clean-up and maintenance of seed treatment equipment must wear:

- Chemical-resistant gloves (minimum length 12 inches) made of: barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils. Important: Tape gloves down to the sleeves of the coveralls.
- Protective eyewear
- Chemical-resistant, hooded coveralls worn over long-sleeved shirt and long pants
- Shoes and socks
- A minimum of a NIOSH-approved elastomeric half mask respirator with organic vapor (OV) cartridges and combination N, R or P filters; OR a NIOSH-approved gas mask with OV canisters; OR a NIOSH-approved powered air purifying respirator with OV cartridges and combination HE filters.

Multiple Task Workers must wear:

(Multiple task workers perform multiple tasks in one day such as mixing, product application, filling seed containers/ bagging, bag sewing, and clean-up)

- Chemical-resistant gloves made of: barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils. Important: when performing seed treatment equipment clean-up procedures, gloves must be at least 12 inches long and taped down to the sleeves of the coveralls.
- Protective eyewear
- Long-sleeved shirt and long pants
- When performing seed treatment equipment clean-up procedures, chemical-resistant, hooded coveralls over long-sleeved shirt and long pants are required.
- Shoes and socks
- A minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.

User Safety Requirements

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

User Safety Recommendations

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is toxic to fish and wildlife and highly toxic to aquatic invertebrates. Treated seeds exposed on the soil surface may be hazardous to wildlife. Cover or collect seeds spilled during loading. Do not contaminate water when disposing of equipment washwater or rinsate.

Pollinator Precautions

Abamectin and thiamethoxam are highly toxic to bees exposed to direct treatment or residues in blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Groundwater Advisory

This product has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

RESTRICTED USE PESTICIDE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Restrictions:

- Avicta Duo COT202 is only for use in Syngenta-certified seed treatment facilities: Use in hopper box, planter box, slurry box, or other farmer-applied applications is not permitted.
- Treatment of highly mechanically scarred or damaged seed or seed known to be of low vigor and poor quality may result in reduced germination and/or reduction of seed and seedling vigor. Conduct germination tests with a representative sample of the seed lot before committing the total seed lot to a selected seed treatment. Because seed quality and seed storage conditions are beyond the control of Syngenta, no claims are made to guarantee the germination of seed or propagating material for all crops.
- Seed treated with this product must be visually identifiable from untreated seed by the use of an approved colorant or dye to prevent accidental use of treated seed as food for humans or feed for animals. Refer to 21 CFR, Part 2.25. Any colorant or dye added to treated seed must be cleared for use in accordance with 40 CFR, Part 180.1001. Refer to the section on this label **REQUIRED SEED CONTAINER STATEMENTS** for language that must appear on the tag that is attached to the container of treated seed.

PRODUCT INFORMATION

Avicta Duo COT202 is a cotton seed treatment nematicide/insecticide that contains the active ingredients abamectin and thiamethoxam. When Avicta Duo COT202 is applied to cotton seed, seedlings are protected from early season nematode damage and damage from the insect pests specified in the Crop Use Directions.

To add early season disease protection, the seed treatment fungicides specified in the Crop Use Directions are approved for use on cotton seed and may also be mixed with Avicta Duo COT202. Follow the most restrictive requirements and precautions for all products used in all mixtures. (See **Avicta Duo COT202 in Tank Mixtures**.)

REQUIRED SEED CONTAINER STATEMENTS

User is responsible for ensuring that the seed container meets all requirements under the Federal Seed Act.

The Federal Seed Act requires that the container of seed treated with Avicta Duo COT202 must be labeled with the following statements:

- This seed has been treated with abamectin nematicide and thiamethoxam insecticide.
- Do not use treated seed for feed, food, or oil purposes.

In addition, the U.S. Environmental Protection Agency requires the following statements on the container of cotton-seed treated with Avicta Duo COT202.

- Store away from food and feedstuffs.
- Do not allow children, pets or livestock to have access to treated seeds.
- Wear long-sleeved shirt, long pants, and chemical-resistant gloves made of: barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, polyvinyl chloride (PVC) \geq 14 mils, or Viton \geq 14 mils when handling treated seed.
- **Groundwater Advisory:** This product has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into the groundwater if used in areas where soils are permeable, particularly where the water table is shallow.
- **Pollinator Precautions:** Abamectin and thiamethoxam are highly toxic to bees exposed to direct treatment or residues in blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

- Plant Avicta Duo COT202-treated cottonseed based upon the local planting dates and soil temperatures that are recommended by your state agricultural extension agent.
- Treated seed must be planted into the soil at a depth greater than 1 inch.
- Treated seed exposed on soil surface may be hazardous to wildlife; cover or collect treated seed that are spilled during loading and in areas such as row ends.
- Dispose of all excess treated seed by burying seed away from bodies of water.
- Do not contaminate bodies of water when disposing of planting equipment wash water.
- Dispose of seed packaging in accordance with local requirements.
- Do not use at a rate that will result in more than 0.075 lb thiamethoxam per acre per year (34.0 g ai/A/yr) as a seed treatment application. This seed has been treated with thiamethoxam at 0.375 mg ai/seed, and 0.15 mg of abamectin per seed.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed, and (2) no measurable residues of pesticide remain in the ethanol by-products that are used for agronomic practice.
- Allow a minimum of 45 days after planting Avicta Duo COT202 treated cottonseed before treating the cotton with a neonicotinoid insecticide.
- In areas with a history of high thrips pressure or for cotton grown in North Carolina, South Carolina, and Virginia, apply a foliar insecticide that is approved for thrips control, when cotton is between the 1st and 3rd leaf stage.
- In the event of crop failure or after harvest of cotton grown from Avicta Duo COT202 treated cottonseed, the following crops may be replanted as follows:

| Immediate Plant Back |
|---|
| Alfalfa |
| Cereal Grains: Barley, Buckwheat, Corn (field, pop, and sweet), Oat, Pearl Millet, Proso Millet, Rice (Dry-seeded), Rye, Sorghum, Teosinte, Triticale, Wheat, and Wild Rice |
| Canola |
| Cotton |
| Cucurbit Vegetables Crop Group 9 |
| Fruiting Vegetables Crop Group 8 |
| Head and Stem <i>Brassica</i> Crop Subgroup 5A |
| Leafy <i>Brassica</i> Greens Crop Subgroup 5B |
| Leafy Vegetables (Except <i>Brassica</i> Vegetables) Crop Group 4 |
| Legume Vegetables (Succulent or Dried, Including Soybean) Crop Group 6 |
| Mint: Peppermint and Spearmint |
| Oilseed Crops: Borage, Crambe, Flax Seed, Mustard Seed, and Safflower |
| Onion, Dry Bulb |
| Peanut |
| Potato |
| Root Vegetables Crop Subgroup 1A |
| Strawberry |
| Sunflower |
| Tobacco |
| Tuberous and Corm Vegetables (Except Potato) Crop Subgroup 1D |
| 120-Day Plantback |
| All other crops |
| A cover crop other than the crops listed above that is planted for erosion control or soil improvement, may be planted sooner than the 120-day interval; however, the crop may not be grazed or harvested for food or feed. |

RESISTANCE MANAGEMENT

For resistance management, please note that Avicta Duo COT202 contains both thiamethoxam, a Group 4A insecticide, and abamectin, a Group 6 insecticide. Any insect population may contain individuals naturally resistant to Avicta Duo COT202 and other Group 4A or Group 6 insecticides. The resistant individuals may dominate the insect population if these insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

Thiamethoxam is a systemic insecticide belonging to the neonicotinoid class of chemistry which includes nicotinic acetylcholine receptor (nAChR) agonists. Abamectin belongs to the avermectin class of chemistry which affects nerves and muscles.

To delay insecticide resistance, take the following steps:

- Rotate the use of Avicta Duo COT202 or other Group 4A or Group 6 insecticides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.

Syngenta encourages responsible product stewardship to ensure effective long term control of the insect pests on this label.

For additional information on Insect Resistance Management:

- Contact Syngenta representatives at 1-866-796-4368
- Contact your local Cooperative Extension Service specialist, pest control advisor, or certified crop advisor.
- Visit the Insecticide Resistance Action Committee (IRAC) on the web at: <http://www.irac-online.org/>

ROTATIONAL CROP RESTRICTIONS

In the event of crop failure or after harvest of cotton grown from Avicta Duo COT202 treated cottonseed, the following crops may be replanted as follows:

| Immediate Plant Back |
|---|
| Alfalfa |
| Cereal Grains: Barley, Buckwheat, Corn (field, pop, and sweet), Oat, Pearl Millet, Proso Millet, Rice (Dry-seeded), Rye, Sorghum, Teosinte, Triticale, Wheat, and Wild Rice |
| Canola |
| Cotton |
| Cucurbit Vegetables Crop Group 9 |
| Fruiting Vegetables Crop Group 8 |
| Head and Stem <i>Brassica</i> Crop Subgroup 5A |
| Leafy <i>Brassica</i> Greens Crop Subgroup 5B |
| Leafy Vegetables (Except <i>Brassica</i> Vegetables) Crop Group 4 |
| Legume Vegetables (Succulent or Dried, Including Soybean) Crop Group 6 |
| Mint: Peppermint and Spearmint |
| Oilseed Crops: Borage, Crambe, Flax Seed, Mustard Seed, and Safflower |
| Onion, Dry Bulb |
| Peanut |
| Potato |
| Root Vegetables Crop Subgroup 1A |
| Strawberry |
| Sunflower |
| Tobacco |
| Tuberous and Corm Vegetables (Except Potato) Crop Subgroup 1D |
| 120-Day Plantback |
| All other crops |
| A cover crop other than the crops listed above that is planted for erosion control or soil improvement, may be planted sooner than the 120-day interval; however, the crop may not be grazed or harvested for food or feed. |

MIXING AND APPLICATION INSTRUCTIONS

- Consult the Syngenta Seed Treatment representative or the manufacturer of the application equipment you plan to use regarding the suitability of this application and equipment calibration and operating instructions.
- Always treat seed in a well-ventilated area. Refer to the **Precautionary Statements** section of this label for personal protective equipment requirements and other safety precautions.
- Prepare no more mixture than is needed for the immediate operation.
- Avicta Duo COT202 must be thoroughly recirculated in its original container before using. (See **Mixing Instructions**.)
- Apply Avicta Duo COT202 as a water-based slurry utilizing standard slurry seed treatment equipment that will provide uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of nematode and insect control.

- Thoroughly mix the required amount of Avicta Duo COT202 and other seed treatment products desired with the amount of water required for the slurry treater and the dilution rate. (See **Avicta Duo COT202 in Tank Mixtures**.) Follow the most restrictive requirements and precautions for all products used in mixtures.
- Use an EPA-approved dye or colorant that imparts an unnatural color to the seed.
- Allow seed to dry before bagging /filling containers, and store treated seed away from feed and foodstuffs.

Avicta Duo COT202 in Tank Mixtures:

It is the pesticide user's responsibility to ensure that all products in a tank mix are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, plant back interval from another).

Compatibility:

Before mixing Avicta Duo COT202 with other seed treatment products, determine if all of the products are compatible by conducting a compatibility test:

1. In a clear glass container thoroughly mix all of the intended seed treatment products with the appropriate amount of water, and allow the mixture to sit for one hour.
2. Remix the solution and observe for incompatibility.
3. If compatibility is acceptable, follow the mixing instructions in the **Mixing Instructions** section on this label.

Mixing Instructions:

A. Avicta Duo COT202 Alone

1. To dispense Avicta Duo COT202 into a mix tank, first attach a pump with fittings (example: Micro Matic) that match the fittings on the Avicta Duo COT202 container.
2. Recirculate product. Recirculation time should be calculated based upon the ability of the pump to completely recirculate the volume of the tank 5-10 times within a 30-minute time period.
3. After the initial recirculation, a short recirculation period of 5 minutes per day is recommended.
4. Attach the dispensing hose to a dry-lock type of fitting that is permanently attached to the mix tank lid.
5. Add 1/4-1/2 of the required amount of water to the mix tank.
6. With the agitator running, add the Avicta Duo COT202 to the tank.
7. Continue agitation while adding the remainder of the water.
8. Begin application of the solution after the Avicta Duo COT202 has completely dispersed into the mix water.
9. Maintain agitation until all of the mixture has been applied.

B. Avicta Duo COT202 - Tank Mixing Instructions:

1. Add 1/2 of the required water to the mix tank and turn on the agitation. Mechanical agitation is preferred.
2. If using wettable powders and/or products in water-soluble packaging, add water-soluble packages first to clean water and allow them to completely disperse before adding Avicta Duo COT202 or other products.
3. Allow each tank mix partner to completely disperse before adding the next product.
4. Add the remaining amount of water and agitate.
5. Maintain agitation until the entire slurry mixture has been used.

CROP USE DIRECTIONS

Avicta Duo COT202 on Cottonseed

Avicta Duo COT202 is a seed treatment for the seed of delinted cotton. Avicta Duo COT202 will provide cotton seedlings early season protection from damage caused by nematodes and the insects specified in the following table. Tank mixing Avicta Duo COT202 with Dynasty® CST fungicide will add early season disease protection. Refer to the following table for application directions.

Application

| Pests /Diseases | Amount of Product to Apply* | Instructions |
|---|---|---|
| Nematodes and these insects: Cotton aphid Cotton fleahopper ¹ Plant bugs ¹ Tobacco thrips Western flower thrips Wireworm | Apply Avicta Duo COT202 to cottonseed at a rate to achieve: 0.15 mg abamectin / 0.375 mg thiamethoxam per seed | To control early season nematodes and these insect pests apply the amount of Avicta Duo COT202 specified in the center column. NOTE: • *1 fl oz of Avicta Duo COT202 contains 3.93 grams of abamectin and 9.84 grams of thiamethoxam. • The typical density of Avicta Duo COT202 is 9.93 lb per gallon. |
| Disease Protection – Early Season | Apply 3.1 – 3.95 fl oz/100 lb seed (0.025 – 0.032 mg ai/seed) of Dynasty CST. | To add early season disease control in cotton, cottonseed may be treated with a mixture of Avicta Duo COT202 plus Dynasty CST. Follow the most restrictive requirements and precautions for all products used in mixtures. |

¹Avicta Duo COT202 only provides suppression of these pests. Suppression means either erratic control ranging from good to poor or consistent control at a level below that which is generally considered acceptable for commercial control.

CLEANING SEED TREATMENT EQUIPMENT

Do not use compressed air hoses to remove the dust from the seed treating equipment or the area around the equipment; use a central vacuum system or a HEPA filter portable vacuum cleaner.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in a cool, dry place.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling

Nonrefillable container. Do not reuse or refill this container. Syngenta's authorized agent will pick up the empty container and handle its disposal. Contact Syngenta or its authorized agent to make arrangements for this service and store empty container in a secure area until pick up.

Triple rinse container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then contact Syngenta or its authorized agent for disposal.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean-up immediately. Take special care to avoid contamination of equipment and facilities during clean-up procedures and disposal of wastes.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

Avicta®, Dynasty®, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company

Viton™ is a trademark of The Chemours Company FC, LLC

Protected by U.S. Patent Nos. 5,852,012, & 6,875,727.

©2026 Syngenta

For non-emergency (e.g., current product information) call
Syngenta Crop Protection at 1-866-796-4368.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

**SCP 1538A-L1C 1025
4247976**

RESTRICTED USE PESTICIDE DUE TO ACUTE INHALATION TOXICITY

FOR SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

| | | | |
|--------------|-------|----|-------------|
| THIAMETHOXAM | GROUP | 4A | INSECTICIDE |
| ABAMECTIN | GROUP | 6 | INSECTICIDE |



Nematicide/ Insecticide

A Seed Treatment Product to Protect
Cotton Seedlings from Early Season
Nematode and Insect Damage

For use only in Syngenta-certified cottonseed treatment facilities and only in seed treatment equipment with closed transfer and application systems. Not for use in hopper box, planter box, slurry box, or other farmer-applied applications.

Active Ingredients:
Abamectin (CAS No. 71751-41-2) 11.2%
Thiamethoxam (CAS No. 153719-23-4) 28.0%

Other Ingredients: 60.8%

Total: 100.0%

Avicta Duo COT202 is a flowable concentrate containing 1.11 pounds of abamectin and 2.78 pounds of thiamethoxam per gallon.

See additional precautionary statements and directions for use in booklet.

EPA Reg. No. 100-1538

EPA Est. 100-NE-001

55887

Product ID

Avicta® and the Syngenta Logo are Trademarks of a Syngenta Group Company

Protected by U.S. Patent Nos. 5,852,012, & 6,875,727.

©2026 Syngenta

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1538A-L1C 1025
4247976

15 gallons
Net Contents

KEEP OUT OF REACH OF CHILDREN.

WARNING/ AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING/AVISO

May be fatal if inhaled. Do not breathe vapor. May be fatal if swallowed. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

This product must be used only in seed treatment equipment with closed transfer application systems.

FIRST AID

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. If swallowed: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

SYNGENTA HOTLINE NUMBER: For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372

Environmental Hazards: This pesticide is toxic to fish and wildlife and highly toxic to aquatic invertebrates. Treated seeds exposed on the soil surface may be hazardous to wildlife. Cover or collect seeds spilled during loading. Do not contaminate water when disposing of equipment washwater or rinsate.

Pollinator Precautions: Abamectin and thiamethoxam are highly toxic to bees exposed to direct treatment or residues in blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Groundwater Advisory: This product has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling

Nonrefillable container. Do not reuse or refill this container. Syngenta's authorized agent will pick up the empty container and handle its disposal. Contact Syngenta or its authorized agent to make arrangements for this service and store empty container in a secure area until pick up.

Triple rinse container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then contact Syngenta or its authorized agent for disposal.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean-up immediately. Take special care to avoid contamination of equipment and facilities during clean-up procedures and disposal of wastes.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!