

# FIFRA Section 24(c) Special Local Need (SLN) Label

CLOPYRALID	GROUP	4	HERBICIDE
------------	-------	---	-----------

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Stinger®

EPA Reg. No. 62719-73

EPA 24(c) Special Local Need Registration SLN No. FL-110010  
For Distribution and Use Only Within the State of Florida

This label valid until December 31, 2027, or until otherwise amended, withdrawn, canceled, or suspended.

### Postemergence Broadleaf Weed Control in Annual Strawberry

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Stinger® herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- In addition to restrictions and limitations imposed by this supplemental labeling, use of Stinger is subject to all use precautions and limitations imposed by the label affixed to the container for Stinger.

#### Directions for Use

Active Ingredient:	% by Weight
clopyralid: 3,6-dichloro-2- pyridinecarboxylic acid, monoethanolamine salt.....	40.9%
Other Ingredients .....	59.1%
Total .....	100.0%

Acid Equivalent: clopyralid: 3,6-dichloro-2-pyridinecarboxylic acid - 31% (3 lb/gal)

Refer to product label for Additional Precautionary Statements, Use Precautions and Restrictions, Mixing Instructions and Application Directions.

**Keep Out of Reach of Children**

## CAUTION

#### Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters. Do not contaminate water used for irrigation or domestic purposes.

Target Weeds	Stinger (pint/acre)	Use Restrictions
<b>Control</b> clovers common ragweed curly dock giant ragweed red sorrel vetch <b>Suppression</b> black medic Carolina geranium cutleaf evening primrose smartweed	1/3 (0.125 lb ae clopyralid/acre)	<ul style="list-style-type: none"> <li>• <b>Preharvest Interval:</b> Do not apply within 2 days of harvest.</li> <li>• <b>Do not</b> make more than 2 applications per crop.</li> <li>• <b>Minimum Retreatment Interval:</b> 14 days.</li> <li>• <b>Do not</b> exceed 2/3 pint (0.25 lb ae clopyralid) per acre per year per crop.</li> <li>• <b>Restricted Entry Interval:</b> 48 hours.</li> <li>• <b>Do not</b> use Stinger with a surfactant on strawberries.</li> <li>• <b>Do not</b> tank mix with other herbicides registered for use on strawberry.</li> <li>• <b>Do not</b> apply by aircraft.</li> <li>• <b>Chemigation:</b> Do not apply Stinger through any type of irrigation system.</li> <li>• <b>Do not</b> apply within 8 hours of expected rainfall or irrigation.</li> </ul>

**Note:** Temporary leaf cupping and/or berry distortion may occur. Therefore, some berries may not be marketable.

**Broadcast Applications:** Apply uniformly with ground equipment in 20 to 60 gallons of water per acre.

**Spot Applications:** Refer to the Hand-Held Sprayers section of the label for Stinger for spot application instructions.

**Precautions:**

- Allow strawberries to become well-established prior to application to new plantings.

**Sprayer Clean-Out:** To avoid injury to desirable plants, equipment used to apply Stinger must be thoroughly cleaned before re-using with any other chemicals. Read the product label for specific clean-out instructions.

**Crop Rotation Interval**

- Read the Crop Rotation Interval section of the section 3 label before rotating to any crop. In many situations, it will not be possible to plant another crop in the same field due to carry over injury to the rotated crop. Consult with your local county agent before planting subsequent crops.
- A field bioassay must be performed prior to planting any crops that are not listed in the Crop Rotation Interval section of the product label. An 18-month rotation is recommended for those crops due to the potential of crop injury.
- A field bioassay must be performed prior to planting any sensitive crops.

**Special Conditions and Risks of Use**

**USE OF Stinger® herbicide (THE “PRODUCT”) ON STRAWBERRY (THE “CROP”) MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS. READ AND UNDERSTAND THESE SPECIAL CONDITIONS AND RISKS OF USE BEFORE USING THE PRODUCT ON THE CROP.**

Corteva Agriscience makes the Product available for use in the manner described in this Special Local Need (SLN) Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury or loss. The decision to use this Product in the manner described in this SLN Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls, and (iv) any other relevant factors.

**By purchasing the Product for use, or using the Product, in the manner described in this SLN Labeling, you acknowledge and accept that:**

- (1) You assume all risk of Crop injury, Crop yield reduction and/or Crop loss;**
- (2) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this SLN Labeling;**
- 3) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaims all warranties, express or implied, including any implied warranties of fitness for a particular purpose or merchantability;**
- (4) Corteva Agriscience disclaims all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to inability to sell the Crop, Crop injury, Crop yield reduction and/or Crop loss;**
- (5) These Special Conditions and Risks of Use supersede any contrary representations or recommendations by Corteva Agriscience, its agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.**

**If these Special Conditions and Risks of Use are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container. These Special Conditions and Risks of Use are required by Corteva Agriscience and not specified by the U.S. EPA or the State of Florida.**

<sup>™</sup><sup>®</sup> Trademarks of Corteva Agriscience and its affiliated companies

R043-144

Accepted: 02/27/23

Replaces R043-040