

## SUPPLEMENTAL LABELING

TOLFENPYRAD	GROUP	21A	INSECTICIDE
	GROUP	39	FUNGICIDE



**EPA Reg. No. 71711-36**

**Berry, Low Growing Group (Crop Subgroup 13-07G) excluding cranberry and blueberry, low bush bearberry; bilberry; cloudberry; lingonberry; muntries; partridgeberry; strawberry; cultivars, varieties, and/or hybrids of these**

This supplemental labeling expires 03/31/2022 and must not be used or distributed after this date.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling and the EPA approved container label must be in the possession of the user at the time of application. Read the label affixed to the container for BEXAR® Insecticide before applying. Use of BEXAR Insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for BEXAR Insecticide. New use directions appear on this supplemental labeling that may be different from those that appear on the container label.

### APPLICATION RATE CHART FOR BEXAR INSECTICIDE

Pest	Rate/Acre	Use Directions
Aphids Eriophyid mites Katydid Leafrollers Lygus Powdery mildew Thrips  Spotted wing drosophila*	27.0 fl oz/acre (0.28 lb ai/acre)	<b>USE RESTRICTIONS</b> <ul style="list-style-type: none"> <li>• <b>Do not apply by air on low growing berry.</b></li> <li>• Apply by ground <b>only</b>, using a minimum of 20 gallons of water per acre.</li> <li>• Do not apply more than 81.0 fl oz (0.85 lb ai) per acre per year.</li> <li>• Do not make more than 3 applications per year.</li> <li>• Allow at least 7 days between applications.</li> <li>• Preharvest Interval (PHI): 1 day</li> </ul> <b>USE RECOMMENDATIONS - DISEASES</b> <ul style="list-style-type: none"> <li>• Begin applications prior to onset of disease.</li> <li>• Use of an agriculturally approved nonionic surfactant at 0.25% v/v may improve disease control.</li> <li>• If weather conditions remain conducive to disease development, apply another registered fungicide product with a different mode of action 7 to 10 days later.</li> <li>• Consult local extension recommendations or your agricultural consultant for information specific to your area.</li> </ul>

\* suppression only

©2019 Nichino America, Inc. Bexar and Nichino America logo are registered trademarks of Nichino America, Inc.

**NICHINO  
AMERICA®**

Nichino America, Inc.  
4550 Linden Hill Road, Suite 501  
Wilmington, DE 19808  
888-740-7700