

**FIFRA SECTION 24(C) SPECIAL LOCAL NEED REGISTRATION
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON**



 FOR ORGANIC PRODUCTION

EPA Reg. No. 10163-330 EPA SLN No. WA-180008

DANGER/PELIGRO

**This label for Previsto expires and must not be distributed or used in accordance with
this SLN registration after December 31, 2023.**

DIRECTIONS FOR USE

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

CROP	PEST	RATE	COMMENTS
Apples and Pears (Bloom, growing season)	Fire Blight	3 – 4 Quarts per Acre	Begin applications at delayed dormancy and repeat as necessary. Use of a reliable fire blight disease model is highly recommended. Use higher rates for more susceptible varieties. If injury is observed, discontinue use immediately. Use on D'Anjou and Comice pears is not recommended due to a high potential for unacceptable phytotoxicity to fruit and foliage.
	<ul style="list-style-type: none"> • Minimum retreatment interval = 2 days. • For bloom/growing season application, do not apply more than 4 quarts of formulated product per acre per application. • For all application timings, do not apply more than 212 quarts of formulated product per acre per year. • Spray volume should not exceed 100 GPA. • Cease spray activities and allow spray to dry before daytime temperatures exceed 85 degrees. • Do not apply when conditions favor slow drying or immediately after or during frost control or crop irrigation, unacceptable markings to fruit and/or foliage may occur. • Do not tank mix with other pesticides, foliar nutrients, adjuvants, spreaders, buffering agents or stickers. 		

Previsto® is a registered trademark used under license by Gowan Company, L.L.C.

RESTRICTIONS/PRECAUTIONS:

- Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.
- This pesticide is toxic to fish and aquatic invertebrates and may contaminate water through runoff. Previsto should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may refer to the WSDA Natural Resource Assessment Section web site (agr.wa.gov/PestFert/NatResources/EndangSpecies.aspx) or contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information regarding aquatic species listed as threatened or endangered. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.
- For use in accordance with this SLN label do not apply this product through any type of irrigation system.

WSDA CONTAINER DISPOSAL GUIDANCE: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA Waste Pesticide Program web site (agr.wa.gov/PestFert/Pesticides/WastePesticide.aspx). Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

Section 24(c) Registrant:
Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569

Rev 11/28/18