



Bayer CropScience LP
800 N. Lindbergh Blvd
St. Louis, Missouri 63167

1-866-99BAYER (1-866-992-2937)

PROPULSE®

For suppression of Nematodes at a reduced
in-furrow application rate in Corn

EPA Reg. No. 264-1084

FIFRA 2(ee) Recommendation

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF ALABAMA,
ARKANSAS, COLORADO, DELAWARE, FLORIDA, GEORGIA, IDAHO, KANSAS,
MARYLAND, MISSISSIPPI, NEBRASKA, NEW JERSEY, NORTH CAROLINA,
NORTH DAKOTA, OKLAHOMA, OREGON, PENNSYLVANIA, SOUTH CAROLINA,
TEXAS, VIRGINIA, WASHINGTON, AND WISCONSIN**

This recommendation is valid until December 31, 2022

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

Tank mixes with Propulse and some fertilizers and micronutrients has been problematic. It is highly recommended that users conduct a “jar-test” to make sure that tank mix partners are compatible before mixing. Contact your local Bayer rep for more information.

CORN

Sweet Corn; Field Corn; Field Corn Grown For Seed, and Popcorn

Disease Suppression	Application Rate	Application Instructions
Nematodes	8.0 fl oz/acre	In-furrow spray during planting directed on or below seed. Tank mixes with Propulse and some fertilizers and micronutrients has been problematic. It is highly recommended that users conduct a "jar-test" to make sure that tank mix partners are compatible before mixing. Contact your local Bayer rep for more information.

Restrictions:

- Do not make more than 2 applications or apply more than 34.2 fl oz of PROPULSE/acre per year.
- Regardless of formulation or method of application, do not apply more than 0.446 lbs fluopyram or 0.712 lbs prothioconazole per acre per year, including soil and foliar uses.
- Apply by either ground, aerial, or chemigation application equipment.
- Pre-Harvest Interval (PHI): 0 day - sweet corn ears and forage and field corn grown for seed; 14 days – field corn, popcorn, (grain and fodder).
- To limit the potential for development of disease resistance to this fungicide, do not make more than 2 sequential applications of PROPULSE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*