

RESTRICTED USE PESTICIDE

Due to Acute Toxicity And Toxicity to Birds and Mammals

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

OXAMYL

GROUP

1A

INSECTICIDE/NEMATOCIDE

FIFRA Section 24(c) Special Local Need Label



Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Vydate L

EPA Reg. No. 352-372

EPA SLN No. NY-990002

For Distribution and Use Only in the State of New York

FOR USE ON GARLIC AND BULB ONIONS IN THE STATE OF NEW YORK

Expiration date: This label expires and must not be distributed or used in accordance with this SLN registration after May 31, 2026.

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read this SLN and the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this FIFRA 24(c) Special Local Need labeling is subject to all use precautions and limitations imposed by the label affixed to the container for this product.

Directions for Use

Vydate® L is recommended for use on garlic and bulb onions to control root knot (except Javanese), stubby root, stem and bulb nematodes in New York. Vydate L Insecticide/Nematicide is recommended for use on garlic and bulb onions to control onion thrips in the State of New York.

Do not use in Suffolk and Nassau Counties, Long Island, New York.

Application

Control of Root Knot (except Javanese), Stubby Root, Stem and Bulb Nematodes

Broadcast Treatment

Apply 2 gals. of Vydate L per acre in a minimum of 20 gals. of water per acre as a broadcast treatment to the soil using ground application equipment. Apply within one week of planting. Thoroughly incorporate to a depth of 4 to 6 inches in the soil.

In-Furrow Treatment

Apply 3/4 to 1 gal. of Vydate L per acre at planting as an in-furrow drench using 100 to 150 gals. of water per acre, or 1 1/2 to 2 gals. per acre as an in-furrow band spray using 20 to 50 gals. of water per acre. Application to soil must be incorporated by water or by mechanical means.

Foliar Treatment for Onion Thrips

Apply by ground or air at 2 to 4 pts/acre (minimum 5 gal/acre by air).

Apply when insects appear in significant numbers. Repeat applications as necessary at 5 to 7 day intervals to maintain control.

Add a wetting agent to improve coverage.

NOTE:

- Do not harvest tops of treated bulb onions.
- Do not apply within 14 days of harvest.
- Do not use on green onions.
- Do not apply more than 18 pts (2 1/4 gals) Vydate L per acre per season.

When making an application to garlic and bulb onions, do not apply this product through any type of irrigation system.

Soil applications must be incorporated into soil by water or mechanical means.

Do not apply more than 8 applications per acre per growing season.

IMPORTANT

BEFORE USING VYDATE L, READ AND CAREFULLY OBSERVE THE CAUTIONARY STATEMENTS AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

This bulletin contains new or supplemental instructions for use of this product which may not appear on package label. Follow the instructions carefully.

This label must be in the possession of the user at the time of pesticide application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

™®Trademarks of Corteva Agriscience and its affiliated companies

R641-010

Approved: 05/09/2023

Replaces R-537