



Bayer CropScience LP  
800 N. Lindbergh Blvd.  
St. Louis, MO 63167  
1-866-99BAYER (1-866-992-2937)

## PROPULSE®

Reduced rate of PROPULSE for diseases of  
Highbush and Lowbush Blueberry

EPA Reg. No. 264-1084

### ***FIFRA 2(ee) Recommendation***

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF ALABAMA, CONNECTICUT, DELAWARE, FLORIDA, GEORGIA, MASSACHUSETTS, MARYLAND, MAINE, MICHIGAN, MISSISSIPPI, NORTH CAROLINA, NEW HAMPSHIRE, NEW JERSEY, NEW YORK, OREGON, PENNSYLVANIA, RHODE ISLAND, SOUTH CAROLINA, VIRGINIA, VERMONT AND WASHINGTON.**

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

#### **DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Blueberry, Highbush; Blueberry, Lowbush;

Disease Control	Application Rate	Application Instructions
Septoria leaf spot ( <i>Septoria</i> spp.)  Monilinia blight ( <i>Monilinia vaccinii-corymbosi</i> )  Valdensinia leaf spot ( <i>Valdensinia heterodoxa</i> )  Leaf rust ( <i>Thekopsora minima</i> )  Anthracnose ( <i>Colletotrichum gloeosporioides</i> )  Botrytis blight ( <i>Botrytis cinerea</i> )  Alternaria fruit rot ( <i>Alternaria</i> spp.)  White pine blister rust ( <i>Cronartium ribicola</i> )	10.0 to 13.6 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 10-day interval. When disease pressure is severe, use the shorter intervals.

**Restrictions:**

- Do not make more than 2 applications or apply more than 27.3 fl oz of PROPULSE/acre per year.
- Regardless of formulation or method of application, do not apply more than 0.446 lbs fluopyram or 0.356 lbs prothioconazole per acre per year.
- Apply by either ground or chemigation application equipment.
- Pre-Harvest Interval (PHI): 7 days
- To limit the potential for development of disease resistance to this fungicide, do not make more than 2 sequential applications of PROPULSE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.  
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

*As with any crop-protection product, always read and follow label directions.  
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*