

RESTRICTED USE PESTICIDE

Due to eye and skin effects. For retail sale to and use by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

FIFRA SECTION 24(c) SPECIAL LOCAL NEED LABEL (SLN) FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON

Comite®

AGRICULTURAL MITICIDE

EPA Reg. No. 400-104

EPA SLN No. OR-200010

REDUCED APPLICATION TIME INTERVAL FOR POTATOES

This label is valid until December 31, 2025 or until otherwise amended, withdrawn, canceled, or suspended.

ACTIVE INGREDIENT:	% BY WT.
Propargite (Sulfurous acid, 2-[4-(1,1-dimethyl-ethyl) phenoxy] cyclohexyl-2- propynyl ester)*	73.6%
OTHER INGREDIENTS:.....	26.4%
TOTAL:.....	100.0%

*Contains 6.55 lbs. active ingredient per gallon.

*Contains petroleum distillates.

KEEP OUT OF REACH OF CHILDREN DANGER / PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may cause reproductive effects in wildlife. Do not apply directly to water or areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This Special Local Need (SLN) label and the federal label for this product must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, precautions, restrictions, and Worker Protection Standard requirements on the main product label for **COMITE®**, EPA Reg. No. 400-104, and on this SLN label.
- Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State Tribe, consult the agency responsible for pesticide regulation.

GENERAL INSTRUCTIONS

COMITE is a liquid emulsifiable concentrate for preparing sprays to control mites.

COMITE is not systemic in action, therefore, complete coverage of both upper and lower leaf surfaces is necessary for effective control.

MIXING INSTRUCTIONS

After three-quarters full of water, add recommended dosage (from table) of **COMITE** to the spray tank. Fill tank, agitate and spray thoroughly to cover foliage for best results.

CROPS	MITES CONTROLLED	APPLIC. RATE (fl. oz. per acre)	APPLICATION INFORMATION
POTATOES (Oregon)	Two-spotted spider	32 - 40	Apply high rate to areas with high mite infestation and to areas which historically have had high mite infestations. See COMPATIBILITIES section of main product label.
	Ground Application: Apply in 20 - 50 gallons total volume per acre. Aerial Application: Apply in a minimum of 10 gallons total volume per acre. Chemigation Application: Apply in 0.1-0.2 acre inches of water. (See CHEMIGATION PRECAUTIONS FOR CORN AND POTATOES section of main product label)		
	POTATO RESTRICTIONS: Apply no more than 80 fl. oz. COMITE (4.1 lbs. propargite a.i.) per acre per year. Do not make more than 2 applications per season. Minimum Spray Interval: 14 days between applications. Restricted Entry Interval (REI): 2 days. See AGRICULTURAL USE REQUIREMENTS section of main product label for general REI exemptions. There is no exception allowed for hand weeding or thinning. Pre-harvest Interval: Do not harvest within 14 days of the last application.		

RESTRICTIONS/PRECAUTIONS

WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS:

- Do not apply by ground within 50 feet or by air within 75 feet of lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries, or commercial fish farm ponds.
- The above excludes irrigation canals and wasteways as well as man-made irrigation conveyance structures and impoundments, unless an exclusion contains water year-round.
- Risk of exposure to sensitive aquatic areas can be reduced by making applications when the wind direction is away from the aquatic area.

RUNOFF PRECAUTIONS:

- Under some conditions, propargite may have a high potential for runoff into surface water for several days after application.
- Do not apply in the following areas:
 - Frequently flooded areas (excluding crop irrigation areas).
 - Areas where intense or sustained rainfall is forecasted to occur within 48 hours.
- Care should be taken when making applications where the following conditions exist:
 - Poorly draining or wet soils with readily visible slopes toward adjacent surface water.
 - Areas with in-field canals or ditches that drain to surface water.
 - Areas not separated from adjacent surface waters with vegetated filter.
- Consult your soil conservation service for recommendations in your use area.

ENDANGERED/THREATENED SPECIES:

- You may contact the Oregon Department of Fish & Wildlife, National Marine Fisheries Service or US Fish & Wildlife Service for information on listed threatened or endangered species.

THIS LABEL IS IN ADDITION TO THE STANDARD FEDERAL LABEL PRINTED ON THE CONTAINER

Follow all applicable directions, restrictions, and precautions on the EPA registered label.

This label must be in the possession of the user at the time of the pesticide application.

SLN 24(c) Registrant:
MacDermid Agricultural Solutions, Inc.
c/o ARYSTA LIFESCIENCE NORTH AMERICA, LLC
15401 Weston Parkway, Suite 150
Cary, NC 27513
For Product Use Information Call 1-866-761-9397
120414V019ORSLN

Arysta LifeScience and the Arysta LifeScience logo are registered trademarks of Arysta LifeScience Corporation.
COMITE is a registered trademark of an Arysta LifeScience Group Company.
©2018 Arysta LifeScience Group Company.