

Product Bulletin



Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

WideARmatch™ with Arylex™ active

EPA Reg. No. 62719-746

Special 2(ee) Recommendation†

(For Distribution and Use Only in the States of Idaho, Montana, Oregon and Washington)

For Improved Control of Narrowleaf Hawksbeard, Canada Thistle, Mayweed Chamomile (Dogfennel), False (Scentless) Chamomile, Cornflower, Meadow Salsify (Goatsbeard), Annual Sowthistle Species, and Volunteer Alfalfa

ATTENTION

†This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of this product according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for this product.

Directions for Use

Refer to WideARmatch™ with Arylex™ active product label for General Use Precautions, Mixing Directions and Application Directions.

Apply at 14 to 19.6 fl oz of WideARmatch with Arylex active per acre to actively growing wheat (including spring, winter and durum), barley, and triticale from the 2-leaf crop growth stage up to flag leaf emergence (Zadoks scale 39). Use the high rate for improved control of narrowleaf hawksbeard, Canada thistle, mayweed chamomile (dogfennel), false (scentless) chamomile, cornflower, meadow salsify (goatsbeard), annual sowthistle species, and volunteer alfalfa. Apply to susceptible broadleaf weed seedlings that are actively growing and less than 8 inches tall. For best results apply when susceptible broadleaf weed seedlings are actively growing and less than 4 inches tall. Only weeds emerged at the time of treatment will be controlled.

Warm, moist growing conditions promote active weed growth and enhance the activity of WideARmatch with Arylex active by allowing maximum foliar uptake and contact activity. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and re-growth may occur. For best results, ensure thorough spray coverage of target weeds.

™® Trademarks of Corteva Agriscience and its affiliated companies

R454-003

Issued: 09/09/22

Replaces: R454-001 and R454-002