

Stinger® herbicide (clopyralid) is a state restricted use pesticide and is to be distributed only by licensed pesticide dealers.

CLOPYRALID | GROUP 4 | HERBICIDE

FIFRA Section 24(c) Special Local Need (SLN) Label

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Stinger®

EPA Reg. No. 62719-73

EPA SLN No. WA-230007

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

This label for Stinger expires and must not be distributed or used in accordance with this SLN registration after December 31, 2027.

Postemergence Broadleaf Weed Control in Strawberry

Use of Stinger® herbicide on strawberry may result in crop injury, crop yield reduction, and/or crop loss. Read and understand the Special Conditions and Risks of Use section on this label before using this product in accordance with this FIFRA Section 24(c) Special Local Need.

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and the federal label for this product must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the EPA registered label.

Directions for Use

Stinger is a residual herbicide and applications must be made based on accurate rate per acre calibrations. Refer to the product label for General Use Precautions, Restrictions, Mixing and Application instructions, and a complete list of weeds controlled or suppressed.

Target Broadleaf Weeds, Application Rates, and Use Restrictions:

Target Broadleaf Weeds	Application Rate (pint/acre)	Use Restrictions
clover, white dandelion groundsel, common hawksbeard, smooth pineappleweed sowthistle, annual (suppression) thistle, bull (suppression) thistle, Canada (suppression) vetch	Post-harvest application: 1/3 to 2/3 pint/acre (0.125 – 0.25 lb ai clopyralid/acre) Spring application: 1/3 pint/acre maximum (0.125 lb ai clopyralid/acre)	<ul style="list-style-type: none"> • Make 1 to 2 applications per year, not to exceed a total of 2/3 pint per acre (10.67 fl oz per acre; 0.25 lb ai per acre) per year. Make only 1 application in the spring. • Do not use surfactants when applying Stinger to strawberry. • Do not tank mix Stinger with other herbicides registered for use on strawberry. • Do not apply Stinger within 6 to 8 hours of expected rainfall or irrigation. • Do not apply within 4 months of planting. • Post-Harvest Application: Apply between September 15 and November 15. • Spring Application: Apply prior to first bloom. Do not apply within 30 days of harvest.

Broadcast Application:

For difficult to control weeds, best results will be achieved with a post-harvest application followed by a spring application. Apply the spray uniformly in 30 to 75 gallons of water per acre using ground equipment only.

Post-Harvest Application: After harvest, make a ground application between September 15 and November 15 at a rate of 1/3 to 2/3 pint of Stinger (0.125 to 0.25 lb ai clopyralid) per acre. For control of Canada thistle, apply Stinger after the majority of basal leaves have emerged but prior to bud stage.

Spring Application: Prior to first bloom, make a ground application in the spring at a maximum rate of 1/3 pint of Stinger (0.125 lb ai clopyralid) per acre. Do not apply within 30 days of harvest.

Spot Application: Use a rate per acre equivalent to a broadcast application and follow the same use restrictions. Refer to the Hand-Held Sprayers section on the product label for Stinger for spot application instructions. Apply the spray uniformly.

Specific Use Restrictions:

- **Chemigation:** Do not apply Stinger through any type of irrigation system.
- Do not Stinger apply by aircraft.
- Refer to the product label for Stinger for crop rotation intervals.
- This product is persistent and may be present in treated plant materials for months to years after application. Do not sell or transport treated plant materials off-site for compost distribution for 18 months after application.
- Do not apply Stinger to strawberry if you are unwilling to accept minor leaf injury. Minor leaf cupping may occur.

WSDA Container Disposal Guidance: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA website under the Waste Pesticide Program at <https://agr.wa.gov/wastepesticide>. Cleaned containers may also be disposed of in a sanitary landfill if permitted by the county. Burning is not a legal method of container disposal in Washington.

Special Conditions and Risks of Use

USE OF Stinger (THE “PRODUCT”) ON STRAWBERRY (THE “CROP”) MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS. READ AND UNDERSTAND THESE SPECIAL CONDITIONS AND RISKS OF USE BEFORE USING THE PRODUCT ON THE CROP.

Corteva Agriscience makes the Product available for use in the manner described in this Special Local Need Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury or loss. The decision to use this Product in the manner described in this Special Local Need Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls, and (iv) any other relevant factors.

By purchasing the Product for use, or using the Product, in the manner described in this Special Local Need Labeling, you acknowledge and accept that:

- (1) You assume all risk of Crop injury, Crop yield reduction and Crop loss;
- (2) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this Special Local Need Labeling;
- (3) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaims all warranties, express or implied, including any implied warranties of fitness for a particular purpose or merchantability;
- (4) Corteva Agriscience disclaims all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to Crop injury, Crop yield reduction and/or Crop loss;
- (5) These Special Conditions and Risks of Use supersede any contrary representations or recommendations by Corteva Agriscience, its agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

If these Special Conditions and Risks of Use are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container. These Special Conditions and Risks of Use are required by Corteva Agrisciences and not required or enforced by the U.S. EPA or the WSDA.

24(c) Registrant: Corteva Agriscience LLC
9330 Zionsville Road
Indianapolis, IN 46268

TM® Trademarks of Corteva Agriscience and its affiliated companies

R043-188
Accepted: 04/26/23
Replaces R043-113