

**RESTRICTED USE PESTICIDE  
DUE TO TOXICITY TO NONTARGET ORGANISMS**

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certificate.

## Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

# rozol<sup>®</sup> VOLE BAIT

**FOR CONTROL OF VOLES AND DEER MICE IN HOP FIELDS (POST-HARVEST OR DORMANT-SEASON USE ONLY).**

**EPA Reg. No. 7173-242 EPA SLN No. WA-180006**

This label for Rozol Vole Bait expires and must not be distributed or used in accordance with this SLN registration after December 31, 2024.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions and precautions on the EPA registered label.

**GENERAL INFORMATION:** Experts contacted by WSDA indicate that certain varieties of hops (e.g., Cascade, Centennial, Chinook, CTZ, Simcoe) appear to be preferred by voles. However, voles will also feed on other varieties of hops. Preferably, make application when the grass has not yet matted. Bait is more readily accepted when voles are protected by vegetation.

**DIRECTIONS FOR USE:** This product may only be used to control voles (*Microtus spp.*) and deer mice (*Peromyscus spp.*) in hop fields (post-harvest or dormant-season use only).

**APPLICATION DIRECTIONS:** Before application, locate vole trails, runway systems and harborage areas to be treated.

**Hand spot baiting:** Place 1-1/2 ounces (6 tablespoons) of bait in each active hole, trail or runway. Cover each placement with grass or shingle to avoid exposing nontarget organisms, or place bait in a tamper resistant bait station.

**Ground or Aerial broadcast baiting:** Broadcast evenly at 10 pounds of bait per acre using commercial bait dispensing equipment. Concentrate baiting in areas with heaviest vegetative cover. Do not broadcast bait on bare ground. Do not apply bait in piles or permit piles to be formed by equipment. If populations are high, make a second application 1 to 2 months after the first.

**RESTRICTIONS/PRECAUTIONS:**

- Do not apply more than 10 pounds of bait per acre per application.
- Do not apply more than 20 pounds of bait per acre per season.
- Do not apply bait during the growing season.
- Only apply bait after fall harvest, before spring growth and when no rain or snow is expected for three consecutive days.
- Do not apply within 50 feet of any body of surface water or where raptors are actively feeding on voles.
- Do not apply bait on bare ground or to crops or sites not specified above.
- Do not allow pets, livestock, or other non-target animals to be exposed to bait. Do not allow animals to graze in treated areas. Do not use hay cut after application for feed or bedding. To avoid exposing nontarget organisms, follow the instructions in the "Pesticide Disposal" section of the federal label to ensure proper clean-up of any bait for reuse or disposal.
- Do not apply bait in a manner that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Keep all other persons out of the treated area during application.
- This product is extremely toxic to fish and wildlife. Dogs and predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten the bait. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.
- Rozol Vole Bait should not be used under this SLN label where impact on threatened or endangered species is likely. You may refer to the WSDA Natural Resources Assessment Section web site (<https://agr.wa.gov/departments/land-and-water/natural-resources/endangered-species-program>) for additional information related to pesticide use and endangered species protection. Information from the EPA about no-spray buffer zones is available at <https://agr.wa.gov/departments/land-and-water/natural-resources/endangered-species-program/buffers>. Pesticide applicators may use the Washington Department of Fish and Wildlife's SalmonScape mapping tool at <https://apps.wdfw.wa.gov/salmonscape> to determine if listed salmonid species occur in or near a proposed application site. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

**WSDA Container Disposal Guidance:**

Completely empty the contents of the container into application equipment, and apply to a labeled crop or site. The empty container may be disposed of in a sanitary landfill. Burning is not a legal method of container disposal in Washington.

Section 24(c) SLN registrant:  
Liphatech, Inc.  
3600 W. Elm Street  
Milwaukee, WI 53209

**LIPHATECH<sup>®</sup>**

Rev Date 01/16/2020