

BROAD SPECTRUM FOLIAR INSECTICIDE, MITICIDE AND FUNGICIDE FOR AGRICULTURAL, ORNAMENTAL AND FOREST CROPS.

ACTIVE INGREDIENTS:

Cinnamon Oil 60.0 %

OTHER INGREDIENTS: 40.0 %

Potassium oleate

TOTAL 100.0 %

CINNERATE is an innovative formulation with high essential oils content, providing a knock-down and eradication effect towards several species of mites, insects and fungus.

CINNERATE disrupts the functioning of cell membranes as well as inhibits the growth and promotes dehydration.

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information, call the National Pesticides Information Center at 1-800-858-7378, 7:30 AM to 3:30 PM, Pacific time (PT), Mon. - Fri. During other times, call the Poison Control Center at 1-800-222-1222.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Caution. Causes moderate eye irritation. Prolonged or frequent contact may cause allergic reactions in some individuals. Avoid contact with eyes, skin and clothing. Wear protective eyewear and gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and handlers must wear:

- Long-sleeved shirt and long pants
- Socks and shoes
- Protective eyewear
- Chemical resistant gloves made of neoprene, nitrile or natural rubber

User Safety Recommendations

Users should remove clothing/PPE immediately if the product gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of the gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

The application of CINNERATE does not modify odor or taste of commodities. Crops can be consumed without a security period of time after application.

CINNERATE



Net content:

2.5 Gallons

Batch:

Packaging date:

seipasa®

DIRECTIONS FOR USE

CINNERATE is a contact miticide, insecticide and fungicide, therefore direct contact with the target organism is required for control.

CINNERATE must be applied using a sufficient volume of water to insure complete crop coverage. Not for aquatic use.

APPLICATION

For best results, spray coverage should be uniform and complete. Spray droplet size should be adjusted to maximize coverage, minimize drift, and allow direct contact with as many pathogens (insects, mites, fungi) as possible. **Use lower rates for light to moderate pest pressure and higher rates for heavy infestations.**

CINNERATE can be applied to crops at any time up to and including the day of harvest. **Begin spraying when pest first appear; do not wait until plants are heavily infested. Repeat as needed (5-day minimum interval) to maintain desired level of control.** There are no restrictions on the number of applications per season.

Always follow these directions: Apply in early morning or late afternoon / evening. Spray directly onto pest(s) or pathogen(s). Apply before noticeable foliar pest damage occurs. Thorough spray coverage is essential for good pest control.

MIXING INSTRUCTIONS

Shake well before using. The use of adjuvants (spreaders and/or penetrants) is highly recommended for improving performance. Fill spray tank ½ full with intended amount of water (>45°F). Add CINNERATE at recommended rate, then other products as desired, and continue filling with water. In areas with hard water (>200 ppm), addition of a water softener is recommended before adding this product to the spray tank. Maintain agitation while mixing and during application for best results. Always clean tank, pump and lines thoroughly with water after use. **DO NOT MIX CINNERATE WITH Captan, Bordeaux mixtures, oxidizing agents such as bleach or highly alkaline or acid products.**

Mix CINNERATE with only those chemicals that are registered for use on specific crop to be treated.

GROUND SPRAY

Spray-to-wet applications of this product may be made with standard field sprayers and backpack or other hand-held spray equipment.

AERIAL SPRAY

Prior to application, check the aircraft/helicopter spray equipment for uniformity of spray pattern, spray width and output. Apply a minimum of 10 gallons per acre of diluted product. Optimum results are obtained when plant foliage is not too dense. **COVERAGE IS ESSENTIAL FOR GOOD CONTROL.** Use an output that will ensure optimum product penetration and coverage. Make aerial applications when the wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Application more than 10 feet above the crop canopy is not recommended.

DRIP IRRIGATION

For soil inhabiting insects and insect stages, use 2-4 pints per acre of plant bed in 2-3 applications over a 7-14-day schedule. Set emitter spacing and line pressure to give best coverage and reduce leaching.



MANUFACTURED BY:

SEIPASA, S.A.

22240 Tardienta

(Huesca)

Tel. +34 974 253 433

www.seipasa.com

MADE IN SPAIN

PLANT SAFETY AND COMPATIBILITY

Rate recommendations are based on product testing on a wide variety of plants. However, not all plant species/varieties/cultivars have been tested with this product, applied alone or with possible spray mix combinations (adjuvants, fertilizers, or other pesticides). We recommend testing all tank mix combinations for physical compatibility and plant safety. Test spray a few leaves with intended spray mix 1 day before spraying entire plant. Do not use if leaf burn or spotting is observed. Use with care on plants with tender tissue. Do not apply to wilted or otherwise stressed plants, or to newly transplanted material prior to root establishment. Do not apply when the temperature is >90°F. Consult your dealer or distributor for specific recommendations. This product cannot be mixed with any product containing a label prohibition against such mixing. Do not exceed label dosage rate.

TARGETED PESTS

MITES: including but not limited to: **Two spotted spider mite, European red mite, Pear Leaf Blister Mite, Pear Rust Mite, etc.**

INSECTS: including but not limited to **Scales** (such as **San Jose Scale, California Red Scale, Yellow Scale, Black Scale, Soft Scale, etc.**), **Aphids** (such as **cotton aphid, green peach aphid** among others), **Thrips** (such as **Thrips spp., Frankliniella spp., Heliothrips spp., Scirtothrips citri** among others.) **Psyllids** (such as **Bactericera spp., Diaphorina citri, Cacopsylla spp.**) **Leafhoppers** (such as **Empoasca spp., Macrosteles spp., and Graphocephala spp.**) **White flies** (such as **Dialeurodes spp., Trialeurodes spp., Bemisia spp.**) **Flies** (such as **walnut husk fly, fungus gnats**) **Armyworms, Caterpillars and loopers** (such as **cabbage looper, tent caterpillar, fruit worms, alfalfa caterpillar, caterpillar larvae** among others) **Moth** (such as **codling moth, grape berry moth, Other pests** such as **borers** (**Peach twig borer, Hypothenemus hampei**), **mealybugs** (**Planococcus spp., Pseudococcus spp., Dysmicoccus brevipes**) and **apple maggots, lygus bugs, blueberry maggot, leafrollers**).

DISEASES: such as **powdery mildew and rusts. and sporulated forms of fungi** as **Botrytis cinerea, Fulvia fulva, etc.**

APPLICATION RATE TABLE

For the control of the different pests, use CINNERATE as indicated in the following table for each crop. Use the higher rate when disease pressure is high or a severe infection has occurred.

CROPS	RATES (fl.oz. /100 gallons)	GALLONS PER ACRE	
Grown outdoors, in greenhouses, shadehouses and nurseries; including but not limited to			
Vegetables: Asparagus, Bean, Beet, Broccoli, Brussels sprouts, Cabbage, Carrot, Cauliflower, Celery, Cucumber, Eggplant, Endive, Garlic, Leek, Lettuce, Melon, Okra, Onion, Pepper, Potato, Pumpkin, Radicchio, Shallots, Spinach, Squash, Swiss chard, Tomato	16-64	10-20	Low volume applications
		25-100	< 20 inches
		100-160	20-50 inches
		160-210	> 50 inches
Fruits and nut trees: Almonds, Apricots, Apples, Cherries, Citrus, Nectarines, Peaches, Pears, Pecans, Pistachio, Plums, Prunes, Walnuts	16-64	50-500	
		25-45	200-1000
	32-64	100-300	Pear psylla, Pear leaf Blister Mite and Pear Rust Mite
Small fruits and Berries: Blackberries, Blueberries, Bushberries, Caneberries, Cranberries, Currants, Guava, Raspberries, Strawberries	16-64	30-300	
Grapes	16-45	25-300	
	32-64	50-200	Mealybugs
Field crops: Cereal grain, Cotton, Dry beans, Field corn, Grasses grown for seed, Lentils, Sugar beets, Sweet corn, Tobacco, Hops	16-64	10-210	
		>15	Aerial application
Tropical and other fruits: Avocados, Bananas, Dates, Figs, Mangoes, Papaya, Pomegranate, Pineapple, Kiwi, Coffee	16-64	50-200	
Herbs and pasture: Alfalfa, Basil, Chives, Cilantro, Dill, Mint, Oregano, Parsley, Pasture, Thyme, Hemp, Cannabis	16-45	10-100	
Ornamentals: Turf, Lawn, Golf courses, and landscape areas	16-50	50-210	
Forest: Pine trees	16-45	85-125	

Note to User: Unintentional consequences such crop injury may result due to environmental or growing conditions, manner of use or application. Because plant tolerance of pesticides varies as conditions vary, treat a few plants under conditions unexpected to be encountered and observe for plant damage prior to full-scale application to large number of plants.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container in a cool, well-ventilated area away from children, animals, foods, feeds and seeds. Avoid excessive heat and freezing temperatures. Keep container tightly closed during storage.

Product Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Triple rinse (or equivalent) during mixing and loading. Recycling decontaminated containers is the best option of container disposal. The Agricultural Container Recycling Council (ACRC) operates the national recycling program. To contact your state or local ACRC recycler, visit the ACRC web page at www.acrcycle.org. Decontaminated containers may also be disposed of in a sanitary landfill.

WARRANTY AND CONDITIONS OF SALE

SELLER OFFERS AND THE BUYER ACCEPTS THIS PRODUCT UNDER THE FOLLOWING AGREED TERMS AND CONDITIONS OF SALE AND WARRANTY.

The directions for the use of this product are believed to be reliable and should be followed carefully. However, it is impossible to take into account all the variables and to eliminate risks associated with its use. Injury or damage may result because of conditions that are beyond the control of the SELLER.

SELLER warrants that this product conforms to the chemical description on this label and is believed to be reasonably fit for the purposes stated on this label when used in accordance with the directions on this label under conditions of normal use.

SELLER MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

To the fullest extent permitted by law, the SELLER shall not be liable for consequential, special or indirect damages resulting from the use or handling of the product. Any variation or exemption from this warranty must be in writing and signed by authorized representative or the seller.

This product has not been registered by the U.S. Environmental Protection Agency (EPA). SEIPASA SA represents that this product qualifies for 25(b) exemption from registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).