

RESTRICTED USE PESTICIDE / PESTICIDA DE USO RESTRINGIDO

(GROUND AND SURFACE WATER CONCERNS) / (PREOCUPACIONES POR LAS AGUAS SUBTERRÁNEAS Y SUPERFICIALES)

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION. USERS MUST READ AND FOLLOW ALL PRECAUTIONARY STATEMENTS AND INSTRUCTIONS FOR USE IN ORDER TO MINIMIZE POTENTIAL FOR ATRAZINE TO REACH GROUND AND SURFACE WATER. / SOLO PARA VENTA AL POR MENOR Y USO POR APLICADORES CERTIFICADOS O PERSONAS BAJO SU SUPERVISIÓN DIRECTA, Y SOLO PARA LOS USOS PREVISTOS EN LA CERTIFICACIÓN DEL APLICADOR CERTIFICADO. LOS USUARIOS DEBEN LEER Y SEGUIR TODAS LAS INDICACIONES DE PRECAUCIÓN E INSTRUCCIONES DE USO PARA MINIMIZAR LA POSIBILIDAD DE QUE LA ATRAZINA LLEGUE A LAS AGUAS SUBTERRÁNEAS Y SUPERFICIALES.

Atrazine GROUP 5 HERBICIDE

ATRAZINE 90 WDG HERBICIDE

For Season-Long Weed Control in Corn, Sorghum and Certain Other Crops.

ACTIVE INGREDIENTS:

Atrazine (2-chloro-4-ethylamino-6-isopropylamino-s-triazine)	88.2%
Related compounds	1.8%
OTHER INGREDIENTS	10.0%
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN / MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS

CAUTION / PRECAUCIÓN

See Inside This Label Booklet For Additional Precautionary Statements, Complete First Aid, Directions for Use, Storage and Disposal and Other Use Information.

FOR A MEDICAL EMERGENCY OR HELP WITH ANY SPILL, LEAK, FIRE OR EXPOSURE INVOLVING THIS MATERIAL, CALL DAY OR NIGHT CHEMTREC 1-800-424-9300.

EPA REG. NO. 34704-622

L111021-082825

MANUFACTURED FOR: LOVELAND PRODUCTS, INC.®, P.O. BOX 1286, GREELEY, COLORADO 80632-1286



FIRST AID	
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If Inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<p>FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-800-424-9300. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Note to Physician: There is no specific antidote for atrazine. If this product is ingested, induce emesis or lavage stomach. The use of an aqueous slurry of activated charcoal may be considered.</p>	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid breathing dust. Causes eye irritation. Avoid contact with eyes, skin or clothing.

Personal Protective Equipment:

Mixers, loaders, cleaners of equipment or spills, and other handlers exposed to the concentrate must wear:

- Coveralls over long-sleeved shirt and long pants,
- Chemical-resistant gloves made of Barrier Laminate, Butyl Rubber \geq 14 mils, Nitrile Rubber \geq 14 mils, Neoprene Rubber \geq 14 mils, Polyvinyl Chloride (PVC) \geq 14 mils, Viton \geq 14 mils,
- Chemical-resistant footwear plus socks,
- Chemical-resistant apron, and
- Wear a minimum of a NIOSH-approved elastomeric half mask respirator with organic vapor (OV) cartridges and combination R, or P filters; OR a NIOSH-approved gas mask with OV canisters; OR a NIOSH-approved powered air purifying respirator with OV cartridges and combination HE filters.

Applicators using spray equipment mounted on their backs must wear:

- Coveralls over long-sleeved shirt and long pants,
- Chemical-resistant gloves made of Barrier Laminate, Butyl Rubber \geq 14 mils, Nitrile Rubber \geq 14 mils, Neoprene Rubber \geq 14 mils, Polyvinyl Chloride (PVC) \geq 14 mils, Viton \geq 14 mils,
- Chemical-resistant footwear plus socks.

All other applicators and all other handlers exposed to the dilute must wear:

- Long-sleeved shirt and long pants,
- Shoes plus socks, and
- Chemical-resistant gloves made of Barrier Laminate, Butyl Rubber \geq 14 mils, Nitrile Rubber \geq 14 mils, Neoprene Rubber \geq 14 mils, Polyvinyl Chloride (PVC) \geq 14 mils, Viton \geq 14 mils

See engineering controls for additional requirements.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Engineering controls statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets with

requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and can enter groundwater which may be used as drinking water. Atrazine has been found in groundwater. Users are advised not to apply atrazine to sand and loamy sand soils where the water table (groundwater) is close to the surface and where these soils are very permeable, i.e., well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

NON-TARGET ORGANISM ADVISORY STATEMENT: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

Product must not be mixed or loaded within 50 feet of intermittent streams and rivers, natural or impounded lakes and reservoirs. Product must not be applied within 66 feet of points where field surface water runoff enters perennial or intermittent streams and rivers or within 200 feet of natural or impounded lakes and reservoirs. If this product is applied to highly erodible land, the 66 foot buffer or setback from runoff entry points must be planted to crop, or seeded with grass or other suitable crop.

Product must not be mixed or loaded, or used within 50 feet of all wells, including abandoned wells, drainage wells, and sink holes. Operations that involve mixing, loading, rinsing, or

washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited, unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water, and rain water that may fall on the pad.

Surface water shall not be allowed to either flow over or from the pad which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above-specified minimum containment capacities do not apply to vehicles when delivering pesticide to the mixing/loading sites.

Additional State imposed requirements regarding well-head setbacks and operational area containment must be observed.

One of the following restrictions must be used in applying atrazine to tile-terraced fields containing standpipes:

- Do not apply within 66 feet of standpipes in tile-outletted terraced fields.
- Apply this product to the entire tile-outletted terraced field and immediately incorporate it to a depth of 2 to 3 inches in the entire field.
- Apply this product to the entire tile-outletted terraced field under a no-till practice only when a high crop residue management practice is practiced. High crop residue management is described as a crop management practice where little or no crop residue is removed from the field during and after crop harvest.

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Restrictions:

- **Aerial application is prohibited.**
- Not for use in the states of Hawaii or Alaska, or in the U.S. territories (Puerto Rico, Guam, American Samoa, the U.S. Virgin Islands, and the North Mariana Islands)
- Use on roadsides; Conservation Reserve Program (CRP) land; conifers, including Christmas Tree plantings; timber; forestry; and, Miscanthus and other perennial bioenergy crops is prohibited.
- Users must only apply to fallow land in the following states according to the prescribed rotation pattern in the table below:

Fallow Rotation Pattern	Fallow Use Authorized in these States Only
Wheat-Corn-Fallow	CO, KS, ND, NE, SD & WY
Wheat-Fallow-Wheat	CO, KS, ND, NE, SD & WY
Wheat-Sorghum-Fallow	AR, CO, GA, IL, KS, LA, MS, MO, NE, NM, NC, OK, SD & TX

ANY USE OF THIS PRODUCT IN AN AREA WHERE USE IS PROHIBITED IS A VIOLATION OF FEDERAL LAW. Before using this product, you must consult the Atrazine Watershed Information Center (AWIC) to determine whether the use of this product is prohibited in your watershed. AWIC can be accessed through www.atrazine-watershed.info, or 1-866-365-3014. If use of this product is prohibited in your watershed, you may return this product to your point of purchase or contact Loveland Products, Inc. for a refund.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: if the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls,
- Shoes plus socks, and
- Chemical-resistant gloves made of Barrier Laminate, Butyl Rubber ≥ 14 mils, Nitrile Rubber ≥ 14 mils, Neoprene Rubber ≥ 14 mils, Polyvinyl Chloride (PVC) ≥ 14 mils, Viton ≥ 14 mils.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter until sprays have dried.

PRODUCT INFORMATION

Atrazine 90 WDG Herbicide controls* many annual broadleaf and grass weeds in the crops

included on this label. Atrazine 90 WDG Herbicide may be applied before or after weeds emerge.

WEED RESISTANCE MANAGEMENT MODE OF ACTION (MOA)

Atrazine 90 WDG Herbicide contains the active ingredients atrazine. Atrazine is a triazine herbicide (Group 5 mode of action).

Atrazine's main mechanism of action is the Inhibition of photosynthesis.

Contact your local extension agent, crop advisor, or sales representative to find out if suspected resistant weeds to this MOA has been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective mechanisms of actions for each target weed. Use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner.

A given weed population may contain or develop resistance to an herbicide or herbicide MOA after repeated use. Appropriate resistance-management strategies should be followed to mitigate or delay resistance. If levels of control provided by applications of this product is reduced and cannot be accounted for by factors such as misapplication, abnormal levels of target species or extremes of weather, it may be the case that target species have developed a strain resistant to applications of this product.

Suspected herbicide-resistant weeds may be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

If resistance develops, this product may not provide sufficient control of target species. Where you suspect target species are developing resistance, contact State/local agricultural advisors. If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available, or by a mechanical method such as hoeing or tillage. Integrated weed management guidelines promote an economically viable, environmentally

sustainable, and socially acceptable weed control program regardless of the herbicide(s) used. The highlights of successful integrated weed management include:

1. Correctly identify weeds and look for trouble areas within field to identify resistance indicators.
2. Rotate crops.
3. Start the growing season with clean fields.
4. Rotate herbicide modes of action by using multiple modes of action during the growing season and apply no more than 2 applications of a single herbicide mode of action to the same field in a 2-year period. One method to accomplish this is to rotate herbicide tolerant trait systems.
5. Apply listed rates of herbicides to actively growing weeds at the correct time with the right application techniques.
6. Control any weeds that may have escaped the herbicide application.
7. Thoroughly clean field equipment between fields.
8. Scout before and after application to monitor weed populations for early signs of resistance development.
9. Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.

Contact your local agronomic advisor for more specific information on integrated weed management for your area. Users should report lack of performance to registrant or their representative. For mixtures including this herbicide note that each listed weed may not be controlled by multiple mechanisms of action. Refer to crop specific directions (below) for maximum application rates and number of applications.

***Important Note:** Following many years of continuous use of this product and chemically related products, biotypes of some of the weeds listed on this label have been reported which cannot be effectively controlled by this and related herbicides. These weeds may include lambsquarters, pigweed, black nightshade, kochia, and others. Where this is known

or suspected, we recommend the use of this product in combination with other registered herbicides which are not triazines. Consult your State Agricultural Extension Service about specific weed resistance to atrazine in your area and weed control recommendations. Within the specific crop directions on this label, certain weeds are indicated for suppression only. Weed suppression is a visual reduction in weed competition (reduced population and/or vigor) as compared to an untreated area. Degree of suppression will vary with rate used, size of weeds, and environmental conditions following treatment.

In each case where a range of rates is given, the lower rate should be used on light soils and soils low in organic matter, and the higher rate should be used on heavy soils high in organic matter.

Since Atrazine 90 WDG Herbicide acts mainly through root absorption, its effectiveness depends on rainfall or irrigation to move it into the root zone. Should weeds develop, a shallow cultivation or rotary hoeing will generally result in better weed control. Care should be taken to avoid using Atrazine 90 WDG Herbicide where adjacent desirable trees, shrubs, or plants might be injured.

Endangered Species: It is a Federal offense to use any pesticide in a manner that results in an unauthorized "take" (e.g., kill or otherwise harm) of an endangered species under the Endangered Species Act section 9. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the area in which you are applying the product. You must obtain a Bulletin no earlier than six months before using this product. To obtain Bulletins, consult <http://www.epa.gov/espp/>, call 1-844-447-3813, or email ESPP@epa.gov. You must use the Bulletin valid for the month in which you will apply the product.

Note: Loveland Products Inc. does not recommend the use of Atrazine 90 WDG Herbicide in combination with other herbicides or oils except as specifically described on this label or supplemental labeling distributed by Loveland Products, Inc.

APPLICATION PROCEDURES

Ground application: Use conventional ground sprayers equipped with nozzles that provide accurate and uniform application. Be certain that nozzles are uniformly spaced and are the same size. Calibrate sprayer before use and recalibrate at the start of each season and when

changing carriers. Unless otherwise specified, use a minimum of 10.0 gallons of spray mixture per acre for all preplant incorporated, preplant surface, preemergence, and postemergence applications (with or without oil or surfactant) with ground equipment.

Use a pump with capacity to:

1. Maintain 35 to 40 psi at nozzles,
2. Provide sufficient agitation in tank to keep mixture in suspension, and
3. To provide a minimum of 20% bypass at all times. Use centrifugal pumps which provide propeller shear action for dispersing and mixing this product. The pump should provide a minimum of 10.0 gallons per minute per 100 gallons tank size circulated through a correctly positioned sparger tube or jets.

Use screens to protect the pump and to prevent nozzles from clogging. Screens placed on the suction side of the pump should be 16-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles. Check nozzle manufacturer's recommendations.

For band applications, calculate amount to be applied per acre as follows:

$$\begin{array}{rclcl} \text{band width in inches} & \times & \text{broadcast rate} & = & \text{amount needed} \\ \text{row width in inches} & & \text{per acre} & & \text{per acre of field} \end{array}$$

Chemigation: Do not apply this product through any type of irrigation system.

Application in water: Mixing procedures:

1. Be sure sprayer is clean and not contaminated with 2,4-D, residual oil, or other materials.
2. Fill tank 1/2 to 2/3 full with clean water.
3. Start agitation.
4. Pour this product directly from package into tank. Let it wet and settle into the water.
5. Finish filling tank with water. Continue agitation during mixing and application.

Application in liquid fertilizer: Nitrogen solution or complete liquid fertilizer may replace all or part of the water as a carrier for preemergence, preplant incorporated, or preplant surface ground application on corn and sorghum. Check the compatibility of this product with liquid fertilizer and/or nitrogen solution as shown below before use. Do not apply in nitrogen

solution or complete liquid fertilizer after corn or sorghum emerges, except as noted under **Postemergence** for corn, or crop injury may occur.

Compatibility Test: Since liquid fertilizers can vary, even within the same analysis, always **check compatibility with herbicide(s) each time before use.** Be especially careful when using complete suspension or fluid fertilizers as serious compatibility problems are more likely to occur. Commercial application equipment may improve compatibility in some instances. The following test assumes a spray volume of 25.0 gallons per acre. For other spray volumes, make appropriate changes in the ingredients. Check compatibility using this procedure:

1. Add 1.0 pint of fertilizer to each of 2 one-quart jars with tight lids.
2. To **one** of the jars add 1/4 teaspoon or 1.2 milliliters of a compatibility agent approved for this use, such as Compex®, E-Z Mix, or Unite™ (1/4 teaspoon is equivalent to 2.0 pints per 100 gallons spray). Shake or stir gently to mix.
3. To **both** jars add the appropriate amount of herbicide(s). If more than one herbicide is used, add them separately with dry herbicides first, flowables next, and emulsifiable concentrates last. After each addition, shake or stir gently to thoroughly mix. The appropriate amount of herbicides for this test follows:

Dry herbicides: For each pound to be applied per acre, add 1.5 level teaspoons to each jar.

Liquid herbicides: For each pint to be applied per acre, add 0.5 teaspoon or 2.5 milliliters to each jar.

4. After adding all ingredients, put lids on and tighten, and invert each jar ten times to mix. Let the mixtures stand 15 minutes and then look for separation, large flakes, precipitates, gels, heavy oily film on the jar, or other signs of incompatibility. Determine if the compatibility agent is needed in the spray mixture by comparing the two jars. If either mixture separates, but can be remixed readily, the mixture can be sprayed as long as good agitation is used. If the mixtures are incompatible, test the following methods if improving compatibility: A) slurry the dry herbicide(s) in water before addition, or B) add 1/2 of the compatibility agent to the fertilizer and the other 1/2 to the emulsifiable concentrate or flowable herbicide before addition to the mixture. If incompatibility is still observed, do not use the mixture.

Application in water plus emulsifiable oil or oil concentrate: Adding emulsifiable oil (petroleum-derived, petroleum-derived oil concentrate, or single or mixed crop-derived oil

concentrate) to postemergence water-based sprays in corn and sorghum may improve weed control. However, under certain conditions, the use of either type of oil may seriously injure the crop. To minimize this possibility, observe all following directions.

Use one of the following properly emulsified:

- A suitable oil concentrate containing at least 1% but not more than 20% suitable emulsifier or surfactant blend.
- Petroleum-derived oil containing at least 1% suitable emulsifier.

Note: In the event of a compatibility problem when mixing oil with Atrazine 90 WDG Herbicide and water, a compatibility agent such as Compex, E-Z Mix, or Unite should be used. Any of the above oils contaminated with water or other materials can cause compatibility problems and/or crop injury.

Mixing procedures-all uses:

1. Be sure sprayer is clean and not contaminated with any other materials, or crop injury or sprayer clogging may result.
2. Fill tank 1/4 full with clean water, nitrogen solution, or complete liquid fertilizer.
3. Start agitation.
4. Be certain that the agitation system is working properly and creates a rippling or rolling action on the liquid surface.
5. Pour product directly from package into tank.
6. Continue filling until 90% full. Increase agitation if necessary to maintain surface action.
7. Add emulsifiable oil, oil concentrate, or tank mix herbicide(s) after this product is thoroughly suspended.
8. Finish filling tank.
9. Empty tank as completely as possible before refilling to prevent buildup of oil or emulsifiable concentrate residue. Maintain agitation to avoid separation of materials.
10. If an oil or emulsifiable concentrate film starts to build up in tank, drain it and clean with strong detergent solution or solvent.
11. Clean sprayer thoroughly immediately after use by flushing system with water containing a detergent.

APPLICATION RESTRICTIONS

- Aerial application is prohibited.

- Do not apply this product through any type of irrigation system.
- When tank-mixing or sequentially applying atrazine or products containing atrazine to corn or sorghum, the total pounds atrazine applied (pounds active ingredient per acre) must not exceed 2.5 pounds active ingredient per acre per year.
- When tank-mixing or sequentially applying atrazine or products containing atrazine to crops other than corn or sorghum, the total pounds of atrazine applied (pounds active ingredient per acre) must not exceed the specific seasonal rate limits as noted in the use directions.

MANDATORY SPRAY DRIFT MANAGEMENT

Ground Boom Applications:

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to use a coarse or coarser droplet size (ASABE S572).
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- User must maintain a 15 foot (4.6 m) in-field downwind buffer (in the direction in which the wind is blowing) from the edge of streams and rivers, as well as high-tide line for all estuarine/marine environments.
- Do not apply during temperature inversions.

Boomless Ground Applications:

- Applicators are required to use a medium or coarser droplet size (ASABE S572) for all applications.
- User must maintain a 15 foot (4.6 meter) in-field downwind buffer (in the direction in which the wind is blowing) from the edge of streams and rivers, as well as high-tide line for all estuarine/marine environments.
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

BOOM HEIGHT – Ground Boom

- For ground equipment, the boom should remain level with the crop and have minimal bounce.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

BOOMLESS GROUND APPLICATIONS:

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

HANDHELD TECHNOLOGY APPLICATIONS:

Take precautions to minimize spray drift.

ROTATIONAL CROP RESTRICTIONS FOR ALL USES

1. Land treated with Atrazine 90 WDG Herbicide should not be planted to any crop except corn or sorghum until the following year or injury may occur.
2. If Atrazine 90 WDG Herbicide is applied after June 10, do not rotate with crops other than corn or sorghum the next year or injury may occur.
3. In the High Plains and Intermountain area of the West where rainfall is sparse and erratic or where irrigation is required, use Atrazine 90 WDG Herbicide only when corn or sorghum is to follow corn or sorghum, or a crop of untreated corn or sorghum is to precede other rotational crops.
4. In eastern parts of the Dakotas, KS, western MN, and NE, do not rotate to soybeans if the rate applied to corn or sorghum was more than 2.2 pounds per acre or equivalent band application rate, as soybean injury may occur.
5. Injury may occur to soybeans planted in north-central and northwest IA, southcentral and southwest MN, northeast NE, southeast SD, and other areas the year following application on soils having a calcareous surface layer.
6. Do not plant sugar beets, tobacco, vegetables (including dry beans), spring-seeded small grains or small-seeded legumes and grasses the year following Atrazine 90 WDG Herbicide application for injury may occur.

CORN AND SORGHUM WEED CONTROL/SUPPRESSION

Preplant Surface-Applied, Preplant Incorporated, or Pre-emergence (or Postemergence at 2.2 pounds per acre with Oil)

Broadleaf and Grass Weeds Controlled

annual morningglory	groundcherry	mustards	velvetleaf**
barnyardgrass**	jimsonweed	nightshade*	(buttonweed)
(watergrass)	kochia*	pigweed*	wild oats
cocklebur*	lambsquarters*	purslane	witchgrass**
giant foxtail*	large (hairy)	ragweed	(<i>Panicum capillare</i>)
green foxtail**	crabgrass*	sicklepod	yellow foxtail*

Postemergence with Emulsifiable Oil or Oil concentrate in Water (at 1.3 pounds per acre)

Broadleaf Weeds Controlled

annual morningglory	lambsquarters*	ragweed	wild buckwheat
cocklebur*	mustards	smartweed	
jimsonweed	pigweed*	velvetleaf*	

Note: For best control of Cocklebur and Velvetleaf, do not apply less than the specified 2.2 pounds per acre.

* Maximum rate limitations and local resistance may result in suppression only or lack of control. Refer to IMPORTANT NOTE within PRODUCT INFORMATION section of this label.

** Suppression or partial control only on medium and fine textured soils.

MAXIMUM RATES PRIOR TO CORN OR SORGHUM EMERGENCE

For single preemergent broadcast applications (including early preplant, preplant incorporated, preplant surface, at planting or preemergence) the following maximum use rates apply:

Soil Erodibility Classification*	Plant Residue Amount	Maximum Rate/Acre
Highly erodible	30% or more (conservation tillage program)	2.0 lbs AI
	Less than 30% (conventional tillage program)	1.6 lbs AI
Not highly erodible	No limit	2.0 lbs AI

*As defined by the Natural Resource Conservation Service

MAXIMUM RATES AFTER CORN OR SORGHUM EMERGENCE

For postemergence applications, the following maximum rates apply to the total of all atrazine treatments:

Previous Soil Application Of Atrazine To Crop	Total Maximum Rate/Acre/Calendar Year
No	2.0 lbs AI postemergence
Yes	2.5 lbs AI combined from soil and postemergence applications

Atrazine 90 WDG Herbicide contains 0.9 lb AI/lb.

CORN

Atrazine 90 WDG Herbicide may be applied before planting, at planting or after planting, subject to maximum rate limitations.

Postemergence applications to corn must be made before corn reaches 12 inches in height.

Preplant Surface-Applied (Broadleaf and grass control): Use on medium- and fine-textured soils with minimum-tillage or no-tillage systems only in CO, IL, IN, IA, KS, KY, MN, MO, MT, NE, ND, SD, WI, and WY. Apply the specified rate of Atrazine 90 WDG Herbicide shown in Table 1 up to 30 days prior to planting. On *coarse-textured* soils, do not apply more than two weeks prior to planting.

If weeds are present at the time of treatment, apply in tank mix combination with a contact herbicide (for example, paraquat [Gramoxone Inteon®] or glyphosate [Makaze®]). Observe directions for use, precautions, and restrictions on the label of the contact herbicide.

Note: To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.

Preplant Incorporated (Broadleaf and grass control): Broadcast in spring after plowing, at rate in Table 1A. Apply to the soil and incorporate before, during, or after final seedbed preparation. Avoid deep incorporation. For best results, apply within two weeks prior to planting.

Preemergence or At-planting (Broadleaf and grass control): Apply during or shortly after planting before weed emergence, at rate in Table 1A.

Postemergence (Broadleaf and grass control): Apply before weeds are 1.5 inches tall and before corn reaches 12 inches tall, at rate in Table 1B. When using nitrogen solutions, direct the spray to avoid corn foliage injury.

TABLE 1: ATRAZINE 90 WDG HERBICIDE RATES FOR BROADLEAF AND GRASS CONTROL IN CORN

A. FOR ALL SOIL APPLICATIONS PRIOR TO CORN EMERGENCE

(including early preplant, preplant incorporated, preplant surface, at planting or preemergence)

Soil Erodibility Classification*	Plant Residue Amount	Maximum Rate/Acre ¹
Highly erodible	30% or more (conservation tillage program)	2.2 lbs Atrazine 90 WDG Herbicide
	Less than 30% (conventional tillage program)	1.77 lbs Atrazine 90 WDG Herbicide
Not highly erodible	No limit	2.2 lbs Atrazine 90 WDG Herbicide

*As defined by the Natural Resource Conservation Service

B. FOR CORN POSTEMERGENCE APPLICATION

Previous Soil Application Of Atrazine To Crop	Total Maximum Rate/Acre/Calendar Year
No	2.2 lbs this product postemergence
Yes	2.77 lbs this product combined from soil and postemergence applications

Postemergence applications to corn must be made before corn reaches 12 inches in height.

¹**Broadleaf control (eastern CO, western KS, western NE, NM, OK panhandle, west TX, and eastern WY):** On sand, loamy sand, sandy loam, mild to strongly alkaline soil, and all recently leveled soil, apply no more than 1.3 pounds this product per acre, either preplant surface, preplant incorporated, or preemergence. On other soils in these areas, apply rate in

Table 1 for broadleaf and grass control.

Postemergence with emulsifiable oil or oil concentrate in water: Add the following volume of one of the type oils indicated for ground application unless the oil label specifies otherwise:

Type Oil	Ground Application
Oil concentrate (Crop or Petroleum-derived)	1.0 qt/acre
Petroleum-derived oil	1.0 gal/acre

Note: Crop-derived or petroleum-derived oil concentrates should contain at least 1% but not more than 20% suitable emulsifier or surfactant blend. Petroleum-derived oils should contain at least 1% suitable emulsifier.

Broadleaf and grass control: For postemergence control of those weeds listed under **Preplant Incorporated and Preemergence**, broadcast 2.2 pounds this product per acre plus emulsifiable oil or oil concentrate after weed emergence, but before weeds reach 1.5 inches in height and before corn is 12 inches tall.

Broadleaf control: For postemergence control of those weeds listed under **Postemergence with emulsifiable oil or oil concentrate in water**, broadcast 1.3 pounds this product per acre plus emulsifiable oil or oil concentrate before pigweed and lambsquarters reach 6 inches in height, before all other weeds reach 4 inches in height and before corn is 12 inches tall. A cultivation may be necessary if all weeds are not controlled or if weeds regrow.

Use precautions for applications with emulsifiable oil or oil concentrate in water:

1. Inbred lines or any breeding stock may be severely injured by applications with emulsifiable oil or oil concentrate.
2. Adding other insecticides, herbicides, liquid fertilizers, or other materials is not recommended, because they may cause compatibility problems or crop injury.
3. Store and handle emulsifiable oil and oil concentrate carefully. Oil contaminated with even a small amount of water may not emulsify properly when added to the tank. To avoid crop injury,
4. Do not apply when crop is under stress from prolonged cold, wet weather, poor fertility, or other factors, or when crop is wet and succulent from recent rainfall.

5. Do not make more than one application per season.

TANK MIXTURES FOR CORN

This product may be tank mixed with other herbicides for control of certain broadleaf and grass weeds in corn: Use tank mix directions appearing on the labels of other herbicides when tank mixing with this product. Observe all precautions and limitations on labeling of products used in a particular tank mix.

Atrazine 90 WDG Herbicide + Simazine 4L or Simazine 90DF

Use a tank mixture of Atrazine 90 WDG Herbicide + simazine for the control of crabgrass, fall panicum and carpetweed in addition to the control of the weeds listed for Atrazine 90 WDG Herbicide alone.

Make broadcast application of tank mix before planting, at planting, or after planting, but prior to emergence of corn and weeds. Cultivate shallowly if weeds develop.

PREPLANT SURFACE-APPLIED: Use on medium- and fine-textured soils in minimum-tillage or no-tillage systems only in CO, IL, IN, IA, KS, KY, MN, MO, MT, NE, ND, SD, WI, and WY. Apply up to 30 days before planting. Applications made less than 30 days before planting may be made as either a split or single treatment. On coarse-textured soils, do not apply more than two weeks before planting.

If weeds are present at time of treatment, apply in a tank mix combination with a contact herbicide (for example, paraquat [Gramoxone Inteon] or glyphosate [Makaze]). All applicable directions, use precautions and limitations on the contact herbicides' labels must be followed.

Note: To the extent possible do not move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.

PREPLANT INCORPORATED: Make soil application and incorporate in the spring before, during, or after final seedbed preparation. Avoid deep incorporation. For best results, apply within two weeks before planting.

PREEMERGENCE: Apply during or shortly after planting, but prior to crop and weed emergence.

All applicable directions, use precautions, and limitations on this and the simazine label

must be followed.

When tank mixing or sequentially applying simazine and/or atrazine or products containing either active ingredient to corn, the total pounds of simazine and/or atrazine applied must not exceed 2.5 pounds active ingredient per acre per calendar year.

Atrazine 90 WDG Herbicide + Simazine 4L, or Simazine 90DF with glyphosate (Makaze)

Use a tank mixture of Atrazine 90 WDG Herbicide + simazine + glyphosate for control of certain broadleaf weeds and grasses where corn will be planted directly into a cover crop, established sod, or in previous crop residues. Refer to glyphosate label for rates and tank mix directions. All applicable directions, limitations, and use precautions on this, the simazine, and the glyphosate label must be followed.

Atrazine 90 WDG Herbicide + Simazine 4L, or Simazine 90DF with paraquat (Gramoxone Inteon)

Use a tank mixture of Atrazine 90 WDG Herbicide + simazine + paraquat to control most emerged weeds and for residual weed control where corn will be planted directly into a cover crop, established sod or in previous crop residues. Add Atrazine 90 WDG Herbicide and simazine to water in spray tank. Agitate until thoroughly mixed. Next add paraquat and a nonionic surfactant, such as Loveland Activator 90, or similar nonionic surfactant approved for agricultural use.

Continue agitation during application. Apply 1.1 pounds of Atrazine 90 WDG Herbicide plus 2.0 pints Simazine 4L, or 1 1/10 pounds Simazine 90DF plus the specified rate of paraquat in 20.0 to 60.0 gallons of water per sprayed acre. Make broadcast application before, during, or after planting, but before emergence of corn. Add 1/2 pint of a nonionic surfactant per 100 gallons of spray mixture. Use the higher rate of paraquat if existing weeds are 4 to 6 inches tall. Weeds taller than 6 inches will not be controlled.

All applicable directions, limitations, and use precautions on this, the simazine and the paraquat labels must be followed.

Atrazine 90 WDG Herbicide + paraquat (Gramoxone Inteon)

For control of existing vegetation and residual control where corn will be planted directly into cover crop, established sod or in previous crop residues-Broadcast 2.2 pounds Atrazine 90

WDG Herbicide and specified label rate of paraquat in 20.0 to 60.0 gallons of water per sprayed acre. Following the paraquat label, add specified rate of nonionic surfactant per 100 gallons of diluted spray. Add Atrazine 90 WDG Herbicide to spray tank first and thoroughly mix with water. Add the paraquat and surfactant last.

Refer to the paraquat label for further directions, limitations and cautions.

Atrazine 90 WDG Herbicide + alachlor (Micro-Tech®)

Use a tank mixture of Atrazine 90 WDG Herbicide + alachlor for the control of most annual broadleaf and grass weeds in corn (field and silage corn only) such as annual morningglory, barnyardgrass, black nightshade, brachiaria, buttonweed (velvetleaf), carpetweed, cocklebur, crabgrass, fall panicum, Florida pusley, giant foxtail, green foxtail, yellow foxtail, goosegrass, lambsquarters, pigweed, purslane, mustard, common ragweed, smartweed and witchgrass. Applications may be made preplant, within 7 days of planting; preemergence, or postemergence, until weeds reach the two leaf stage and the corn is not more than 5 inches tall.

Refer to the alachlor label for application rates and directions. Applicable limitations and use precautions on this and the alachlor label must also be followed.

Atrazine 90 WDG Herbicide + alachlor (Micro-Tech) + paraquat (Gramoxone Inteon) or glyphosate (Makaze)

For Minimum-Tillage or No-Tillage Systems

For control of many emerged annual weeds, suppression of many emerged perennial weeds, and preemergence control of many annual grasses and weeds in areas where corn will be planted directly into a cover crop, established sod or in previous crop residues. Regrowth from perennial weeds will not be controlled. Do not apply by air.

Make application immediately before, during, or after planting, but before crop emergence. Refer to the alachlor label for application rates and directions. All applicable directions, limitations, and use precautions on this and the alachlor and paraquat or glyphosate labels must be followed.

Atrazine 90 WDG Herbicide + metolachlor/S-metolachlor (Dual II MAGNUM®) or Atrazine 90 WDG Herbicide + metolachlor/S-metolachlor (Dual II MAGNUM) + simazine

PREPLANT/PREEMERGENCE

Use a tank mixture of Atrazine 90 WDG Herbicide + metolachlor/S-metolachlor or Atrazine 90 WDG Herbicide + metolachlor/S-metolachlor + simazine as a preplant surface, preplant incorporated or preemergence application for the control of barnyardgrass, crabgrass, crowfootgrass, fall panicum, foxtail millet, giant foxtail, goosegrass, green foxtail, prairie cupgrass, red rice, signalgrass (*Brachiaria*), southwestern cupgrass, witchgrass, yellow foxtail, yellow nutsedge, black nightshade, carpetweed, Florida pusley, galinsoga, pigweed, browntop panicum, cocklebur, common purslane, hairy nightshade, lambsquarters, morningglory, ragweed, smartweed, and velvetleaf in corn.

To apply, refer to the tank mix directions appearing on the metolachlor/S-metolachlor label. Note, where directions specify AAtrex® Nine-O®, substitute Atrazine 90 WDG Herbicide. All applicable directions, limitations, and use precautions on this and the metolachlor/S-metolachlor and simazine labels must be followed.

POSTEMERGENCE

Use a tank mixture of Atrazine 90 WDG Herbicide plus metolachlor/S-metolachlor as a postemergence application for the control of barnyardgrass, crabgrass, crowfootgrass, fall panicum, foxtail (giant, green and yellow), jimsonweed, mustard, pigweed, prickly sida, purslane, ragweed, smartweed, and velvetleaf and for partial control of cocklebur, morningglory, and yellow nutsedge. To apply, follow the tank mix directions appearing on the metolachlor/S-metolachlor label. Note, where directions specify AAtrex Nine-O, substitute Atrazine 90 WDG Herbicide. All applicable directions, limitations, and precautions on the metolachlor/S-metolachlor label must be followed.

Atrazine 90 WDG Herbicide + metolachlor/S-metolachlor (Dual II MAGNUM) + paraquat (Gramoxone Inteon) or Atrazine 90 WDG Herbicide + metolachlor/S-metolachlor (Dual II MAGNUM) + glyphosate (Makaze) or Atrazine 90 WDG Herbicide + metolachlor/S-metolachlor (Dual II MAGNUM) + simazine + paraquat (Gramoxone Inteon) or Atrazine 90 WDG Herbicide + metolachlor/S-metolachlor (Dual II MAGNUM) + simazine + glyphosate (Makaze).

For Minimum-Tillage or No-Tillage Systems

Use a tank mixture of Atrazine 90 WDG Herbicide + metolachlor/S-metolachlor with paraquat or glyphosate, or Atrazine 90 WDG Herbicide + metolachlor/S-metolachlor + simazine with paraquat or glyphosate in minimum-tillage or no-tillage systems where corn is planted directly into a cover crop, stale seedbed, established sod, or previous crop residues.

The paraquat tank mixtures will control most emerged annual weeds and suppress many perennial weeds, in addition to controlling most preemergence annual broadleaf weeds and grasses. The glyphosate tank mixtures will control emerged annual and perennial weeds, in addition to controlling most preemergence annual broadleaf weeds and grasses. Make application before, during, or after planting but before the corn emerges. To apply, follow the tank mix directions appearing on the metolachlor/S-metolachlor label. Note, where directions specify AAtrex Nine-O, substitute Atrazine 90 WDG Herbicide. All application directions, limitations, and use precautions on this and the metolachlor/S-metolachlor, simazine, paraquat and glyphosate labels must be followed.

USE RESTRICTIONS FOR ALL APPLICATIONS TO CORN AND SORGHUM

- Do not exceed an application rate of 2.0 pounds active ingredient of atrazine per acre for any single application, and the total pounds of atrazine applied (pounds active ingredient per acre) must not exceed 2.5 pounds active ingredient per acre per year.
- Do not apply product 60 days prior to harvest for field corn forage and preemergent sorghum forage.
- Do not apply product 45 days prior to harvest for sweet corn forage and postemergent sorghum.
- Do not apply atrazine and propazine products to the same sorghum acre.
- Following harvest of a treated crop, plow (moldboard or disk-plow) and thoroughly till the soil in the fall or spring to minimize possible injury to rotational spring-seeded crops, regardless of the rate used.
- Do not graze treated area or feed treated forage to livestock for 21 days following application.
- For postemergence applications plus emulsifiable oil, see additional use precautions under that section of this label.
- Postemergence application to corn and sorghum must be made before crop reaches 12

inches in height.

- Do not apply with mechanically pressurized handguns to sweet corn.

SORGHUM AND SORGHUM-SUDAN HYBRIDS (GRAIN AND FORAGE TYPES):

Postemergence applications to sorghum must be made before sorghum reaches 12 inches in height.

Atrazine 90 WDG Herbicide may be applied before planting, at planting or after planting, subject to maximum rate limitations.

Use only on medium- and fine-textured soils having a minimum of 1% organic matter, except as separately specified for furrow-irrigated bedded sorghum grown in Arizona and California.

Preplant Surface-Applied (Broadleaf and grass control): Use on medium- and fine-textured soils with minimum-tillage or no-tillage systems only in CO, IL, IN, IA, KS, KY, MN, MO, MT, NE, ND, SD, WI, and WY. Apply 2/3 the specified rate of Atrazine 90 WDG Herbicide shown in Table 3 for the soil texture as a split treatment 30 to 45 days prior to planting and the remainder at planting. Applications made less than 30 days prior to planting may be as either a split or single treatment. Under dry conditions, irrigation after application is recommended to move Atrazine 90 WDG Herbicide into the soil.

If weeds are present at time of treatment, apply in a tank mix combination with a contact herbicide (for example, paraquat [Gramoxone Inteon] or glyphosate [Makaze]). Observe directions for use, precautions, and restrictions on the label of the contact herbicide.

Note: To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.

Preplant Incorporated (Broadleaf and grass control): Broadcast in spring after plowing at rate shown in Table 3. Apply before, during, or after final seedbed preparation. If soil is tilled or worked after applications, avoid deep incorporation. For best results, apply within two weeks prior to planting.

Preemergence or At-planting (Broadleaf and grass control): Apply during or shortly after planting, but prior to weed or crop emergence at rate shown in Table 3.

TABLE 3: ATRAZINE 90 WDG HERBICIDE RATES FOR BROADLEAF AND GRASS CONTROL IN SORGHUM.

FOR ALL SOIL APPLICATIONS PRIOR TO SORGHUM EMERGENCE (including early preplant, preplant incorporated, preplant surface, at planting or preemergence)

Soil Erodibility Classification*	Plant Residue Amount	Maximum Rate/Acre ¹
Highly erodible	30% or more (conservation tillage program)	2.2 lbs Atrazine 90 WDG Herbicide
	Less than 30% (conventional tillage program)	1.77 lbs Atrazine 90 WDG Herbicide
Not highly erodible	No limit	2.2 lbs Atrazine 90 WDG Herbicide

* As defined by the Natural Resource Conservation Service

¹ Do not apply preplant surface or preplant incorporated in AL, AR, FL, GA, LA, MS, NC, NM, OK, SC, TN, or TX. Do not apply preemergence in NM, OK, or TX, except in northeast OK and the TX Gulf Coast. In case of planting failure, sorghum or corn may be replanted. Do not make a second broadcast application, or injury may occur. If originally applied in a band and sorghum or corn is replanted in untreated row middles, this product may be applied in a band to the second planting provided the maximum application rate of 2.5 pounds active ingredient of this product per acre atrazine per calendar year is not exceeded.

Preemergence broadleaf weed control in furrow irrigated bedded sorghum (AZ and CA only): For preemergence control of many broadleaf weeds, broadcast 0.9 to 1.3 pounds of this product per acre after bed preparation, during or after planting, but before sorghum and weeds emerge and before the first furrow irrigation. Follow with several regular irrigations, making sure to thoroughly wet all soil.

Use precautions for preemergence application to furrow irrigated bedded sorghum in AZ and CA: To avoid possible sorghum injury, do not use on sand or loamy sand soil or on sorghum planted in furrows. Applications to sorghum growing on alkali soils or where

cuts, fills, or erosion have exposed calcareous or alkaline subsoils may cause crop injury. In case of crop failure, do not replant sorghum for 8 months following application. Corn may be planted immediately.

Postemergence (Broadleaf and grass control): Apply at rate shown in Table 4 before weeds exceed 1.5 inches in height. Applications may be made before crop height of 12 inches.

TABLE 4: ATRAZINE 90 WDG HERBICIDE RATES FOR BROADLEAF AND GRASS CONTROL IN SORGHUM.

FOR SORGHUM POSTEMERGENCE APPLICATION	
Previous Soil Application Of Atrazine To Crop	Total Maximum Rate/Acre/Calendar Year
No	2.2 lbs postemergence
Yes	2.77 lbs combined from soil and postemergence applications

Postemergence applications to sorghum must be made before sorghum exceeds 12 inches in height.

Postemergence broadleaf weed control with emulsifiable oil or oil concentrate in water: Broadcast 1.3 pounds this product per acre for control of many broadleaf weeds. Apply before pigweed and lambsquarters reach 6 inches in height, before all other weeds reach 4 inches in height and before sorghum is 12 inches tall. In CO, western KS, NM, OK, TX, and desert regions of AZ and CA, apply when sorghum is about 6 to 10 inches in height, but before sorghum is 12 inches tall. In all other areas, apply after sorghum reaches the 3-leaf stage and before sorghum is 12 inches tall. Add 1.0 gallon of emulsifiable oil per acre for ground application or add 1.0 quart of oil concentrate for ground application. A cultivation may be necessary if all weeds are not controlled or if weeds regrow.

For the list of weeds controlled, see Atrazine 90 WDG Herbicide Applied Alone-Corn or Grain Sorghum-Postemergence with Emulsifiable Oil or Oil Concentrate in Water.

Use precautions for applications with emulsifiable oil or oil concentrate in water: See "Use precautions for applications with emulsifiable oil or oil concentrate in water" in Corn section.

Postemergence broadleaf weed control with surfactant (CO, western KS, NM, OK, TX, and desert regions of AZ and CA only): Broadcast 1.3 pounds this product per acre plus 0.75 to 1.5 pints of surfactant when sorghum is 6 up to 12 inches in height, but before weeds exceed 1.5 inches in height. Apply only on sandy loam and finer textured soil.

Use precautions for all applications to sorghum:

1. Heavy rain immediately following application tends to cause excessive concentrations of herbicide in seed furrow, resulting in possible crop injury. Do not apply to furrow-planted sorghum until furrows are leveled (plowed in). Level deep planter marks or seed furrows before application.
2. Application to sorghum growing under stress caused by minor element deficiency or to sorghum growing on highly calcareous soil may result in crop injury.
3. Following harvest, plow (moldboard or disk-plow) and thoroughly till soil in fall or spring to minimize possible injury to spring-seeded rotational crops, regardless of rate used.
4. Injury may occur if both this herbicide, preplant surface, preplant incorporated, or preemergence, and an at-planting systemic insecticide applied in-furrow are used.
5. Do not apply more than 2.2 to 2.77 pounds of this product per acre per calendar year (See MAXIMUM RATES section of label).
6. For all soil applications prior to crop emergence (except for preemergence use on bedded sorghum in AZ and CA), do not apply to coarse-textured soils, i.e., sand, loamy sand, sandy loam, or to medium- and fine-textured soils having less than 1% organic matter, or injury may occur.
7. For postemergence applications, do not apply to sand or loamy sand, or injury may occur.

Note: Do not graze or feed forage from treated areas for 21 days following application, or illegal residues may result.

TANK MIXTURES FOR GRAIN SORGHUM

Metolachlor/S-metolachlor (Dual MAGNUM®): Use as tank mixture with metolachlor/S-metolachlor for control of those weeds listed on the metolachlor/S-metolachlor label, as well as on this label. Use this tank mixture only on sorghum seed treated with Concep®. Refer to the

metolachlor/S-metolachlor label for all directions, precautions, and limitations.

WINTER WEED CONTROL IN TEXAS

For postemergence control of winter weeds only, such as henbit, seedling dock, and annual thistle on fall bedded land in the Gulf Coast and Blacklands of Texas. Apply 0.9 to 1.1 pounds of this product per acre postemergence to the weeds in November or December to land that will be planted to corn, grain sorghum, or forage sorghum the following spring. For best results add a suitable surfactant at the rate of 0.5% of the spray volume, an emulsifiable oil at the rate of 1.0% of the spray volume, or an oil concentrate at the rate of 1.0 quart per acre. Normal weed control programs may be used in the following corn, grain sorghum or forage sorghum crop.

Note: Do not plant any crops except corn, grain sorghum, or forage sorghum the spring following this treatment, or illegal residues may result.

ATRAZINE 90 WDG HERBICIDE ALONE-CHEMICAL FALLOW

Wheat-Sorghum-Fallow (AR, CO, GA, IL, KS, LA, MS, MO, NE, NM, NC, OK, SD & TX): To control annual broadleaf and grass weeds following wheat harvest and in the following sorghum crop when grown under minimum tillage, broadcast 2.5 pounds this product per acre to wheat stubble immediately following wheat harvest. If weeds are present at application, remove them with a sweep plow or other suitable implement after application, or use an approved contact herbicide before or after the application of Atrazine 90 WDG Herbicide. Plant sorghum into wheat stubble the following spring with minimum soil disturbance. Use a surface planter or a planter leaving a shallow furrow. If weeds are present at planting, remove them with a sweep plow or other suitable implement before planting.

For the list of weeds controlled, see **Atrazine 90 WDG Herbicide Applied Alone-Corn or Grain Sorghum-Preplant Surface-Applied, Preplant Incorporated, or Preemergence.**

Use precautions:

1. Use only on silt loam or finer textured soil, or crop injury may result.
2. Wheat-sorghum-fallow cropping sequence must be followed.
3. Do not apply following sorghum harvest.

Notes: To avoid illegal residues, do not graze or feed forage from treated area to livestock.

To avoid illegal residues and crop injury, do not plant any crop other than those on this label within 18 months following treatment.

Wheat-Corn-Fallow (CO, KS, ND, NE, SD, and WY): This product controls cheatgrass (downy brome, chess), Kochia*, mustards, pigweed*, Russian thistle, wild lettuce, wild sunflower, and volunteer wheat during period after wheat harvest. Weed control may extend into following corn crop grown under minimum tillage.

*(Local resistance may result in suppression only or lack of control. Refer to IMPORTANT NOTE within PRODUCT INFORMATION SECTION of this label.)

Follow directions for use, notes, and use precautions in the "Wheat-Sorghum-Fallow" section above, substituting corn for references to sorghum.

Wheat-Fallow-Wheat (CO, KS, ND, NE, SD, and WY): For preemergence control of cheatgrass (downy brome, chess), common lambsquarters*, field pennycress, Kochia*, mustard, Russian thistle, wild lettuce, and suppression of volunteer wheat during fallow period of a wheat-fallow-wheat rotation, broadcast 0.5 to 1.1 pounds of this product per acre on all soils except those listed under "Use Precautions." For control of pigweed and wild sunflower, use the higher rate. Apply to stubble ground. Treat only once during same fallow period.

*(Local resistance may result in suppression only or lack of control. Refer to IMPORTANT NOTE within PRODUCT INFORMATION SECTION of this label.)

TANK MIXTURES FOR CHEMICAL FALLOW

Wheat-Sorghum-Fallow or Wheat-Corn-Fallow

Paraquat (Gramoxone Inteon): If weeds are present at application, a tank mix with paraquat may be used. Broadcast 2.5 pounds of Atrazine 90 WDG Herbicide plus specified label rate of paraquat in 20.0 to 60.0 gallons of water per acre by ground equipment. Add 0.5 to 1.0 pint of nonionic surfactant per 100 gallons of spray mixture. Add Atrazine 90 WDG Herbicide to spray tank first and thoroughly mix with water. Then add paraquat, followed by surfactant. Use the higher rate of paraquat if weeds are 4 to 6 inches tall. This mixture will not control weeds taller than 6 inches. Apply to stubble ground. Treat only once during same fallow period. Refer to paraquat label for further directions, precautions, and limitations.

Wheat-Fallow-Wheat

Paraquat (Gramoxone Inteon): If weeds are present at application, a tank mix with paraquat may be used. Broadcast 0.5 to 1.1 pounds of Atrazine 90 WDG Herbicide plus specified label rate of paraquat in 20.0 to 60.0 gallons of water per acre by ground equipment. Add 0.5 to 1.0 pint of a nonionic surfactant per 100 gallons of spray mixture. Add Atrazine 90 WDG Herbicide to spray tank first and thoroughly mix with water. Then add paraquat, followed by surfactant. Use the higher rate of paraquat if weeds are 4 to 6 inches tall. This mixture will not control weeds taller than 6 inches. Apply to stubble ground. Treat only once during same fallow period. Refer to paraquat label for further directions, precautions, and limitations. If weeds are present at application and this product is used alone, use either an approved contact herbicide before or after treatment, or tillage after treatment. Use tillage to control weeds which escape during fallow period. Till before planting. For this product applied alone or in tank mixture with paraquat, plant at least 2 inches deep and 12 months or more after application.

Use precautions: To avoid crop injury:

1. Do not use on sand soil.
2. Do not treat eroded hillsides, caliche and rocky outcroppings, or exposed calcareous subsoil.
3. Do not treat soils of the Rosebud and Canyon Series in western NE and adjoining counties in CO and WY.
4. Do not treat soils with calcareous surface layers.
5. Avoid spray overlap.

Note: Do not graze treated areas within 6 months after application or illegal residues may result.

CHEMICAL FALLOW USE RESTRICTIONS

For soils in North and South Dakota with a pH of 7.5 or greater:

- Do not apply more than 1.5 pounds active ingredient per acre for any application. Do not apply more than one application per cycle.

For soils in North and South Dakota with a pH of less than 7.5:

- Do not apply more than 2.0 pounds active ingredient per acre for any application. Do not

apply more than one application per cycle.

For all other locations:

- Do not apply more than 2.25 pounds active ingredient per acre for any application. Do not apply more than one application per cycle.

SUGARCANE

For control of many broadleaf and grass weeds, including amaranths*, crabgrass, fireweed, flora's paintbrush, foxtails*, junglerice* and wiregrass, broadcast 2.2 to 4.4 pounds of Atrazine 90 WDG Herbicide at time of planting or ratooning, but before sugarcane emerges. Broadcast or band by ground equipment in a minimum of 20.0 gallons per acre, unless indicated otherwise. One additional application may be made over the sugarcane as it emerges, and two additional applications may be made interline after emergence as directed sprays. Repeat treatments, where needed, may be applied broadcast, band, or interline as suggested with the final application being prior to close-in. Do not exceed the rate of herbicide suggested for any one crop of sugarcane.

*Local resistance may result in suppression only or lack of control. Refer to IMPORTANT NOTE within PRODUCT INFORMATION SECTION of this label.)

Use precautions for all states:

1. Do not apply more than 4.0 pounds active ingredient per acre for any application.
2. Do not apply more than 10.0 pounds active ingredient per acre per crop.

Note: Where high rates of Atrazine 90 WDG Herbicide are used, apply in a minimum of 1.0 gallon of water for each 1.0 pound of product applied per acre.

Florida

For control of emerged pellitory weed: Apply 0.4 to 0.6 pound of this product per acre in at least 40.0 gallons of water as a directed spray by ground equipment prior to close-in. Add 4.0 quarts of surfactant for each 100 gallons of spray. Thoroughly cover weed foliage.

For control of alexandergrass*, large crabgrass, pellitory (artillery) weed, and spiny amaranth*, use one of the following methods at planting or ratooning:

1. Apply 4.4 pounds of this product per acre preemergence. Follow with one or two applications, as needed, postemergence to sugarcane and weeds at 2.2 pounds this product per

- acre. Treat before weeds exceed 1.5 inches in height.
2. Apply 1 to 3 times, as needed, at 2.2 pounds this product per acre postemergence to sugarcane and weeds. Treat before weeds exceed 1.5 inches in height.

Louisiana

For control of annual weeds during the summer fallow period, apply 2.2 pounds of this product per acre to weed-free beds immediately after bed formations. Follow normal weed control program after planting.

Texas

Use Atrazine 90 WDG Herbicide for control of barnyardgrass*, pigweed*, purslane and sunflower, in plant or ratoon sugarcane.

Apply 4.4 pounds per acre of Atrazine 90 WDG Herbicide preemergence. Follow with one or two applications, as needed, at 3.3 pounds per acre postemergence to sugarcane and weeds. For best results when weeds are emerged, add a nonionic surfactant at a concentration of 2.0 quarts per 100 gallons to the spray and apply before weeds exceed 1.5 inches in height.

*(Local resistance may result in suppression only or lack of control. Refer to IMPORTANT NOTE within PRODUCT INFORMATION SECTION of this label.)

Use precautions:

1. Injury to sugarcane may occur when under moisture stress, when soil is of low absorptive capacity, or when land is first cropped to sugarcane.
2. Do not apply after close-in.
3. Do not apply more than 11.0 pounds per acre to any one crop of sugarcane, or crop injury may result.

TURFGRASSES FOR SOD

St. Augustinegrass, Centipedegrass, and Zoysia Grass

Broadcast 2.2 to 4.4 pounds of this product per acre according to soil texture to control those weeds listed under Atrazine 90 WDG Herbicide Applied Alone-Corn or Grain Sorghum-Preplant Surface-Applied, Preplant Incorporated, or Preemergence.

Muck or peat	4.4 lbs product	Old beds: Within 2 days after lifting sod
		New beds: 3 to 4 days after sprigging or plugging
Sandy soil	2.2 lbs product	Old beds: Within 2 days after lifting sod
		New beds: 7 to 10 days after sprigging or plugging

If weeds regrow, apply an additional 2.2 pounds per acre on muck or peat, or 1.1 pounds per acre on sandy soil.

Use precautions: To avoid crop injury:

1. Do not apply within 30 days prior to cutting or lifting.
2. Do not apply in combination with surfactants or other spray additives.
3. Use only on turfgrass reasonably free of infestations of insects, nematodes, and diseases.
4. For muck or peat soils, do not apply more than 4.0 pounds active ingredient per acre for any application, do not apply more than 6.0 pounds active ingredient per year.
5. For sandy soil, do not apply more than 2.0 pounds active ingredient per acre for any application, do not apply more than 3.0 pounds active ingredient per year.
6. On newly sprigged turfgrass, temporary slowing of growth may follow application.

TURFGRASS at Residential sites (including homes, daycare facilities, schools, playgrounds, parks, recreational areas, and sports fields).

Bermudagrass, Centipedegrass, St. Augustinegrass, and Zoysia Grass

Apply Atrazine 90 WDG Herbicide after October 1 before emergence of winter annual weeds for control of annual blugrass*, burclover, carpet burweed, chickweed, corn speedwell, henbit, hop clover, and spurweed. Atrazine 90 WDG Herbicide will control annual bluegrass even if it is emerged at time of treatment. For control of summer annual weeds listed in the preemergence portion of the Atrazine 90 WDG Herbicide Applied Alone-Corn or Grain Sorghum section of this label, also apply Atrazine 90 WDG Herbicide in late winter before the weeds emerge. Apply in a minimum of 15.0 gallons of water per acre or 1.0 gallon per 1,000 square feet.

Where annual bluegrass is the major weed, use 1.1 pounds of Atrazine 90 WDG Herbicide per acre (0.4 ounce per 1,000 square feet). Use 1.1 pounds of this product (0.4 ounce per 1,000 square feet) for control of the other weeds named above. Do not exceed 1.1 pounds per acre per treatment on newly sprigged turfgrass or on hybrid bermudagrass such as Tiflawn, Tifway, and Ormond.

On newly sprigged turfgrass and hybrid bermudagrass, temporary slowing of growth and yellowing may occur following application. To avoid turf injury.

*(Local resistance may result in suppression only or lack of control. Refer to IMPORTANT NOTE within PRODUCT INFORMATION SECTION of this label.)

For continued summer annual weed control, apply another 1.1 pounds of this product per acre at least 30 days after the previous application, but not after April 15. However, do not make more than two applications of this product per year.

Use Restrictions:

- Do not apply more than 1.0 pound active ingredient per acre for (1.1 lb of this product) any application, do not apply more than 2.0 pounds active ingredient per acre per year.
- Applications made by backpack-spray to landscape turf- Restrict backpack application to landscape turf to spot treatments only.
- Use only on turfgrass reasonably free of infestations of insects, nematodes, and diseases.
- Do not use on golf greens.
- Do not use north of NC (except may be used in Virginia Coastal Plains) or west of the high rainfall areas of eastern OK and eastern TX.
- Do not apply over the rooting area of trees or ornamentals not listed on this label.
- Do not overseed with desirable turfgrass within 4 months before or 6 months after treatment.
- Do not apply this product to newly seeded bermudagrass until it has overwintered and has a well-developed rhizome system. Do not exceed 2.0 pounds product per acre within 12 months of seeding bermudagrass.

Note: Do not graze or feed turf clippings to animals, or illegal residues may result.

MACADAMIA NUTS

For preemergence control of many broadleaf and grass weeds, including crabgrass, foxtail*, wiregrass, flora's paintbrush, spanishneedles, and fireweed, broadcast 2.2 to 4.4 pounds of this product per acre before harvest and before weeds emerge. Repeat as necessary. Do not

spray when nuts are on ground during harvest period. Do not apply by air.

Restrictions:

- Do not apply more than 4.0 pounds active ingredient per acre for any application.
- Do not apply more than 8.0 pounds active ingredient per year.
- Do not apply with mechanically pressurized handguns to macadamia nuts.

*(Local resistance may result in suppression only or lack of control. Refer to IMPORTANT NOTE within PRODUCT INFORMATION SECTION of this label.)

GUAVA

Use only on established plantings which are at least 18 months old. Apply as a directed spray at 2.2 to 4.4 pounds per acre of Atrazine 90 WDG Herbicide in 20.0 to 50.0 gallons of spray mix preemergence or early postemergence to weeds. When applying postemergence, the use of a surfactant and greater spray volume (80.0 to 100 gallons of spray mix per acre) may enhance weed control. This product controls many annual broadleaf and grass weeds, including fireweed, purslane, scarlet pimpernel, spanishneedles, and sowthistle.

Restrictions:

- Do not allow spray to contact foliage or fruit.
- Do not apply more frequently than at 4 month intervals.
- Do not apply more than 4.0 pounds active ingredient per acre for any application.
- Do not apply more than 8.0 pounds active ingredient per year.
- Do not apply with mechanically pressurized handguns to guava.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in safe manner. Store in original container only. Store in cool, dry place. Reduce stacking height where local conditions, such as humidity or pallet overhang, can affect package strength. Personnel should use clothing and equipment consistent with good pesticide handling.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment by shaking and tapping sides and bottom to loosen clinging particles. Offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration. Do not burn, unless allowed by state and local ordinances.

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC – 1-800-424-9300.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with

such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS," AND LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. BOX 1286, GREELEY, CO 80632-1286.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

AAtrix, Concep, Dual MAGNUM, Dual II MAGNUM, Gramoxone Inteon and Nine-O are registered trademarks of a Syngenta Group Company.

Complex is a registered trademark of Kalo, Inc.

Unite is a trademark and Makaze is a registered trademark of Loveland Products, Inc.

Micro-Tech is a registered trademark of Monsanto Technology, LLC.

AAtrix Nine-O, Atrazine 90 WDG, Gramoxone Inteon, and Micro-Tech are restricted use pesticides.

PRIMEROS AUXILIOS	
En caso de ingesta:	<ul style="list-style-type: none"> • Llame de inmediato a un centro toxicológico o a un médico para que le recomienden un tratamiento. • Haga que la persona beba un vaso de agua a sorbos si puede tragar. • No induzca el vómito, a menos que así lo indique el centro toxicológico o un médico. • No administre nada por vía oral a una persona inconsciente.
En caso de inhalación:	<ul style="list-style-type: none"> • Traslade a la persona a un área con aire fresco. • Si la persona no respira, llame al 911 o a una ambulancia y, a continuación, administre respiración artificial, de preferencia boca a boca, si es posible. • Llame a un centro toxicológico o a un médico para que le recomienden un tratamiento adicional.
En caso de contacto con la piel o la ropa:	<ul style="list-style-type: none"> • Quítese la ropa contaminada. • Enjuague la piel de inmediato con bastante agua por 15 a 20 minutos. • Llame a un centro toxicológico o a un médico para que le recomienden un tratamiento.
En caso de contacto con los ojos:	<ul style="list-style-type: none"> • Mantenga los ojos abiertos y enjuáguelos lenta y cuidadosamente con agua, durante 15 a 20 minutos. • Si usa lentes de contacto, retírelos después de los primeros 5 minutos, luego continúe enjuagando los ojos. • Llame a un centro toxicológico o a un médico para que le recomienden un tratamiento.
ANTE UNA EMERGENCIA MÉDICA RELACIONADA CON ESTE PRODUCTO, LLAME AL: 1-800-424-9300. Lleve con usted el envase o la etiqueta del producto cuando llame a un centro toxicológico o a un médico, o cuando vaya a recibir tratamiento. Nota para el médico: No existe un antídoto específico para la atrazina. Si se ingiere este producto, induzca la emesis o el lavado de estómago. Se puede considerar el uso de una suspensión acuosa de carbón activado.	

INDICACIONES DE PRECAUCIÓN RIESGOS PARA SERES HUMANOS Y ANIMALES DOMÉSTICOS PRECAUCIÓN

Es nocivo si se ingiere. Evite respirar el polvo. Provoca irritación ocular. Evite el contacto con los ojos, la piel o la ropa.

Equipo de protección personal:

Los mezcladores, cargadores, limpiadores de equipos o derrames y otros manipuladores expuestos al concentrado deben usar:

- Overol (mameluco) encima de una camisa de manga larga y pantalones largos.
- Guantes resistentes a productos químicos hechos de laminado impermeable, goma de butilo ≥ 14 mils, goma de nitrilo ≥ 14 mils, goma de neopreno ≥ 14 mils, cloruro de polivinilo (PVC) ≥ 14 mils, Viton ≥ 14 mils.
- Medias y zapatos resistentes a sustancias químicas.
- Delantal resistente a productos químicos.
- Use como mínimo un respirador elastomérico de media máscara aprobado por NIOSH con cartuchos de vapor orgánico (VO) y filtros combinados R o P; O una máscara antigas aprobada por NIOSH con cartuchos de VO; o un respirador para suministro de aire aprobado por NIOSH equipado con cartuchos de VO en combinación con filtros HE.

Los aplicadores que usen equipos de pulverización en la espalda deben usar:

- Overol (mameluco) encima de una camisa de manga larga y pantalones largos.
- Guantes resistentes a productos químicos hechos de laminado impermeable, goma de butilo ≥ 14 mils, goma de nitrilo ≥ 14 mils, goma de neopreno ≥ 14 mils, cloruro de polivinilo (PVC) ≥ 14 mils, Viton ≥ 14 mils.
- Zapatos y medias resistentes a sustancias químicas.

Todos los demás aplicadores y manipuladores expuestos al producto diluido deben usar:

- Camisa de manga larga y pantalones largos.
- Zapatos con medias.
- Guantes resistentes a productos químicos hechos de laminado impermeable, goma de butilo ≥ 14 mils, goma de nitrilo ≥ 14 mils, goma de neopreno ≥ 14 mils, cloruro de polivinilo (PVC) ≥ 14 mils, Viton ≥ 14 mils.

Consulte los controles de ingeniería para requisitos adicionales.

Siga las instrucciones del fabricante para la limpieza y el mantenimiento de los equipos de protección personal (EPP). En caso de no existir dichas instrucciones de lavado, use detergente y agua caliente. Mantenga y lave el EPP separado del resto de la ropa para lavar. Deseche la ropa y otros materiales absorbentes que se hayan empapado o que estén muy contaminados con el concentrado de este producto. No vuelva a usarlos.

Indicaciones de controles de ingeniería:

Cuando los manipuladores usan sistemas cerrados, cabinas cerradas o aeronaves de manera tal que se cumplan los requisitos enumerados en la Norma de Protección del Trabajador (WPS, por sus siglas en inglés) para pesticidas agrícolas (40 CFR 170.240 (d) (4-6)), los requisitos del EPP del manipulador pueden reducirse o modificarse según lo especificado en la WPS.

RECOMENDACIONES DE SEGURIDAD PARA EL USUARIO

Los usuarios deben hacer lo siguiente:

- Lavarse las manos antes de comer, beber, mascar chicle, consumir tabaco o usar el baño.
- Quitarse la ropa o el EPP de inmediato si el pesticida se mete dentro. Luego, deberá lavarse por completo y ponerse ropa limpia.
- Quitarse el equipo de protección personal de inmediato después de manipular este producto. Lavar el exterior de los guantes antes de quitárselos. Tan pronto como sea posible, lavarse por completo y ponerse ropa limpia.

PELIGROS AMBIENTALES

La atrazina puede viajar (filtrarse o lixiviarse) a través del suelo y puede ingresar al agua subterránea que puede usarse como agua potable. Se encontró atrazina en las aguas subterráneas. Se recomienda a los usuarios no aplicar atrazina en suelos arenosos y franco-arenosos donde el nivel freático (aguas subterráneas) esté cerca de la superficie y donde estos suelos sean muy permeables, es decir, bien drenados. Los organismos agrícolas locales pueden brindar más información sobre el tipo de suelo en su zona y la ubicación de las aguas subterráneas.

INDICACIÓN DE ADVERTENCIA PARA ORGANISMOS NO OBJETIVO: Este producto es tóxico para las plantas y puede afectar de forma negativa el forraje y el hábitat de organismos

no objetivo, incluidos los polinizadores, en zonas adyacentes al sitio tratado. Proteja el forraje y el hábitat de los organismos no objetivo siguiendo las instrucciones de la etiqueta destinadas a minimizar la deriva de la pulverización.

El producto no debe mezclarse ni cargarse a menos de 15 metros (50 pies) de arroyos y ríos intermitentes, lagos y embalses naturales o presas. El producto no debe aplicarse a menos de 20 metros (66 pies) de los puntos donde la escorrentía de agua superficial del campo ingresa a arroyos y ríos perennes o intermitentes o a menos de 61 metros (200 pies) de lagos y embalses naturales o presas. Si este producto se aplica en tierras altamente erosionables, la franja de protección o separación de 20 metros (66 pies) desde los puntos de entrada de la escorrentía debe plantarse con cultivos, o sembrarse con césped u otro cultivo adecuado.

El producto no debe mezclarse, cargarse, ni usarse a menos de 15 metros (50 pies) de todos los pozos, incluidos los pozos abandonados, los pozos de drenaje y los sumideros. Se prohíben las operaciones que impliquen mezclar, cargar, enjuagar o lavar este producto en equipos o envases de manipulación o aplicación de pesticidas o desde ellos a menos de 15 metros (50 pies) de cualquier pozo, a menos que se realicen en una plataforma impermeable construida para soportar el peso de la carga más pesada que pueda colocarse o moverse por la plataforma. Dicha plataforma se diseñará y mantendrá para contener cualquier derrame de producto o fuga del equipo, el agua de enjuague o lavado del envase o equipo y el agua de lluvia que pueda caer sobre la plataforma.

No se permitirá que el agua superficial fluya por encima o desde la plataforma, lo que significa que la plataforma debe ser independiente. La plataforma se inclinará para facilitar la eliminación de material. Una plataforma no techada deberá tener capacidad suficiente para contener como mínimo el 110 % de la capacidad del envase del pesticida o equipo de aplicación más grande de la plataforma. Una plataforma que esté cubierta por un techo de tamaño suficiente para excluir por completo su contacto con la precipitación, tendrá una contención mínima del 100 % de la capacidad del envase del pesticida o equipo de aplicación más grande de la plataforma. Las capacidades de contención descritas más arriba deberán mantenerse en todo momento. Las capacidades mínimas de almacenamiento especificadas líneas arriba no se aplican a los vehículos que entregan pesticidas en los lugares de mezcla/carga. Deben cumplirse los requisitos adicionales impuestos por el estado en relación con la separación de las bocas del pozo y la contención de la zona de operaciones.

Se debe usar una de las siguientes restricciones en la aplicación de atrazina en campos con terrazas de baldosas que contengan tuberías verticales:

- No se debe aplicar a menos de 20 metros (66 pies) de distancia de los tubos verticales en campos con terrazas de baldosas.
- Aplique este producto a todo el campo con terrazas de baldosas e incorpórelo de inmediato a una profundidad de 5 a 7 centímetros (2 a 3 pulgadas) en todo el campo.
- Aplique este producto a todo el campo con terrazas de baldosas con una práctica de cultivo sin labranza solo cuando se realice el manejo de residuos de cultivos de alto volumen. El manejo de residuos de cultivos de alto volumen se describe como una práctica en la que se eliminan pocos o ningún residuo del campo durante y después de la cosecha.

Este pesticida es tóxico para los invertebrados acuáticos. No aplique el producto de forma directa al agua, en zonas donde habrá agua superficial ni en zonas intermareales que estén debajo de la marca de agua máxima promedio. No aplique el producto cuando las condiciones climáticas favorezcan la deriva de las zonas tratadas. La deriva y fuga de las zonas tratadas pueden ser peligrosos para los organismos acuáticos en zonas adyacentes. No contamine el agua al desechar el agua de lavado o enjuague del equipo.

El uso de este producto de forma contraria a lo indicado en su etiqueta constituye una infracción de la ley federal.

No aplique este producto de manera que entre en contacto con los trabajadores u otras personas, ya sea de forma directa o a causa de la deriva por el viento. Solo los manipuladores protegidos pueden permanecer en la zona durante la aplicación.

Para conocer cualquier requisito específico de su estado o tribu, consulte al organismo responsable de la normativa sobre pesticidas.

REQUISITOS PARA EL USO AGRÍCOLA

Use este producto solo de conformidad con su etiquetado y con la Norma de Protección del Trabajador, 40 CFR Parte 170. Esta norma contiene los requisitos para la protección de los trabajadores agrícolas en granjas, bosques, viveros e invernaderos, y para las personas que manipulan pesticidas agrícolas. Contiene los requisitos para la capacitación, descontaminación, notificación y asistencia de emergencia. Además, contiene las instrucciones específicas y excepciones relativas a las indicaciones de esta etiqueta sobre el equipo de protección personal (EPP) y el intervalo de ingreso restringido. Los requisitos en esta sección de la etiqueta (requisitos para uso agrícola) aplican únicamente a los usos de este producto que están cubiertos por la Norma de Protección del Trabajador.

No ingrese o permita el ingreso de trabajadores a las zonas tratadas durante el intervalo de ingreso restringido (IIR) de 12 horas.

Excepción: si el producto se inyecta en el suelo o se incorpora al suelo, la Norma de Protección del Trabajador, bajo ciertas circunstancias, permite que los trabajadores ingresen a la zona tratada si no habrá contacto con nada que haya sido tratado.

El EPP necesario para el ingreso anticipado a las zonas tratadas que se permite según la Norma de Protección del Trabajador y que implica el contacto con todo lo que ha sido tratado, como plantas, tierra o agua, es el siguiente:

- Overol (mameluco).
- Zapatos con medias.
- Guantes resistentes a productos químicos hechos de laminado impermeable, goma de butilo ≥ 14 mils, goma de nitrilo ≥ 14 mils, goma de neopreno ≥ 14 mils, cloruro de polivinilo (PVC) ≥ 14 mils, Viton ≥ 14 mils.

REQUISITOS PARA USOS NO AGRÍCOLAS

Los requisitos de esta sección se aplican a los usos de este producto que NO están dentro del ámbito de la Norma de Protección del Trabajador para pesticidas agrícolas (40 CFR Parte 170). La Norma de Protección del Trabajador se aplica cuando este producto se usa para cultivar plantas agrícolas en granjas, bosques, viveros o invernaderos.

No ingrese ni permita que otras personas ingresen hasta que la pulverización haya terminado.

ALMACENAMIENTO Y ELIMINACIÓN

No contamine el agua, los alimentos o forrajes al almacenar o desechar el producto.

ALMACENAMIENTO DE PESTICIDAS: almacénelo de manera segura. Almacénelo solo en el envase original. Conserve en un lugar fresco y seco. Reduzca la altura de apilamiento cuando las condiciones locales, como la humedad o el voladizo del palé, puedan afectar la resistencia del envase. El personal debe usar ropa y equipo acorde con la buena manipulación de pesticidas.

ELIMINACIÓN DE PESTICIDAS: los residuos resultantes del uso de este producto pueden eliminarse en el lugar o en una instalación de residuos aprobada.

ELIMINACIÓN DEL ENVASE: envase no rellenable. No vuelva a usar ni llenar este envase. Vacíe por completo la bolsa en el equipo de aplicación agitando y golpeando los lados y el fondo para aflojar las partículas adheridas. Entréguela para su reciclaje si existe esta opción o deseche la bolsa vacía en un vertedero sanitario o incinérela. No la queme, a menos que lo permitan las ordenanzas estatales y locales.

Para obtener ayuda en casos de derrame, fuga, incendio o exposición que involucren este material, comuníquese de día o de noche con CHEMTREC al 1-800-424-9300.