SUPPLEMENTAL LABELING

RESTRICTED USE PESTICIDE DUE TO ACUTE TOXICITY

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS – **NOT TO BE USED BY UNCERTIFIED PERSONS WORKING UNDER THE SUPERVISION OF A CERTIFIED APPLICATOR.**

Syngenta Crop Protection, LLC

P. O. Box 18300

Greensboro, North Carolina 27419-8300

SCP 1652A S14 0919

Gramoxone® SL 3.0 HERBICIDE

EPA Reg. No. 100-1652

A Weed, Grass, and Harvest Aid Desiccant/Defoliant Herbicide

Active Ingredients:

Total:	100.0%
Other Ingredients:	56.1%
Paraquat dichloride*	43.9%

*CAS No. 4685-14-7

Gramoxone SL 3.0 is formulated as a liquid containing 3.0 pounds paraquat cation per gallon as 4.134 pounds paraquat dichloride per gallon.

Gramoxone SL 3.0 contains alerting agent (odor), emetic, and dye.

KEEP OUT OF REACH OF CHILDREN / MANTENER FUERA DEL ALCANCE DE LOS NIÑOS.



Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.) See additional precautionary statements and directions for use inside booklet

All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Gramoxone SL 3.0 as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



DIRECTIONS FOR USE RESTRICTED USE PESTICIDE

Macadamia Nuts, HI Only

Crops (Including all cultivars, varieties, and/or hybrids of these)

Macadamia Nuts

Target Weed	Rate pt/A	Application Timing	Use Directions
Refer to Section 7.0.	1.3	Postemergence (Directed Spray)	For use in Hawaii Only to control weeds while macadamia nuts are on the ground. Apply as a ground directed spray application in a
			minimum of 30-150 gal/A of spray volume when weeds and grasses are 1-6" high.
			Add Spreader (nonionic) at 8 to 32 fl oz per 100 gallons.
			Treatments should be made immediately after harvest to minimize the number of nuts on the ground at the time of application.
			For mature woody weeds, green suckers, late germinating weeds and grasses and for perennials – retreatment or spot treatment may be necessary.
			See Section 4.4.4 of the Gramoxone SL 3.0 label for adjuvant recommendations.

Tank Mix Options:

 Gramoxone SL 3.0 can be tank mixed with other herbicides labeled for use on macadamia nuts applied by the same methods and at the same timings. Refer to **Section 4.4.2** for tank-mix precautions in the Gramoxone SL 3.0 label.

Resistance Management:

Refer to Section 3.1.

Precautions:

- Do not allow spray to contact green stems or foliage, as injury may result.
- Do not spray under windy conditions and use a shield for young trees.

USE RESTRICTIONS

- 1. Refer to **Section 6.1** for additional product use restrictions.
- 2. Maximum Single Application Rate: 1.3 pt/A
- 3. **REI:** 24 hours
- 4. Minimum Application Interval: 7 days
- 5. Maximum Annual Rate: 5.2 pt/A
 - a. **DO NOT** exceed 1.95 lb ai/A/year of paraquat-containing products.
- 6. **DO NOT** make more than 4 application per year.
- 7. **DO NOT** apply this product through any type of irrigation system.
- 8. **DO NOT** allow animals to graze on treated areas.
- 9. Pre-Harvest Interval (PHI): 1 day

Gramoxone® and the SYNGENTA Logo are Trademarks of a Syngenta Group Company.

©2019 Syngenta

For non-emergency (e.g., current product information), call Syngenta Crop Protection at 1-800-334-9481.

Manufactured for: Syngenta Crop Protection, LLC P.O. Box 18300 Greensboro, North Carolina, 27419-8300