

FIFRA Section 24(c) Special Local Need Label

METHOXYFENOZIDE	GROUP 18	INSECTICIDE
SPINETORAM	GROUP 5	INSECTICIDE



Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Intrepid Edge

EPA Reg. No. 62719-666

EPA SLN No. TN-230001

For Distribution and Use Only in the State of Tennessee

Control of Thrips in Cotton

Expiration date: This label expires and must not be distributed or used in accordance with this SLN registration after April 4, 2028.

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read this SLN and the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this FIFRA 24(c) Special Local Need labeling is subject to all use precautions and limitations imposed by the label affixed to the container for this product.

Directions for Use

Refer to the Intrepid Edge® product label for Application Directions, Mixing Directions, and Restrictions.

Application Rate: The amount of Intrepid Edge® applied per acre will depend upon pest infestation level. Use a lower rate in the rate range for light infestations and a higher rate in the rate range for heavy infestations. Use of an adjuvant may increase the level of thrips control.

Application Timing: Make applications during late cotyledon stage cotton and then continue to manage populations until cotton is no longer susceptible to early season thrips damage. Under heavy pressure multiple applications may be needed to control pest. Using Intrepid Edge® is not a replacement for insecticide seed treatments or in-furrow insecticide applications at planting. Early season, foliar applications of Intrepid Edge® to suppress thrips in cotton should only be made to fields that have already received an insecticide seed treatment or in-furrow insecticide application for thrips control.

Resistance Management: Do not make more than two consecutive applications of Group 5 or 18 insecticides per crop. If additional treatments are required after two consecutive applications of Group 5 or 18 insecticides, rotate to another class of effective insecticide for at least one application. Consult your Corteva Agriscience representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area. Avoid applying Intrepid Edge and other Group 5 and 18 insecticides to consecutive generations of all target insect pests.

Pest	Intrepid Edge® (fl oz/acre)
Thrips	3 – 6

Restrictions:

- **Preharvest Interval:** Do not apply within 28 days of harvest.
- Do not apply more than 8 fl oz of Intrepid Edge® per acre per application.
- Do not apply more than a total of 48 fl oz of Intrepid Edge® (0.94 lb ai methoxyfenozide, 0.19 lb ai spinetoram) per acre per year.
- **Minimum Treatment Interval:** Do not make applications less than 4 days apart.
- **Maximum Number of Applications:** Do not make more than 6 applications per year.
- Do not make more than two consecutive applications.

™®Trademarks of Corteva Agriscience and its affiliated companies

R911-019

Approved: 04/04/2023

Initial printing