

FIFRA SECTION §24(C) SPECIAL LOCAL NEED LABEL

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON

THIS LABEL IS VALID UNTIL DECEMBER 31, 2023 UNLESS OTHERWISE AMENDED, WITHDRAWN, CANCELED, OR SUSPENDED

EPTAM® 7F

Selective Herbicide - Emulsifiable Liquid

EPA Reg. No. 10163-283, EPA SLN No. OR-160009

FOR CONTROL OF ITALIAN RYEGRASS, ANNUAL BLUEGRASS, COMMON CHICKWEED AND HENBIT IN TABLE BEETS GROWN FOR SEED AND SWISS CHARD GROWN FOR SEED.

ACTIVE INGREDIENT:

Total 100.0%

Contains 7 pounds active ingredient per gallon. Contains petroleum distillates

KEEP OUT OF REACH OF CHILDREN WARNING-AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

ENVIRONMENTAL HAZARDS

This chemical is toxic to mammals. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Follow all applicable directions, restrictions, and precautions, including statements pertaining to the Worker Protection Standards, on the EPA-registered main label and this label.
- This label must be in the possession of the user at the time of pesticide application.

CROP	PINTS/ACRE	COMMENTS
SWISS CHARD GROWN FOR SEED	2 1/4 - 3 1/2	PREPLANT INCORPORATION: Apply and incorporate 2 ¼ pints per acre EPTAM 7E on course textured soils, or 3 ½ pints per acre on medium and fine textured soils just before planting. Crop injury and stunting may occur if conditions for germination and growth are not optimum (e.g. prolonged periods of cold wet weather or lack of moisture during germination and early growth).
		POSTEMERGENCE INCORPORATION: (After the First True Leaves Have Formed) Apply 3 ½ pints EPTAM 7E per acre after thinning and clean cultivation and incorporate to a depth of 2 to 3 inches. Do not exceed 3 ½ pints EPTAM 7E per acre per crop (except for irrigation applications in the Pacific Northwest Region where 2 applications of 3 ½ pints may be made).
		LAY-BY SUBSURFACE APPLICATION: Prior to application, a clean cultivation must be made for all existing weed growth to be destroyed. Apply 3 ½ pints EPTAM 7E per broadcast acre or in a band treatment (using 2 shanks per row 5 ½ inches apart, centered on the drill row with rows 26 to 42 inches apart) use 1 ¾ pints per acre.
TABLE BEETS GROWN FOR SEED		POSTEMERGENCE INCORPORATION: (After the First True Leaves Have Formed) Apply 3 ½ pints EPTAM 7E per acre after thinning and clean cultivation and incorporate to a depth of 2 to 3 inches. Do not exceed 3 ½ pints EPTAM 7E per acre per crop (except for irrigation applications in the Pacific Northwest Region where 2 applications of 3 ½ pints may be made).
		LAY-BY SUBSURFACE APPLICATION: Prior to application, a clean cultivation must be made for all existing weed growth to be destroyed. Apply 3 ½ pints EPTAM 7E per broadcast acre or in a band treatment (using 2 shanks per row 5 ½ inches apart, centered on the drill row with rows 26 to 42 inches apart) use 1 ¾ pints per acre.

Special Crop use Restrictions: The pesticide applicator, the producer of the crop, and the seed conditioner must be aware that use of this product according to this labeling is deemed a non-feed/non-food use by the Oregon Department of Agriculture, and is regulated by the Oregon Administrative Rule (OAR) 603-057-0535, Pesticide Use On Crops Grown For Seed. If the applicator of this pesticide is not the producer, the applicator must provide a copy of this labeling to the producer of the crop. Producers of this crop who use this product, or cause the product to be used on a field they operate, must provide a copy of this pesticide label to the seed conditioner.

This pesticide does not have an established pesticide residue tolerance for this crop. Consequently, no portion of this seed may be used or distributed for food or feed for 1 year (365 days) after the last application of this product. This restriction pertains to, but is not limited to, green chop, forage, hay, pellets, meal, whole seed, cracked seed, straw, roots, bulbs, foliage or seed screenings, and to the grazing of the crop field, stubble or regrowth. All seed screenings shall be disposed of in such a manner that screenings cannot be distributed or used for food or feed purposes, as indicated in OAR 603-057-0535. Additional regulations concerning seed screenings are stated in OAR 603-057-0535.

Any seed from a field treated with this pesticide product shall bear specific and conspicuous container labeling, or if shipped in bulk, on the shipment invoice or bill of lading. The labeling shall contain the following statement:

"This seed was produced using one or more products for which the United States Environmental Protection Agency has not established pesticide residue tolerances. This seed, in whole, as sprouts, or in any form may not be used for human consumption or animal feed. Failure to comply with this condition may violate requirements of the Federal Food and Drug Administration, the Oregon Department of Agriculture and other regulatory agencies.

24(c) Registrant: Gowan Company P.O. Box 5569 Yuma, AZ 85366 Updated 2-27-18